

RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



COMMUNICATION

Potential Responses to Neutral Thiophenols of Polymeric Membrane Electrodes and Their Applications in Potentiometric Biosensing

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Long Li^{a, b} and Wei Qin^{* a}

www.rsc.org/

The unexpected potential responses to electrically neutral thiophenols (ArSHs) of anion-exchanger doped polymeric membranes are described. Based on the dimerization reactions of ArSHs via horseradish peroxidase (HRP)-catalyzed oxidations, a sensitive and facile potentiometric biosensing platform for HRP has been developed.

For many decades, polymeric membrane ion-selective electrodes have been considered to be feasible only for ionic analytes because a charged species can form a charge separation layer at the interface between a membrane phase and an aqueous phase.¹ This has significantly limited the development and applications of potentiometric membrane electrodes for detection of electrically neutral species and sensing of their related reactions. In recent years, the potential response behaviors and mechanisms of several kinds of electrically neutral species such as phenols (ArOHs)² and boronic acids (BAs)³ on the ion exchanger-doped polymeric membranes have been reported. It has been found that phenols and boronic acids can form the complexes with quaternary ammonium salts via hydrogen bonding in the membrane phase, which could undergo proton dissociation with the concomitant ejection of the formed acids from the membrane phase into the aqueous phase.^{2,3} Based on the oxidations of monomeric phenols to oligomeric phenols and reversible covalent bindings between boronic acid groups and diols, potentiometric sensing platforms for horseradish peroxidase (HRP) mimetics⁴ and for saccharides^{3,5} have been developed, respectively. However, electrically neutral species that can induce potential responses on the polymeric membrane electrodes are rather limited.^{2,3} More examples are still highly desired because they are favourable to ascertain the

generality of this phenomenon and the corresponding response mechanism. Moreover, more electrically neutral species that can induce potential responses on the polymeric membrane electrodes will largely broaden the applications of potentiometric sensors and stimulate the research on electrical chemistry at the interface of two immiscible phases.

Thiols are an important class of molecules in chemical industry and biological systems.⁶ Thiophenols (ArSHs) have been widely used in the production of agrochemicals, polymers and pharmaceuticals.⁷ Aliphatic thiols such as cysteine, homocysteine and glutathione involved in many physiological processes play important roles in living organisms.⁷ ArSHs are more toxic than aliphatic thiols, and it is essential to selectively differentiate ArSHs from aliphatic thiols.^{6,8} Based on the stronger nucleophilicities of the corresponding thiolates from ArSHs, selective fluorescent probes for ArSHs can be developed.^{8,9} However, those fluorescent probes are usually structurally complicated, and the synthesis procedures are time consuming.

In this work, the unexpected potential responses to electrically neutral ArSHs of polymeric membranes doped with lipophilic anion exchangers are shown. The complexations between ArSHs and quaternary ammonium salts have been studied in homogeneous organic solutions. Efforts have been made to relate the response behaviors of ArSHs to their lipophilicities and the association constants between ArSHs and tetraalkylammonium chloride. The heterogeneous two phase extraction experiments indicate the dissociation behaviors of ArSHs in the organic phase and confirm the response mechanism. The potential responses to ArSHs of the polymeric membranes provide a simple way to differentiate ArSHs from biologically important bioactive aliphatic thiols. By using ArSHs as reducing substrates, a potentiometric sensing platform for oxidases and their related reactions can be developed.

Polymeric liquid membranes containing PVC and o-NPOE in a weight ratio of 1:1 and the receptor salt (10 mM/kg) were prepared by the solvent-casting technique with tetrahydrofuran as the casting solvent. After transferring the cocktail into a glass ring fixed on a glass plate and letting tetrahydrofuran evaporate overnight, a uniform membrane of

^a Key Laboratory of Coastal Environmental Processes and Ecological Remediation, Yantai Institute of Coastal Zone Research (YIC), Chinese Academy of Sciences (CAS); Shandong Provincial Key Laboratory of Coastal Environmental Processes, YIC-CAS, Yantai, Shandong 264003, P. R. China. E-mail: wqin@yic.ac.cn; Fax: +86 535 2109000; Tel: +86 535 2109156

^b University of the Chinese Academy of Sciences, Beijing 100049, P. R. China.

†Electronic Supplementary Information (ESI) available. See DOI: 10.1039/x0xx00000x




Fig. 1 Potential responses of five ArSHs in a 50 mM acetic acid buffer of pH 4.0 at the o-NPOE-PVC-TDDA⁺Cl⁻ membrane electrode. All the pKa and log P values were calculated by ACD/Labs 12. All potential values were measured using an electrochemical workstation under zero current conditions.

200 μm thickness was obtained. Disks of 5-mm diameter were punched from the parent membrane and glued to plasticized PVC tubes (i.d. 3 mm, o.d. 5 mm) to fabricate the polymeric membrane electrodes. Fig. 1 shows the potential responses to a number of ArSHs of a TDMA⁺Cl⁻-doped polymeric membrane. At pH 4.0, most of the ArSHs exist in their undissociated (unionized) forms. The potential differences in the concentration range of 10⁻⁴-10⁻³ M were -90 and -140 mv for -Cl and -NO₂ substituted ArSHs, respectively, which are larger than the theoretical value for a singly charged anion. The anionic potential responses induced by the electrically neutral ArSHs are unexpected, since neutral species cannot perturb the distribution of charged species directly at the interface between the immiscible phases.¹⁰ It has been found that phenols and boronic acids can undergo proton dissociation in the organic phase with concomitant ejection of HX to the aqueous phase, thus inducing the anionic responses at the sample-membrane interface.^{2,3}

To investigate the response mechanism of the membrane electrode for ArSHs, efforts were made to understand the interactions between ArSHs and quaternary ammonium salts in homogeneous organic solutions. The association constant (K_a) for the 1:1 complexes between ArSHs and TDMA⁺Cl⁻ were measured in chloroform solutions using UV titration experiments (Fig.S1, ESI⁺). As shown in Tab. 1, the stabilities of the complexes can be determined by the acidities (pK_a) of the ArSH molecules, and the calculated data correlate well with the corresponding Hammett constants for the substituents (Fig. S2, ESI⁺).¹² It should also be noted that the K_a values for the complexes between ArSHs and TDMA⁺Cl⁻ are significantly larger than those for ArOHs, probably due to the higher acidities of ArSHs. Studies on the complexations between ArOHs and quaternary ammonium salts (Q⁺X⁻) show that the K_a values are X⁻ dependent but Q⁺ independent.¹¹ It can be assumed that the interaction between an undissociated -SH group and the anionic component X⁻ of Q⁺X⁻ may be the main

Tab. 1 Association constants (K_a) for ArSHs or ArOHs, acid dissociation constants (pK_a) for ArSHs and Hammett constants (H) for different substituents.


ArSH	K _a (M ⁻¹)	pK _a ^c	H ^d
<i>p</i> -nitro(thio)phenol	^a 4.3 × 10 ⁵ / ^b 1.3 × 10 ⁵	4.5	0.78
<i>p</i> -fluorothiophenol	2.4 × 10 ⁵	6.2	0.06
<i>p</i> -chlorothiophenol	1.5 × 10 ⁵	6.1	0.23
<i>p</i> -toluenethiol	1.3 × 10 ⁵	6.8	-0.17
<i>p</i> -methox(thio)phenol	^a 1.7 × 10 ⁵ / ^b 7.9 × 10 ³	6.8	-0.27

^aThe association constants of the 1:1 complexes for ArSHs with TDMA⁺Cl⁻ were determined at room temperature (ca. 25 °C) by UV titration experiments. ^bAssociation constants for ArOHs and TDMA⁺Cl⁻ from ref 2. ^cAcid dissociation constant calculated by ACD/Labs 12. ^dHammett constants from ref 12.

association form for the complexation between Q⁺X⁻ and an ArSHs compound. This is reasonable since the dissociated ArS⁻ group could hardly complex with the bulky Q⁺ group to form an ion pair, let alone the thermodynamically unfavourable product H⁺X⁻.¹³

Extraction experiments in two phase systems were carried out with aqueous solutions containing *p*-chlorothiophenol and chloroform solutions with or without TDMA⁺Cl⁻. UV spectra were measured to study the extraction behaviors of ArSHs and their deprotonation processes. As shown in Fig.S3 (ESI⁺), when *p*-chlorothiophenol in water (at pH 4) is extracted into pure CHCl₃, the UV spectrum of the CHCl₃ phase shows a peak at 250 nm, which originates from the undissociated neutral *p*-chlorothiophenol. However, when CHCl₃ with TDMA⁺Cl⁻ is used to extract *p*-chlorothiophenol, the peak at 250 nm decreases along with an increase at 275 nm, indicating the complexation occurs between *p*-chlorothiophenol and TDMA⁺Cl⁻. Experiments also show that the pH of the aqueous phase decreases from 4.0 to 3.5 while the concentration of chloride increases after extraction of *p*-chlorothiophenol from the aqueous into the CHCl₃ phase containing TDDA⁺Cl⁻ (see Tab. S1, ESI⁺). According to the pH change, it can be estimated that ca. 40% *p*-nitrothiophenol extracted into the organic phase dissociates. The extraction experiments confirm the dissociation of ArSHs in the organic phase and the ejection of H⁺X⁻ into the aqueous phase, which causes the deportation process thermodynamically favourable and induces the anionic potential responses. The above results indicate that the response mechanism for ArSHs is similar to those for ArOHs and BAs (Scheme S1, ESI⁺). In these processes, BAs behavior as Lewis acids, while ArSH and ArOHs work as Bronsted acids.

In consistency with the potential responses to ArOHs and BAs of the polymeric membranes doped with lipophilic quaternary ammonium salts, the sensitivity of the ArSH membrane electrode is dependent on the acidity (pK_a) and lipophilicity (log P) of each ArSH. As shown in Fig. 1, the magnitudes of the potentiometric responses to the ArSHs are in the order of *p*-NO₂ > *p*-Cl > *p*-CH₃ > *p*-F > *p*-OCH₃. Indeed, a



Scheme 1 Response mechanism of the proposed potentiometric platform.

thiophenol derivative with a stronger acidity and higher Log P induces a larger anionic response. The response order of p -NO₂ > p -Cl > p -CH₃ is induced by the acidity effect, and the order of p -Cl > p -CH₃ > p -F > p -OCH₃ is due to the lipophilicity effect. Similar potentiometric selectivities were also reported by Umezawa.² It was found that alkythiols could not induce potential responses on the polymeric membrane (data not shown), probably due to their weak acidities and low lipophilicities. Therefore, it can be concluded that the aromatic thiol moiety is crucial to the potential responses. Considering that alkythiols cannot induce potential response on the membrane electrodes, polymeric membrane electrode may be used as a simple and efficient sensor for distinguishing ArSHs from alkythiols.

The responses of the membranes doped with different quaternary ammonium cations (i.e., TDMA⁺ and TBuA⁺) and anions with (i.e., TCPB⁻, SCN⁻ and Cl⁻) were investigated. The results are shown in (Fig.S4, ESI[†]). In contrast to negligible responses of the blank and the hydrophilic TBuA⁺ doped membranes, all membranes containing lipophilic TDMA⁺ show anionic responses, which are dependent on the hydrophilicities and H-bond basicities of the anions (i.e., Cl⁻ > SCN⁻ > TCPB⁻). The order of the magnitude of potentiometric response is parallel to that of the charge-dipole and/or hydrogen bonding interactions between X⁻ and ArOH as expected from the ion radiuses of X⁻ (1.8 and 2.08 Å for Cl⁻ and SCN⁻, respectively²). It is clear that lipophilic cations dominate and the anions regulate the potential responses.

ArSHs can be used as oxidase substrates due to their high reducibilities. The oxidations of ArSHs induce S-S coupling and no thiol groups remain in the oxidation products. Therefore, these oxidation processes could “turn off” the anionic potential responses of the reporter thiol groups and the negative signals would return to the positive ones. The potentiometric platform for oxidase/mimic enzymes and their related reactions can be developed as shown in Scheme 1. As a proof of concept experiment, the potential traces to 4 kinds of ArSHs at the polymeric membrane electrode before and after the oxidation reactions are shown in Fig.S5 (ESI[†]). It can be seen that anionic potential responses can be obtained after ArSHs are added into the solutions, and the positive responses are indeed observed after H₂O₂ and horseradish peroxidase (HRP) are added to induce the oxidation reactions of the ArSHs. These results confirm that the formed S-S coupling products c-




Fig. 2 Potentiometric responses to 10⁻³ M p -nitrothiophenol of polymeric membrane electrodes in acetic acid buffer (pH=4.0) in the absence (a) or presence (b) of 10 mM H₂O₂ and different concentrations of HRP. All potential values were measured using an electrochemical workstation under zero current conditions.

annot induce potential responses at the polymeric membrane electrodes. Our previous studies indicate that the potentiometric sensing of HRP/mimic enzymes can be done via the oligomerization reactions of ArOHs based on the large lipophilicities of the formed oligomeric ArOHs.⁴ Unlike those of ArOHs which involve C-O and C-C coupling reactions, the oxidations of ArSHs produce the S-S coupling products. For the ArOH based platform using the “negative to negative” signal mode, the initial negative potential response could be a large background for the following detection, so that the substrate should be carefully selected to insure not only a rapid reaction kinetic but also a large Log P change. However, for the ArSHs based method using the “negative to positive” signal mode, only a fast reaction kinetic is required since the S-S coupling product without the thiol moiety cannot induce any potential response on the polymeric membrane. As shown in Figure S3, the large potential change before and after oxidation reactions indicates that p -nitrothiophenol is the most suitable substrate in these 4 kinds of ArSHs for HRP detection.

As a preliminary application, the potentiometric platform was used to detect HRP in the absence and presence of H₂O₂ (Fig.2). It has been reported that thiols can be oxidized to

disulfides via the thiol-mediated autocatalytic generation of H_2O_2 from oxygen.¹⁴ As shown in Fig.2(a) and Fig.S6(a) (ESI⁺), HRP can be detected in the range of $1 \times 10^{-3} \sim 1 \times 10^{-2}$ U/mL with a detection limit of 4×10^{-4} U/mL in the absence of H_2O_2 . Such sensitivity is comparable to traditional colorimetric method for HRP detection. It has been shown that H_2O_2 assists the oxidization of thiols and thus can improve the sensitivity for HRP detection.¹⁴ Indeed, the presence 10 mM H_2O_2 allows HRP to be detected in the range of $1 \times 10^{-8} \sim 2 \times 10^{-7}$ U/mL with a detection limit of 5×10^{-9} U/mL (Fig.2(b) and Fig.S6(b) (ESI⁺)). The detection limit is 1.5 orders of magnitudes lower than that of the method based on the oxidation of *o*-phenylenediamine.¹⁵ Besides HRP detection, this versatile platform can also be used for other peroxidases and peroxidase mimics (e.g., hemin/G-quadruplexes) and their related biosensing applications.

Conclusions

In conclusion, we report here for the first time the unexpected anionic potentiometric responses to electrically neutral ArSHs of the polymeric membranes doped with lipophilic anion exchangers. The complexations between ArSHs and quaternary ammonium salts were studied in homogeneous organic solutions. The association constants between the ArSHs and TDMA^+Cl^- obtained from titration experiments correlate with the Hammett constants for the para substituents and with the response magnitudes of the ArSHs. The heterogeneous two phase extraction experiments confirm the dissociation of ArSHs in the organic phase and the ejection of H^+X^- into the aqueous phase. The response mechanism for ArSHs is similar to those of ArOHs and boronic acids. The anion exchanger doped polymeric membranes can be readily used to distinguish ArSHs from aliphatic thiols. Based on the HRP-catalyzed oxidations of ArSHs, a sensitive and facile potentiometric platform for HRP has been developed, which can be extended for applications in peroxidase/peroxidase mimetic related biosensing.

This work was financially supported by the National Natural Science Foundation of China (21475148) and the Taishan Scholar Program of Shandong Province.

Notes and references


- 1 E. Bakker, P. Buhlmann and E. Pretsch, *Chem. Rev.*, 1997, **97**, 3083-3132; (b) J. Bobacka, A. Ivaska and A. Lewenstam, *Chem. Rev.*, 2008, **108**, 329-351; (c) P. Buhlmann and L. D. Chen, Ion-Selective Electrodes with Ionophore-Doped Sensing Membranes. In *Supramolecular Chemistry: From Molecules to Nanomaterials*: Wiley, 2012.
- 2 T. Ito, H. Radecka, K. Tohda, K. Odashima and Y. Umezawa, *J. Am. Chem. Soc.*, 1998, **120**, 3049-3059.
- 3 X. W. Wang, D. F. Yue, E. G. Lv, L. Wu and W. Qin, *Anal. Chem.*, 2014, **86**, 1927-1931.
- 4 X. W. Wang, Z. F. Ding, Q. W. Ren and W. Qin, *Anal. Chem.*, 2013, **85**, 1945-1950.
- 5 H. D. Chen, L. Li, H. M. Guo, X. W. Wang and W. Qin, *Rsc Adv.*, 2015, **5**, 13805-13808.
- 6 X. M. Shao, R. X. Kang, Y. L. Zhang, Z. T. Huang, F. F. Peng, J. Zhang, Y. Wang, F. C. Pan, W. J. Zhang and W. L. Zhao, *Anal. Chem.*, 2015, **87**, 399-405.
- 7 (a) J. C. Love, L. A. Estroff, J. K. Kriebel, R. G. Nuzzo and G. M. Whitesides, *Chem. Rev.*, 2005, **105**, 1103-1169; (b) J. Yin, Y. Kwon, D. Kim, D. Lee, G. Kim, Y. Hu, J.-H. Ryu and J. Yoon, *J. Am. Chem. Soc.*, 2014, **136**, 5351-5358; (c) H. S. Jung, X. Chen, J. S. Kim and J. Yoon, *Chem. Soc. Rev.*, 2013, **42**, 6019-6031; (d) L. A. Herzenberg, S. C. DeRosa, J. G. Dubs, M. Roederer, M. T. Anderson, S. W. Ela, S. C. Deresinski and L. A. Herzenberg, *P. Natl. Acad. Sci. USA*, 1997, **94**, 1967-1972.
- 8 J. Li, C.-F. Zhang, S.-H. Yang, W.-C. Yang and G.-F. Yang, *Anal. Chem.*, 2014, **86**, 3037-3042.
- 9 (a) H. Maeda, M. Matsuno, M. Ushida, K. Katayama, K. Saeki and N. Itoh, *Angew. Chem., Int. Ed.*, 2005, **44**, 2922-2925; (b) A. Taglieber, H. Höbenreich, J. D. Carballeira, R. J. G. Mondière, M. T. Reetz, *Angew. Chem., Int. Ed.*, 2007, **46**, 8597-8600.
- 10 P. Buhlmann, M. Hayakawa, T. Ohshiro, S. Amemiya and Y. Umezawa, *Anal. Chem.*, 2001, **73**, 3199-3205.
- 11 K. Odashima, T. Ito, K. Tohda and Y. Umezawa, *Chem. Pharm. Bull.*, 1998, **46**, 1248-1253.
- 12 C. Hansch, A. Leo and A. W. Taft, *Chem. Rev.*, 1991, **91**, 165-195.
- 13 A. J. Fry, *J. Org. Chem.*, 2013, **78**, 5476-5481.
- 14 E. Golub, R. Freeman and I. Willner, *Anal. Chem.*, 2013, **85**, 12126-12133.
- 15 X. W. Wang and W. Qin, *Chem. Eur. J.*, 2013, **19**, 9979-9986.

Potential Responses to Neutral Thiophenols of Polymeric Membrane Electrodes and Their Applications in Potentiometric Biosensing

Long Li^{a,b} and Wei Qin^{*a}

^aKey Laboratory of Coastal Environmental Processes and Ecological Remediation, Yantai Institute of Coastal Zone Re-search (YIC), Chinese Academy of Sciences (CAS); Shandong Provincial Key Laboratory of Coastal Environmental Pro-cesses, YICCAS, Yantai, Shandong 264003, P. R. China. E-mail: wqin@yic.ac.cn; Fax: +86 535 2109000; Tel: +86 535 2109156

^bUniversity of the Chinese Academy of Sciences, Beijing 100049, P. R. China.



The unexpected potential responses to electrically neutral thiophenols of anion-exchanger doped polymeric membranes have been observed and applied in potentiometric biosensing of horseradish peroxidase.