



Cite this: *Environ. Sci.: Processes Impacts*, 2020, 22, 1355

Interactions of ferrous iron with clay mineral surfaces during sorption and subsequent oxidation†

Natacha Van Groeningen, ^a Laurel K. ThomasArrigo, ^a James M. Byrne, ^b Andreas Kappler, ^b Iso Christl *^a and Ruben Kretzschmar ^a

In submerged soils and sediments, clay minerals are often exposed to anoxic waters containing ferrous iron (Fe^{2+}). Here, we investigated the sorption of Fe^{2+} onto a synthetic montmorillonite (Syn-1) low in structural Fe (<0.05 mmol Fe per kg) under anoxic conditions and the effects of subsequent oxidation. Samples were prepared at two Fe-loadings (0.05 and 0.5 mol Fe added per kg clay) and equilibrated for 1 and 30 days under anoxic conditions ($\text{O}_2 < 0.1$ ppm), followed by exposure to ambient air. Iron solid-phase speciation and mineral identity was analysed by ^{57}Fe Mössbauer spectroscopy and synchrotron X-ray absorption spectroscopy (XAS). Mössbauer analyses showed that $\text{Fe}(\text{II})$ was partially oxidized (14–100% of total added Fe^{2+}) upon sorption to Syn-1 under anoxic conditions. XAS results revealed that the added Fe^{2+} mainly formed precipitates (layered Fe minerals, $\text{Fe}(\text{III})$ -bearing clay minerals, ferrihydrite, and lepidocrocite) in different quantities depending on the Fe-loading. Exposing the suspensions to ambient air resulted in rapid and complete oxidation of sorbed $\text{Fe}(\text{II})$ and the formation of $\text{Fe}(\text{III})$ -phases ($\text{Fe}(\text{III})$ -bearing clay minerals, ferrihydrite, and lepidocrocite), demonstrating that the clay minerals were unable to protect ferrous Fe from oxidation, even when equilibrated 30 days under anoxic conditions prior to oxidation. Our findings clarify the role of clay minerals in the formation and stability of Fe-bearing solid phases during redox cycles in periodically anoxic environments.

Received 4th February 2020
Accepted 14th April 2020

DOI: 10.1039/d0em00063a

rsc.li/espi

Environmental significance

Clay minerals are considered major sorbents in soils and sediments. However, their significance for the redox cycling of iron is still unclear. Under anoxic conditions, elevated concentrations of dissolved ferrous iron are observed. We demonstrate that at circumneutral to alkaline pH, surface-mediated oxidation of ferrous iron can occur under anoxic conditions in the presence of clays that are virtually free of structural iron. The retention of dissolved ferrous iron in the presence of clay minerals was evidenced to be mainly due to the formation of secondary $\text{Fe}(\text{III})$ phases. Subsequent aeration led to the complete oxidation of iron. In contrast to surface precipitation, adsorption of dissolved iron by layered aluminosilicates plays a minor role in iron retention and does not protect ferrous iron from oxidation.

Introduction

Iron is ubiquitous in soils and sediments and of great importance in the biogeochemical cycling of metals.^{1–5} Its solubility in well-aerated soils is mainly controlled by the dissolution and

precipitation of $\text{Fe}(\text{III})$ -(oxyhydr)oxides.⁶ During periods with anoxic conditions, $\text{Fe}(\text{III})$ -(oxyhydr)oxides serve as electron acceptors for dissimilatory metal-reducing bacteria leading to the release of aqueous Fe^{2+} . Ferrous iron (Fe^{2+}), being one of the most abundant redox-active species in anoxic soils, has a crucial role in controlling macronutrient and trace element cycles.⁷ The retention of Fe^{2+} in soils and sediments is, as for most divalent cations, governed by sorption processes (including adsorption as well as surface precipitation and polymerization)⁸ to minerals with a large surface area and specific sorption properties such as clay minerals, carbonates, and $\text{Fe}(\text{III})$ - and $\text{Mn}(\text{IV})$ -(oxyhydr)oxides. Under anoxic conditions, clay minerals are one of the most important sorbent phases as they are commonly found in high abundance in soils but are not susceptible to reductive dissolution. Aqueous Fe^{2+} can adsorb to clay mineral surfaces *via* cation exchange on permanently charged face

^aInstitute of Biogeochemistry and Pollutant Dynamics, Department of Environmental Systems Science, CHN, ETH Zürich, 8092 Zürich, Switzerland. E-mail: iso.christl@env.ethz.ch

^bGeomicrobiology Group, Centre for Applied Geosciences (ZAG), University of Tübingen, Hölderlinstrasse 12, D-72074, Tübingen, Germany

† Electronic supplementary information (ESI) available. Calculation of saturation indices, protocol to ensure anoxic conditions, measured pH after equilibration, percentage of Fe sorbed by oxidation of dissolved ferrous iron, concentration analyses, concentration of Si release analysis, details on XAS data collection and reduction, additional Mössbauer spectra and XAS results. See DOI: 10.1039/d0em00063a



surfaces, by the formation of surface complexes on edge surfaces, or by the formation of surface clusters and precipitates. Fe^{2+} is expected to effectively compete with some trace metal cations (e.g., Zn^{2+} , Cd^{2+}) for sorption sites on clay minerals,^{9,10} leading to mobilization of trace metals under anoxic conditions. On the other hand, immobilization of trace metals can occur through additional pathways such as incorporation into newly formed solids.

Natural clay minerals often contain $\text{Fe}(\text{II})$ and $\text{Fe}(\text{III})$ in their crystal lattice^{11,12} and it has been shown that electrons can be transferred from ferrous iron to structural $\text{Fe}(\text{III})$.^{13–18} Iron-free clay minerals are generally considered to be redox-inactive within the natural redox potential of soils. However, Géhin *et al.*¹⁴ demonstrated that under strictly anoxic conditions $\text{Fe}(\text{II})$ was oxidized to $\text{Fe}(\text{III})$ on surfaces of clay minerals having very low iron contents. Potential electron acceptors in low-iron clay minerals could include trace $\text{Fe}(\text{III})$ impurities, other redox-active elements (e.g., Ti), or hydrogen.¹⁴ Following oxidation, the hydrolysis of $\text{Fe}(\text{III})$ may lead to the formation of oxyhydroxide clusters or precipitates on the clay mineral surface, as has been observed by many studies.^{12,15,19,20} However, uncertainties still prevail about the ability of Fe-free clay mineral to induce electron transfer from adsorbed $\text{Fe}(\text{II})$ to the clay mineral, leading to the formation of adsorbed $\text{Fe}(\text{III})$.^{19,21}

When reduced soils are exposed to air, oxidation of sorbed $\text{Fe}(\text{II})$ by O_2 is expected to occur, which may lead to the formation of polynuclear $\text{Fe}(\text{III})$ -oxyhydroxide clusters and surface precipitates. These oxidation products can effectively immobilize trace elements by co-precipitation.^{22–26} The nature of the surface precipitates formed by oxidation of ferrous iron depends on the reaction conditions (pH, rate of oxidation, suspension concentration, temperature, and concentration of impurities).²⁷ However, if secondary Fe phases were formed first under anoxic conditions, the type of surface precipitates formed during oxidation could also be influenced by the solid phases formed in the absence of O_2 . It has been shown by Géhin *et al.*²⁸ that $\text{Fe}(\text{OH})_2$ oxidizes into chloride green rust (Cl-GR) and further oxidation of Cl-GR leads to the formation of lepidocrocite, goethite, or akaganeite, depending on the ratio of initial concentrations of Cl^- and OH^- . However, it has not been identified which Fe-bearing minerals form by oxidation of $\text{Fe}(\text{II})$ associated with clay mineral surfaces.

Since clay minerals are not susceptible to reductive dissolution, a large portion of the Fe pool in submerged soils is believed to be sorbed or incorporated into clay minerals.²⁹ To date, few studies have been conducted on Fe^{2+} sorption on Fe-free clay minerals,^{14,21,30} which allows the determination of Fe^{2+} adsorption on clay minerals and the relevance of this process for Fe retention. Furthermore, the effects of subsequent oxidation on the speciation of Fe sorbed onto clay surfaces have not yet been determined. Therefore, the aims of this study were to investigate the interactions of Fe^{2+} with a low structural iron montmorillonite under anoxic conditions, and to characterize the Fe surface species and secondary phases formed during sorption. Additionally, we investigated the Fe species formed by oxidation of Fe sorbed on the clay mineral surface. The results

will improve our understanding of the role of clay minerals in Fe redox cycling and phase (trans-)formation.

Experimental

Preparation of clay mineral

A synthetic montmorillonite (Barasym SSM-100), referred to as Syn-1 was purchased from the Source Clays Repository of the Clay Minerals Society (Purdue University, West Lafayette, IN). Syn-1 is very low in structural Fe and was selected for this study to minimize potential oxidation of adsorbed Fe^{2+} by structural $\text{Fe}(\text{III})$. Syn-1, was size fractionated ($< 2 \mu\text{m}$), Ca saturated, and washed free of excess salt. The size-fractionated Ca-saturated Syn-1 was analyzed both as oriented and random oriented powder samples by X-ray diffraction (XRD, D8 Advance, Bruker). Diffractograms were recorded in Bragg–Brentano geometry from 3 to $80^\circ 2\theta$ with a step size of $0.02^\circ 2\theta$ and 4 s acquisition time per step, using Cu $K\alpha$ radiation ($\lambda = 1.5418 \text{ \AA}$, 40 kV, and 40 mA). X-ray fluorescence analysis (XRF, XEPOS, Spectro) was performed for elemental analysis of the clay mineral. The measured diffractograms showed trace amounts of boehmite ($\gamma\text{-AlOOH}$) and total iron content was determined by XRF as 0.026 g kg^{-1} . The specific surface area of Syn-1 was measured by multi-point N_2 -BET analysis (ASiQwin, Quantachrome) as $95 \pm 2 \text{ m}^2 \text{ g}^{-1}$.

Anoxic ferrous iron sorption experiments

Batch sorption experiments were performed to quantify Fe^{2+} sorption on Ca-saturated Syn-1 as a function of pH (4–10) and CaCl_2 concentration (0.1, 1.0, and 10 mM CaCl_2) for two fixed ferrous iron concentrations (0.05 and 0.5 mM) at $25 \pm 1^\circ \text{C}$. The equilibration time was 24 h and the clay suspensions had a solid-to-solution ratio of 1 g L^{-1} , giving Fe/clay ratios of 0.05 and 0.5 mol Fe per kg clay, respectively. Due to the high oxygen sensitivity of Fe^{2+} , all batch sorption experiments were performed under anoxic conditions ($\text{O}_2 < 0.1 \text{ ppm}$). All solutions were prepared with O_2 - and CO_2 -free, doubly deionized (DDI) water ($\geq 18.2 \text{ M}\Omega \text{ cm}$, Milli-Q, Millipore, Merck KGaA, Darmstadt, Germany), which was produced by boiling the water while purging it with pure nitrogen gas for at least 3 h outside the glovebox. This was done to prevent oxidation of dissolved Fe^{2+} and avoid the formation of carbonate solids at alkaline pH values contributing to the removal of Fe^{2+} from solution. All batch experiments were prepared in an anoxic glovebox from a stock clay suspension (25 g clay L^{-1}), which was pre-equilibrated under a N_2 atmosphere in the glovebox for 7 days. Before starting the sorption experiments, clay suspensions were equilibrated for 1 hour at pH 5 to remove CO_2 . Subsequently, aliquots of a Fe^{2+} stock solution (0.25 M made from a FeCl_2 salt) were added to the clay suspension to obtain a total Fe^{2+} concentration of 0.05 and 0.5 mM. The suspension pH was adjusted by adding a pre-determined amount of CO_2 -free base (0.01 or 0.1 M NaOH) to reach the target pH value.

The pH of the clay suspensions was measured for all samples at the end of the equilibration period. The samples were then passed through $0.22 \mu\text{m}$ nylon filters (VWR International) and






Fig. 1 Percentage of Fe^{2+} sorbed on Syn-1 ($\sim 1 \text{ g L}^{-1}$) as a function of pH at different CaCl_2 concentrations. Samples in DDI water had a Ca^{2+} concentration of $\sim 0.1 \text{ mM}$ after 1 day equilibration. Total Fe^{2+} concentrations of (a) 0.05 and (b) 0.5 mM were added and equilibrated for 1 day with 1 g L^{-1} Syn-1.

2.5 mM Fe^{2+} , and 50 mM CaCl_2 . The high CaCl_2 concentration and pH values were selected to focus on sorption processes on edge surfaces, which are expected to dominate under these conditions. Furthermore, at $\text{pH} < 8$ the dissolved Fe^{2+} concentrations in the suspensions were always below saturation with respect to $\text{Fe}(\text{OH})_2$ (Table S1 \dagger).

In the XAS samples, the sorbed fraction of Fe^{2+} was higher at $\text{pH} \sim 8$ than at $\text{pH} \sim 7$ (Table S2 \dagger). The amount of sorbed Fe^{2+} also slightly increased with increasing equilibration time from day 1 to 30 (Table S2 and Fig. S3 \dagger), e.g., by 0.02 mol Fe per kg clay at $\text{pH} 7$. This observation is consistent with findings by Zhu *et al.*,³⁴ who observed an increase in Fe^{2+} sorption to Syn-1 during 5 days at $\text{pH} 7.5$. Such increases of Fe^{2+} sorption suggest that slow sorption processes were taking place. Possible mechanisms explaining slow sorption kinetics include dissolution and precipitation reactions on surfaces.^{35–44}

The derivative of the Fe K-edge XANES spectra of the solids from selected Fe^{2+} sorption samples are shown in Fig. 2, along with relevant Fe reference compounds for comparison. The position of the highest maximum in the derivatives of XANES spectra for the high Fe-loading anoxic sorption sample at 7125 eV suggests an Fe oxidation state of close to +2. However, the presence of a small shoulder at 7128 eV might suggest that a small fraction of Fe^{2+} was oxidized in these samples. For the low Fe-loading anoxic sample at $\text{pH} \sim 8$, the position of the highest maximum is consistent with a valence state between +2 and +3, suggesting that sorbed Fe^{2+} was partially oxidized during sorption experiments. The low Fe-loading anoxic sample at $\text{pH} 7$ exhibited a maximum at 7127 eV, corresponding to a +3 oxidation state.

To quantify the oxidation states of sorbed Fe on Syn-1 under anoxic conditions with greater precision, Mössbauer spectroscopy was performed for selected solids (1 g L^{-1} clay experiments) from low and high Fe-loading samples (Fig. 3 and S8 \dagger). Fit parameters are detailed in Table S6. \dagger Anoxic spectra (Fig. 3a and b) exhibited two doublets revealing the presence of ferric iron ($\text{Fe}(\text{III})$) in addition to ferrous iron ($\text{Fe}(\text{II})$). Accordingly, the spectra were fitted with one paramagnetic doublet with large quadrupole splitting ($\Delta E_Q \sim 2.68 \text{ mm s}^{-1}$) assigned to $\text{Fe}(\text{II})$ and a second doublet with small quadrupole splitting (ΔE_Q




Fig. 2 Derivative of normalized Fe K-edge XANES spectra of Fe^{2+} sorption samples in which Syn-1 ($\sim 5 \text{ g L}^{-1}$) was reacted with low (0.25 mM) or high (2.5 mM) Fe^{2+} concentrations at $\text{pH} \sim 7$ or ~ 8 . Sorption samples were equilibrated for 1 day under anoxic conditions (green lines) and subsequently oxidized for 1 day (orange lines). Superscripts (a–d) indicate the corresponding pairs of anoxic and oxidized samples. Displayed pH values correspond to the pH measured at the end of the equilibration period for each sample. Selected reference spectra of solid phases containing $\text{Fe}(\text{II})$ and/or $\text{Fe}(\text{III})$ are shown for comparison (abbreviations: Nk = nikischerite, Cl-GR = chloride green rust, Fh = ferrihydrite).






Fig. 3 77 K Mössbauer spectra and fits of high Fe-loading solid samples. Syn-1 (1 g L^{-1}) was reacted with 0.5 mM Fe^{2+} (enriched in ^{57}Fe) at pH ~ 7 or ~ 8 under anoxic conditions during 1 day (a and b). Anoxic samples were subsequently exposed to air for 1 day (c and d). Displayed pH values correspond to the pH measured at the end of the equilibration period for each sample. In all graphs, symbols represent data and red lines the model fits. Corresponding fitting parameters are summarized in Table S6.† The Fe(II) doublet is represented as green area and the Fe(III) doublet as orange area.

$\sim 0.81 \text{ mm s}^{-1}$) consistent with Fe(III). At high Fe-loading, 14–28% of the sorbed iron was oxidized to Fe(III) under anoxic conditions and 52–100% for sorption experiments performed at low Fe-loading (Table S6†). The ferrous doublet had similar isomer shift (δ) and quadrupole splitting (ΔE_Q) values as were reported for chloride green rust ($\delta = 1.27 \text{ mm s}^{-1}$; $\Delta E_Q = 2.68 \text{ mm s}^{-1}$).^{45,46} The ferric doublet ($\Delta E_Q \sim 0.81 \text{ mm s}^{-1}$; $\delta \sim 0.46 \text{ mm s}^{-1}$) was consistent with poorly crystallized or amorphous compounds such as ferric hydroxide $\text{Fe}(\text{OH})_3$ or ferrihydrite.⁴⁷ Overall, the data clearly show that the added Fe^{2+} was partially oxidized upon sorption to Syn-1 clay under anoxic conditions. The relative abundance of Fe(III) was consistently higher at the low Fe-loading (Table S6†).

The presence of Fe(III) species in anoxic sorption samples confirms that even under anoxic conditions, Fe(II) can be oxidized to Fe(III) in the presence of clay minerals with very low structural Fe contents. The amount of structural Fe present in the Syn-1 (4.65 mmol Fe per kg) may have contributed to the oxidation of at maximum 0.93 and 9.3% of the added Fe^{2+} for high and low Fe-loading samples, respectively. Oxidation of Fe^{2+} by accidental exposure to O_2 can be excluded as no Fe^{2+} oxidation was observed in Fe^{2+} control solutions without clay. An additional anoxic control experiment on Fe^{2+} sorption to Syn-1 was carried out in a glass bottle to ensure that no residual O_2 sorbed to the walls of Nalgene® plastic bottles induced Fe^{2+} oxidation. The same results were obtained in experiments with glass and Nalgene® bottles, excluding residual sorbed O_2 as a possible electron acceptor. Therefore, other reactive sites on the clay mineral surface must have induced electron transfer from Fe^{2+} to an electron acceptor. As only traces of Ti were

present in the clay mineral (1.3 mmol Ti per kg) it can be excluded as main electron acceptor.

Géhin *et al.*¹⁴ suggested hydrogen to be a potential electron acceptor with the production of H_2 that might be sorbed by the clay mineral and ensure the reversibility of the Fe(II) oxidation. In the study by Géhin *et al.*,¹⁴ the Fe^{2+} concentration added to a Fe-free clay ($\sim 0.06 \text{ mmol Fe per g clay}$) was similar to the low Fe-loading sorption samples of this study. The partial pressure of H_2 in both experiments was very low as they were performed in a 100% N_2 atmosphere.¹⁴ Thermodynamic calculations demonstrate that at very low H_2 partial pressures, H^+ is a potential oxidant for Fe^{2+} at neutral to alkaline pH to form amorphous Fe(III)-hydroxides (Fig. S2a†). In the presence of more crystalline Fe(III) solids having lower iron solubilities such as *e.g.*, lepidocrocite and goethite, H^+ would represent an even more favorable oxidant for Fe^{2+} and might enable Fe^{2+} oxidation already at slightly acidic pH values (Fig. S2b†). The reduction of H^+ would lead to the production of H_2 and an increase of the partial pressure of H_2 . Concurrently, the Fe^{2+} activity is expected to decrease due to the formation of Fe(III) solid phases and progressing Fe^{2+} adsorption to mineral surfaces. Consequently, the ability of H^+ to act as an oxidant for Fe^{2+} decreased during sample equilibration (see arrows in Fig. S2a†). This explains why the oxidation of Fe^{2+} was limited in our experiments. Accordingly, an increased percentage of Fe^{2+} oxidation with lower Fe-loading is expected, which is consistent with previous findings by Soltermann *et al.* and Géhin *et al.*^{14,30} confirming that the oxidation of Fe^{2+} is likely limited by the partial pressure of H_2 and the Fe^{2+} activity in solution. In a control experiment without clay minerals present, Fe^{2+} oxidation was not observed.



Based on this finding, we propose that clay minerals served as a catalyst for Fe^{2+} oxidation by H^+ by facilitating the electron transfer *via* ligation of Fe^{2+} to protonated surface oxygen atoms. This catalytic effect of surfaces in electron transfer reactions may be important in various environmental settings. For instance, it has been shown that the reduction of organic pollutants by Fe^{2+} is enhanced when sorbed to mineral surfaces.^{48–50}

To further investigate the possible solid-phase Fe species formed during the 1 and 30 days equilibration periods, linear combination fits (LCF) of Fe K-edge EXAFS spectra were performed (Fig. 4, Tables S11 and S12†). LCF analysis showed that EXAFS spectra of high Fe-loading anoxic samples were adequately fitted by including Fe(II)-hydroxide ($\beta\text{-Fe}(\text{OH})_2$) (6–13%), chloride green rust ($[\text{Fe}_3^{2+}\text{Fe}^{3+}(\text{OH})_8] \cdot [\text{Cl} \cdot n\text{H}_2\text{O}]^{45}$) (51–68%), and nikischerite ($\text{NaFe}^{2+}_6\text{Al}_3(\text{SO}_4)_2(\text{OH})_{18}(\text{H}_2\text{O})_{12}$)⁵¹ (19–40%). Nikischerite is a natural Fe(II)Al(III)-layered double hydroxide (LDH) mineral with sulfate interlayer anions and is used here as a structural reference representing Fe(II)Al(III)-LDH phases in general.⁵¹ Fitting of the low Fe-loading anoxic samples (Fig. 4b) required the reference compound ferrihydrite, which is in line with the presence of Fe(III) as observed by both Mössbauer and XANES spectroscopy.

An improved fit (>10% decrease in *R*-factor, see Table S12†) was obtained for the low Fe-loading anoxic sorption sample

equilibrated at pH ~ 8 when Fe(II)-containing minerals, namely nikischerite and chloride green rust, were included. This is again consistent with the presence of Fe(II) species, as observed by Mössbauer spectroscopy. A comparison of the Fe oxidation states determined by Mössbauer spectroscopy for the 1 g L⁻¹ batch experiments and by LCF of EXAFS spectra for 5 g L⁻¹ batch samples is provided in Table 1. Both sets of batch experiments are in reasonable agreement and thus confirm that Fe^{2+} in anoxic Fe sorption samples can partially be oxidized in the presence of Syn-1 clay.

For the majority of the samples, the LCF significantly improved when an Fe(III)-containing clay mineral reference (SWy-2, containing 3 wt% Fe) was included as additional reference in the fits. Released Si from Syn-1 after 1 day equilibration between pH 6 and 10 amounted on average to ~ 0.016 mmol Si per g clay compared to up to 8 times lower Si concentrations for anoxic sorption samples with Fe (see Fig. S6a and S7†). This provides additional evidence that the released Si from Syn-1 was removed from solution upon addition of Fe, either by adsorption or incorporation into newly formed Fe phases. It has been suggested by Soltermann *et al.*³⁰ that Fe^{2+} can be taken up by the clay minerals, leading to the formation of Fe-bearing clay minerals.

In presence of elevated divalent metal (Me^{2+}) concentrations (first-row transition series), it has been shown that $\text{Me}(\text{II})\text{Al-LDH}$




Fig. 4 LCF fitting results of Fe K-edge EXAFS spectra of Fe^{2+} sorption samples equilibrated for 1 day under anoxic conditions (green lines) and subsequently oxidized for 1 day (orange lines) and relevant references: (a and b) absolute fractions of Fe references obtained by linear combination fitting of k^3 -weighted Fe K-edge EXAFS spectra for Fe^{2+} sorbed on Syn-1. High Fe load (a and c) and low (b and c) (0.25 mM) sorption samples. The fits were performed over a k -range of 2–11.3 \AA^{-1} ($k_w = 3$) with no fit constraints. The initial fit fractions (70.9–146.2%) were normalized to the sum equal to 100% and are reported in Tables S11 and S12† (abbreviations: Nk = nikischerite, Cl-GR = chloride green rust, Lp = lepidocrocite, Fh = ferrihydrite). (c) k^3 -weighted χ spectra of references (line): SWy-2, $\text{Fe}(\text{OH})_2$, nikischerite, chloride green rust, ferrihydrite and lepidocrocite are shown together with the k^3 -weighted χ spectra (open circles) of 1 day equilibrated samples with corresponding fit (green and orange colors correspond to anoxic and oxidized samples, respectively). Displayed pH values correspond to the pH measured at the end of the equilibration period for each sample. Superscripts (a–d) indicate the corresponding pairs of anoxic and oxidized samples.



Table 2 Shell-fit parameters determined from Fe K-edge EXAFS data of Fe(II) sorption to Syn-1 under anoxic conditions for 1 day.^a Parameter uncertainties are given in parentheses

	Fe–O			Fe–Fe			ΔE_0^e	Red. χ^2^f	R-factor ^g
	CN ^b	R^c (Å)	σ^{2d} (Å ²)	CN ^b	R^c (Å)	σ^{2d} (Å ²)			
High Fe load									
pH 7.2	4.1 (0.8)	2.10 (0.01)	0.009 (0.003)	2.6 (0.9)	3.21 (0.01)	0.006 (0.003)	1.17 (1.79)	110	0.0211
pH 8.1	6.0 (0.5)	2.12 (0.01)	0.009 (0.001)	4.5 (0.5)	3.20 (0.01)	0.007 (0.001)	–1.53 (0.7)	235	0.0032
Low Fe load									
pH 8.1	4.6 (0.9)	2.06 (0.02)	0.012 (0.003)	2.1 (0.4)	3.17 (0.02)	0.009	1.25 (1.81)	14	0.0274

	Fe–O			Fe–Al			Fe–Si			Fe–Fe			ΔE_0^e	Red. χ^2^f	R-factor ^g
	CN ^b	R^c (Å)	σ^{2d} (Å ²)	CN ^b	R^c (Å)	σ^{2d} (Å ²)	CN ^b	R^c (Å)	σ^{2d} (Å ²)	CN ^b	R^c (Å)	σ^{2d} (Å ²)			
Low Fe load															
pH 7.0	4.9 (1.3)	2.02 (0.02)	0.010 (0.003)	2.9 (1.6)	2.82 (0.04)	0.01	1.9 (1.9)	3.03 (0.07)	0.01	1.1 (0.8)	3.48 (0.05)	0.009 (2.70)	0.98	210	0.0335

^a The amplitude reduction factor S_0^2 , was set to 0.9 based on first shell optimization for all sorption samples (R -range 1.1–2.5 Å). ^b Path degeneracy (coordination number). ^c Mean half path length. ^d Debye–Waller parameter. Debye–Waller parameter was fixed to 0.01 Å² for Fe–Al and Fe–Si based on the σ^2 obtained for Fe–Si and Fe–Al path by Soltermann *et al.*³⁰ σ^2 (Fe) was fixed to 0.009 Å² for Fe–Fe of low Fe-load sorption sample like was fitted for edge-sharing Fe–Fe for low Fe-load samples. ^e Energy-shift parameter. ^f Fit accuracy; reduced $\chi^2 = (N_{\text{idp}}/N_{\text{pts}}) \sum_i ((\text{data}_i - \text{fit}_i)/\epsilon_i)^2 (N_{\text{idp}} - N_{\text{var}})^{-1}$. N_{idp} , N_{pts} and N_{var} are, respectively, the number of independent points in the model fit (11.2–13.7), the total number of data points (151–171), and the number of fit variables (6–10). ϵ_i is the uncertainty of the i^{th} data point. ^g R-factor; Normalized sum of squared residuals $(\sum_i (\text{data}_i - \text{fit}_i)^2 / \sum_i \text{data}_i^2)$. Note: The FT of k^3 -weighted EXAFS spectrum of the anoxic low Fe-loading samples at pH 8.1 was calculated over a k -range of 2–9.5 Å^{–1}. The shell-fit analysis of k^3 -weighted Fe EXAFS spectrum of the anoxic low Fe-loading sample at pH 7.0 was performed in R -space $R + \Delta R$ -range of 1.1–4 Å.

between the isosbestic points confirms the successful preparation of a clay film with preferred orientation.^{59–62} The angle dependence of the spectra also confirms the anisotropic formation of the secondary phase aligned with the clay mineral sheets for high Fe-loading anoxic sorption samples.

For the low Fe-loading anoxic samples, the first shell was fitted with 4.6–5.3 O atoms at a radial distance of 2.01–2.05 Å which is an intermediate distance of an octahedral arrangement of O atoms around Fe(II) and Fe(III).^{56,63} The fitting of a CN < 6 may suggest the presence of some tetrahedral Fe as found in ferrihydrite. This result is in agreement with the presence of Fh in the LCF of EXAFS spectra of this sample (Table S12†).

The FT of the EXAFS spectra of the low Fe-loading anoxic samples equilibrated at pH ~8, showed a second peak at ~2.75 Å (uncorrected for phase shift). This second shell was fitted with Fe located at a radial distance of 3.15–3.17 Å and a coordination number of 1.8–2.1. The fitted distance for the second shell corresponds with an intermediate distance of edge-sharing Fe atoms around Fe(II) and Fe(III) as in fougérite⁵⁶ and lepidocrocite,⁶⁴ respectively. This is consistent with the results obtained by LCF and Mössbauer spectroscopy.

The low Fe-loading anoxic sample equilibrated at pH ~7 (Table 2) showed two overlapping small peak after the first shell, one at ~2.3 Å and ~3.1 Å (uncorrected for phase shift). Fitting of

these two shells required a Fe–Al, Fe–Si and Fe–Fe backscattering path with distances of 2.84, 3.08 and 3.48 Å, respectively. Fitted distances for Fe–Al and Fe–Si paths are in good agreement with bond distances in phyllosilicates.⁶⁵ The distance fitted for the Fe–Fe path at 3.48 Å is characteristic for double corner-sharing Fe as in ferrihydrite.^{63,66}

The WT of EXAFS spectra of low Fe-loading anoxic sample equilibrated at pH ~7 (see Fig. S14c†) contained three maxima, one at $k = 3, 7$, and 10 \AA^{-1} suggesting three different backscattering atoms contributing to the second shell. This result confirms that including the Fe–Fe, Fe–Al and Fe–Si backscattering paths in the model to fit low Fe-loading anoxic samples equilibrated at pH ~7 is reasonable. Additionally, F -tests confirmed that the inclusion of Fe–Al and Fe–Si backscattering paths into the fit model significantly improved the goodness of the fit (S.I. Section 11†).

Previous studies have also investigated the sorption of Fe²⁺ to different clay minerals at various Fe-loadings.^{14,30} Collectively, the literature results suggested that depending on the Fe-loading, various Fe(II)- and Fe(III)-containing solids formed. At high Fe-loadings (0.53–2 mol Fe per kg) the formed secondary phases were predominantly Fe(II)-containing phases.^{19,21} At low Fe-loadings (0.002–0.08 mol Fe per kg) mainly Fe(III)-containing phases were observed.^{30,67} Our current study are in agreement



with these results. The formation of an LDH phase with Fe in the octahedral sheet was observed for high Fe-loading anoxic sorption samples as shown by Jones *et al.*¹⁹ Contrary, in the study by Zhu *et al.*²¹ no Fe²⁺ oxidation was observed during sorption of Fe²⁺ onto Syn-1 and the formation of FeAl-LDH was suggested as the main solid phase forming. However, sorption experiments in the study by Zhu *et al.* were performed in an atmosphere with 5% H₂ and 95% N₂. Such elevated H₂ concentration are expected to disable H⁺ to oxidize Fe²⁺ (*cf.* Fig. S2†) and to suppress the formation of green rust as the main secondary phase. In our low Fe-loading samples, over half of Fe(II) was oxidized to Fe(III), from which a large fraction was incorporated into newly formed clay minerals. This result shows that clay minerals can play an important role in the mobility of Fe²⁺ under anoxic conditions.

Effects of aeration of anoxic sorption samples on Fe speciation

To understand the effects of O₂-exposure on the speciation of Fe in clay suspensions (Syn-1) that were anoxically equilibrated for 1 or 30 days with Fe²⁺, the anoxic suspensions were exposed to ambient air for 24 hours. Aeration resulted in a further increase in percent Fe sorption (see Fig. 4a, b and Table S2†), despite a drop in suspension pH (between 0.8 and 3.1 pH values) as a result of hydrolysis of Fe³⁺ (Table S3†). Due to the high O₂ sensitivity of Fe²⁺ at near-neutral pH, rapid oxidation of adsorbed and dissolved Fe²⁺, hydrolysis of Fe³⁺, and precipitation of Fe(III)-(oxyhydr)oxides may occur. This results in the release of protons, explaining the observed decrease of suspension pH.

After the equilibration of Fe²⁺ with Syn-1 under anoxic conditions, substantial amounts of aqueous Fe²⁺ still remained in solution (70–78% of added Fe at pH ~7 and 7–16% at pH ~8, see Table S2†). Additionally, some adsorbed Fe²⁺ may have been desorbed from clay surfaces as pH started to decrease. All dissolved and adsorbed Fe²⁺ is expected to rapidly oxidize upon exposure to O₂ and form iron minerals such as lepidocrocite and/or ferrihydrite, depending on solution composition (*e.g.*, Ca, Si).⁶⁸ However, the remaining 22 to 30% (at pH ~7) or 83 to 93% (pH ~8) of the added Fe was sorbed to the clay, mostly by the formation of Fe(II) solid phases such as chloride green rust, FeAl-LDH, and/or Fe(III)-phyllosilicates, as was shown in the section above. The oxidation kinetics and resulting Fe species formed during aeration of such sorbed ferric Fe on clay minerals is far less understood.

Mössbauer spectroscopy showed that the 24 h aeration of anoxic Fe²⁺ sorption samples led to a complete oxidation of all added Fe, resulting in only a ferric doublet in the spectra of oxic samples (Fig. 3c and d). This finding shows that even 30 days equilibration under anoxic conditions, resulting in mainly the incorporation of Fe(II) into solid phases, did not prevent rapid oxidation to Fe(III). Additional Mössbauer spectra of two high Fe-loading oxic samples were measured at 45, 35, 25 and 15 K (Fig. S10†). The fitting results of the temperature-dependent Mössbauer spectra are given in Tables S7 and S8.† In general, the spectra displayed the characteristic effect of magnetic ordering with decreasing temperature. Ordered sextets had

hyperfine parameters δ (between 0.49 and 0.50 mm s⁻¹), ΔE_Q (between -0.05 and -0.01 mm s⁻¹) and hyperfine field (B_{hf}) (between 43.9 and 44.3 T) for the oxic samples at 15 K, which are in line with values reported for short range ordered Fe(III)-oxyhydroxides such as lepidocrocite and ferrihydrite.⁶⁹ Additionally, temperature dependent Mössbauer spectra allow the determination of the blocking temperature, which is affected by crystallite size or interparticle interactions.^{70,71} We calculated the blocking temperature as the temperature at which the summed areas of the paramagnetic doublets and magnetic sextets (which includes poorly ordered and fully magnetically ordered sextets) of Mössbauer spectra are equal. Fig. S9† shows the relative abundance of doublets and sextets for the two oxic samples at pH 5.1 (pH 7.3 under anoxic conditions) and pH 5.6 (pH 8.4 under anoxic conditions). Both samples displayed a blocking temperature below the literature values for ferrihydrite and lepidocrocite (between 50 and 77 K).^{69,72} A blocking temperature of ~32 K was calculated for the oxic sample pre-equilibrated at pH 7.3, compared to ~42 K for the aerated suspension pre-equilibrated at pH 8.4. This indicates that oxidation of the anoxic sorption samples led to the formation of short range ordered Fe(III)-(oxyhydr)oxides, for which either precipitates had smaller crystallite size or show a lesser degree of interparticle interaction, potentially as a result of surface sorption effects, when formed at pH 7.3 than pH 8.4.

For all oxic sorption samples, the position of the Fe K-edge was at 7127 eV (Fig. 2), which corresponds to the oxidation state Fe(III) and is in line with Mössbauer data (Fig. 2). Linear combination fitting (LCF) results for Fe K-edge EXAFS spectra are presented in Fig. 4 (also Tables S11 and S12†). Oxidized sorption samples were well described (*R*-factor < 2%) with 3 reference compounds: ferrihydrite (45–60%), Fe(III)-containing phyllosilicates (24–36%), and lepidocrocite (13–19%). Essentially, all oxic sorption samples transformed into the same secondary phases regardless of their equilibration conditions (pH, Fe concentration) and time (1 or 30 days) under anoxic conditions (Tables S11 and 12†). Parts of the Fe(III)-containing phyllosilicates that have already formed under anoxic conditions were not affected by oxidation and remained as secondary phase after aeration. It was observed that the fraction of lepidocrocite formed was significantly lower for low Fe-loading samples compared to high Fe-loading samples.

The oxidation of aqueous ferrous iron at pH 7 and in a chloride background is known to lead to the formation of lepidocrocite, with crystallinity decreasing with decreasing pH.²⁷ Below pH 4, oxidation of ferrous iron is expected to precipitate as ferrihydrite.²⁷ As the pH of the oxic sorption samples remained above pH 4 and the concentration of Ca was elevated, the formation of lepidocrocite as the major Fe(III)-(oxyhydr)oxides is expected. However, the presence of dissolved silicate, released from the clay mineral, above 10 $\mu\text{mol L}^{-1}$ has been shown to inhibit the formation of more crystalline iron oxides, such as lepidocrocite,⁷³ and favor the formation of ferrihydrite.^{66,68} Therefore, the observation of ferrihydrite as the dominant secondary phase under oxic conditions in this study was more anticipated, as Si released from the clay mineral amounted to ~80 $\mu\text{mol Si L}^{-1}$ between pH 6 and 10 (Fig. S6a†). The



observation of lepidocrocite as a second Fe(III)-(oxyhydr)oxides in the oxic samples can be explained by two mechanisms: (i) through the oxidation of green rust, which under certain conditions can be oxidized into lepidocrocite²⁸ or (ii) through dissolved ferrous iron catalyzing the transformation of poorly-crystalline ferrihydrite to the more stable mineral lepidocrocite.^{74,75}

A lower amount of lepidocrocite was formed in the low Fe-load samples, as direct oxidation of Fe²⁺ in presence of elevated Si concentrations (high Si/Fe ratio in solution) may mainly precipitate as ferrihydrite.⁷³ However, only a small fraction of the sorbed Fe in the oxic samples was formed by direct oxidation of dissolved Fe²⁺ for low Fe-loading samples equilibrated at pH ~8 (between 7 and 12%, see Fig. S4 and S5†) compared to samples equilibrated at pH ~7 (between 60 and 65%, see Fig. S4 and S5†). Nevertheless both low Fe-loading samples resulted in the presence of similar solid phases. The higher fraction of lepidocrocite observed in the high Fe-loading oxic samples as compared to the low Fe-loading samples may be due to the oxidation of dissolved Fe²⁺ in the presence of a low Si/Fe ratio in solutions, which favors lepidocrocite formation over ferrihydrite or through the transformation of green rust into lepidocrocite.^{66,73} Because only a small fraction of the sorbed Fe (up to 20%, see Fig. S4 and S5†) was formed by direct oxidation of dissolved Fe²⁺ in high Fe-loading samples, lepidocrocite was primarily formed from the transformation of Fe-phases precipitated under anoxic conditions.

The LCF of EXAFS spectra of oxic samples significantly improved when an Fe(III)-containing clay mineral reference (SWy-2, representing Fe(III) in phyllosilicate structures) was included. Furthermore, measured Si in solution of oxic samples was 6–40 times lower than the Si released from Syn-1 after 1 day equilibration between pH 6 and 10 in absence of Fe (~0.016 mmol Si per g clay) (Fig. S6a and S7†). Combined, this provides strong evidence for sorption and/or incorporation of Si into the Fe species formed upon aeration. Based on the LCF results, the amount of Fe sorbed by the formation of Fe-phyllosilicates (the fraction fitted by SWy-2) was calculated. Together with amount Si re-sorbed upon Fe²⁺ addition, this allowed to calculate the ratio of Fe to Si sorbed in the samples. The overall Fe/Si ratio of the newly formed Fe species was estimated to amount to 0.56–1.83 (Tables S4 and S5†). These values are in line with Fe/Si ratios one would expect for 1 : 1 Fe-phyllosilicate, like (Fe²⁺,Fe³⁺)_{3–2}Si₂O₅OH and 2 : 1 Fe-phyllosilicate ((Fe²⁺,Fe³⁺)_{3–2}Si₄O₁₀(OH)₂).⁷⁶

More detailed information about changes in local coordination environment of Fe upon aeration of the suspensions was obtained by shell-by-shell fitting of fourier transformed *k*³-weighted Fe K-edge EXAFS spectra (Fig. S11†). All oxic sorption samples showed a first shell at ~1.5 Å (uncorrected for phase shift), representing the first oxygen shell surrounding the Fe atoms. A second shell was observed between 2 and 3.5 Å, which can be attributed to different Fe–Fe backscattering pairs. The first shell was fitted with 4.9–6.0 O atoms at a radial distance of 1.99–2.00 Å. This is consistent with an octahedral arrangement of the O atoms around a central Fe(III), as in lepidocrocite.⁶³ The fitting of a CN smaller than 6 may suggest the presence of some tetrahedral Fe as found in ferrihydrite.⁷⁷ This result is in

agreement with the presence of Fh in the LCF of EXAFS spectra of oxic sample (Tables S11 and S12†). The second shell was composed of two peaks, one at ~2.65 Å and a second peak at ~3.11 Å (uncorrected for phase shift). These two peaks were successfully fitted with two Fe–Fe single backscattering paths. The first Fe backscattering path (Fe–Fe₁) had a fitted distance of 3.07–3.09 Å with coordination numbers between 1.8–3.7, consistent with edge-sharing Fe octahedral.⁶⁴ The second Fe backscattering path (Fe–Fe₂) had a fitted distance of 3.42–3.45 Å with coordination numbers between 1.2–2.1, characteristic of double corner-sharing Fe octahedral.^{63,66} The fitted coordination numbers of edge-sharing Fe (Fe–Fe₁) atoms increased with increasing Fe-loading, contrary to the fitted coordination numbers of corner-sharing Fe (Fe–Fe₂) atoms (Tables S14 and S15†). These trends correspond well with the increasing fraction of lepidocrocite formed with increasing Fe-loading, as observed by LCF of EXAFS spectra of oxic sorption samples. Triangular multiple scattering (MS) paths Fe–O–O were not implemented in our fit model as including this path decreased the reduced χ^2 -values less than a factor two (see S.I. Section 15 for details†).⁵⁵

Conclusions

Our results show that the formation of Fe-bearing secondary phases is the main sorption mechanism for Fe²⁺ uptake by clay minerals low in structural Fe under anoxic conditions. Clay minerals can enhance the formation of newly formed, layered Fe minerals such as green rusts, FeAl-LDH, and Fe-phyllosilicates, by providing nucleation sites and serving as a source of Al and Si. Furthermore, clay surfaces may act as catalysts for Fe²⁺ oxidation by hydrogen, especially at circum-neutral to alkaline pH values and low H₂ partial pressures. Therefore, the precipitation of Fe-bearing phases in presence of clays may be of importance in controlling Fe²⁺ solubility in anoxic groundwaters and soils.

The nature of the secondary phases formed under anoxic conditions appeared to be controlled by Fe-loading, but also by other factors. Under the conditions used in our study (high Ca²⁺, no CO₂), the formation of layered Fe minerals was favored. These conditions were chosen to ensure saturation of cation exchange sites with Ca²⁺ and exclude formation of iron carbonates, allowing us to identify the sorption mechanisms of Fe²⁺ to edge sites of the clay mineral. However, as submerged soils frequently contain elevated CO₂ concentrations, the precipitation of carbonate green rust and also siderite (FeCO₃) would also be expected in these environments.

After exposure of anoxic sorption samples to ambient air, sorbed Fe(II) was completely oxidized to Fe(III). Aeration led to a rapid transformation of sorbed Fe into ferrihydrite, lepidocrocite, and Fe(III)-bearing clay minerals, independent of the Fe-loading and the duration of the pre-equilibration under anoxic conditions. This demonstrates that clay minerals are unable to stabilize sorbed Fe(II) against oxidation. However, the presence of clay minerals favors the formation of Fe(III)-containing clays both under anoxic and oxic conditions. This process is expected to impact not only the solubility of iron in soils and sediments but also many redox transformations of



organic and inorganic contaminants which can react with Fe-bearing clays.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank M. Fischer and K. Barmettler (Soil Chemistry Group, ETH Zurich) for assistance in the laboratory. We gratefully acknowledge M. Etique (University of Lorraine) for discussions on the synthesis of Fe(II) minerals and M. Siebecker (Texas Tech University) about Wavelet transformation. We acknowledge SOLEIL (proposal no. 20171097) for the provision of synchrotron radiation facilities and thank G. Landrot, E. Fonda and G. Alizon for their support during the synchrotron measurements. We are grateful to E. Elzinga (Rutgers University) for sharing with us the reference XAS spectra of Fe(OH)₂ and nikischerite, T. Borch (Colorado State University) the XAS spectra of chloride green rust and A. Zitolo (SOLEIL) for the XAS spectra of aqueous Fe²⁺ and Fe³⁺. This research was funded by the Swiss National Science Foundation (grant no. 200021_156392).

References

- 1 P. Refait, L. Simon and J. M. R. Genin, Reduction of SeO₄²⁻ anions and anoxic formation of iron(II)-iron(III) hydroxy selenate green rust, *Environ. Sci. Technol.*, 2000, **34**, 819–825.
- 2 V. Badaut, M. L. Schlegel, M. Descostes and G. Moutiers, In situ time-resolved X-ray near-edge absorption spectroscopy of selenite reduction by siderite, *Environ. Sci. Technol.*, 2012, **46**, 10820–10826.
- 3 J. Jonsson and D. M. Sherman, Sorption of As(III) and As(V) to siderite, green rust (fougerite) and magnetite: Implications for arsenic release in anoxic groundwaters, *Chem. Geol.*, 2008, **255**, 173–181.
- 4 S. Sengupta, P. K. Mukherjee, T. Pal and S. Shome, Nature and origin of arsenic carriers in shallow aquifer sediments of Bengal Delta, India, *Environ. Geol.*, 2004, **45**, 1071–1081.
- 5 T. Pal and P. K. Mukherjee, 'Orange sand' - A geological solution for arsenic pollution in Bengal delta, *Curr. Sci.*, 2008, **94**, 31–33.
- 6 W. L. Lindsay and A. P. Schwab, The chemistry of iron in soils and its availability to plants, *J. Plant Nutr.*, 1982, **5**, 821–840.
- 7 W. Stumm and B. Sulzberger, The cycling of iron in natural environments - Considerations based on laboratory studies of heterogeneous redox processes, *Geochim. Cosmochim. Acta*, 1992, **56**, 3233–3257.
- 8 D. L. Sparks, *Environmental soil chemistry*, Academic Press, San Diego, 1995.
- 9 M. H. Bradbury and B. Baeyens, Modelling the sorption of Mn(II), Co(II), Ni(II), Zn(II), Cd(II), Eu(III), Am(III), Sn(IV), Th(IV), Np(V) and U(VI) on montmorillonite: Linear free energy relationships and estimates of surface binding constants for some selected heavy metals and actinides, *Geochim. Cosmochim. Acta*, 2005, **69**, 875–892.
- 10 D. Soltermann, M. Marques Fernandes, B. Baeyens, J. Miehé-Brendlé and R. Dähn, Competitive Fe(II)–Zn(II) uptake on a synthetic montmorillonite, *Environ. Sci. Technol.*, 2014, **48**, 190–198.
- 11 J. B. Dixon, S. B. Weed and R. C. Dinauer, *Minerals in soil environments*, Soil Science Society of America, Madison, Wis., USA, 2nd edn, 1989.
- 12 F. Favre, J. W. Stucki and P. Boivin, Redox properties of structural Fe in ferruginous smectite. A discussion of the standard potential and its environmental implications, *Clays Clay Miner.*, 2006, **54**, 466–472.
- 13 A. Neumann, T. L. Olson and M. M. Scherer, Spectroscopic evidence for Fe(II)–Fe(III) electron transfer at clay mineral edge and basal sites, *Environ. Sci. Technol.*, 2013, **47**, 6969–6977.
- 14 A. Géhin, J.-M. Grenèche, C. Tournassat, J. Brendlé, D. G. Rancourt and L. Charlet, Reversible surface-sorption-induced electron-transfer oxidation of Fe(II) at reactive sites on a synthetic clay mineral, *Geochim. Cosmochim. Acta*, 2007, **71**, 863–876.
- 15 M. V. Schaefer, C. A. Gorski and M. M. Scherer, Spectroscopic evidence for interfacial Fe(II)–Fe(III) electron transfer in a clay mineral, *Environ. Sci. Technol.*, 2011, **45**, 540–545.
- 16 C. A. Gorski, M. Aeschbacher, D. Soltermann, A. Voegelin, B. Baeyens, M. Marques Fernandes, T. B. Hofstetter and M. Sander, Redox properties of structural Fe in clay minerals. 1. Electrochemical quantification of electron-donating and -accepting capacities of smectites, *Environ. Sci. Technol.*, 2012, **46**, 9360–9368.
- 17 C. A. Gorski, L. Klupfel, A. Voegelin, M. Sander and T. B. Hofstetter, Redox properties of structural Fe in clay minerals. 2. Electrochemical and spectroscopic characterization of electron transfer irreversibility in ferruginous smectite, SWa-1, *Environ. Sci. Technol.*, 2012, **46**, 9369–9377.
- 18 C. A. Gorski, L. E. Klupfel, A. Voegelin, M. Sander and T. B. Hofstetter, Redox properties of structural Fe in clay minerals: 3. Relationships between smectite redox and structural properties, *Environ. Sci. Technol.*, 2013, **47**, 13477–13485.
- 19 A. M. Jones, C. A. Murphy, T. D. Waite and R. N. Collins, Fe(II) interactions with smectites: Temporal changes in redox reactivity and the formation of green rust, *Environ. Sci. Technol.*, 2017, **51**, 12573–12582.
- 20 T. B. Hofstetter, R. P. Schwarzenbach and S. B. Haderlein, Reactivity of Fe(II) species associated with clay minerals, *Environ. Sci. Technol.*, 2003, **37**, 519–528.
- 21 Y. Zhu and E. J. Elzinga, Formation of layered Fe(II)-hydroxides during Fe(II) sorption onto clay and metal-oxide substrates, *Environ. Sci. Technol.*, 2014, **48**, 4937–4945.
- 22 T. Huynh, A. R. Tong, B. Singh and B. J. Kennedy, Cd-substituted goethites - A structural investigation by synchrotron X-ray diffraction, *Clays Clay Miner.*, 2003, **51**, 397–402.
- 23 L. N. Moyes, R. H. Parkman, J. M. Charnock, D. J. Vaughan, F. R. Livens, C. R. Hughes and A. Braithwaite, Uranium



- uptake from aqueous solution by interaction with goethite, lepidocrocite, muscovite, and mackinawite: An X-ray absorption spectroscopy study, *Environ. Sci. Technol.*, 2000, **34**, 1062–1068.
- 24 G. Ona-Nguema, G. Morin, F. Juillot, G. Calas and G. E. Brown, EXAFS analysis of arsenite adsorption onto two-line ferrihydrite, hematite, goethite, and lepidocrocite, *Environ. Sci. Technol.*, 2005, **39**, 9147–9155.
- 25 X. W. Xu, C. Chen, P. Wang, R. Kretzschmar and F. J. Zhao, Control of arsenic mobilization in paddy soils by manganese and iron oxides, *Environ. Pollut.*, 2017, **231**, 37–47.
- 26 K. Ehlert, C. Mikutta and R. Kretzschmar, Impact of birnessite on arsenic and iron speciation during microbial reduction of arsenic-bearing ferrihydrite, *Environ. Sci. Technol.*, 2014, **48**, 11320–11329.
- 27 U. Schwertmann and R. M. Cornell, *Iron oxides in the laboratory: preparation and characterization*, Wiley-VCH, Weinheim, New York, 2nd completely rev. and extended edn, 2000.
- 28 J. M. R. Génin, R. Aïssa, A. Géhin, M. Abdelmoula, O. Benali, V. Ernsten, G. Ona-Nguema, C. Upadhyay and C. Ruby, Fougerite and Fe(II)-III hydroxycarbonate green rust; ordering deprotonation and/or cation substitution; structure of hydrocalcite-like compounds and mythic ferrosic hydroxide $\text{Fe}(\text{OH})^{(2+X)}$, *Solid State Sci.*, 2005, **7**, 545–572.
- 29 J. W. Stucki, B. A. Goodman and U. Schwertmann, *Iron in soils and clay minerals*, D. Reidel, Dordrecht, 1988.
- 30 D. Soltermann, M. Marques Fernandes, B. Baeyens, R. Dähn, J. Miehé-Brendlé, B. Wehrli and M. H. Bradbury, Fe(II) sorption on a synthetic montmorillonite. A combined macroscopic and spectroscopic study, *Environ. Sci. Technol.*, 2013, **47**, 6978–6986.
- 31 K. Lagarec and D. G. Rancourt, *Recoil – Mössbauer spectral analysis software for Windows.*, University of Ottawa, Ottawa, 1998.
- 32 D. G. Rancourt and J. Y. Ping, Voigt-based methods for arbitrary-shape static hyperfine parameter distributions in mossbauer-spectroscopy, *Nucl. Instrum. Methods Phys. Res., Sect. B*, 1991, **58**, 85–97.
- 33 H. Funke, A. C. Scheinost and M. Chukalina, Wavelet analysis of extended x-ray absorption fine structure data, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 2005, **71**, 094110.
- 34 Y. Zhu, J. J. Liu, O. Goswami, A. A. Rouff and E. J. Elzinga, Effects of humic substances on Fe(II) sorption onto aluminum oxide and clay, *Geochem. Trans.*, 2018, **19**, 3.
- 35 D. L. Sparks, Kinetics of Ionic Reactions in Clay-Minerals and Soils, *Adv. Agron.*, 1985, **38**, 231–266.
- 36 A. M. Scheidegger, G. M. Lamble and D. L. Sparks, Spectroscopic evidence for the formation of mixed-cation hydroxide phases upon metal sorption on clays and aluminum oxides, *J. Colloid Interface Sci.*, 1997, **186**, 118–128.
- 37 A. C. Scheinost, R. G. Ford and D. L. Sparks, The role of Al in the formation of secondary Ni precipitates on pyrophyllite, gibbsite, talc, and amorphous silica: A DRS study, *Geochim. Cosmochim. Acta*, 1999, **63**, 3193–3203.
- 38 E. J. Elzinga and D. L. Sparks, Nickel sorption mechanisms in a pyrophyllite-montmorillonite mixture, *J. Colloid Interface Sci.*, 1999, **213**, 506–512.
- 39 A. M. Scheidegger, D. G. Strawn, G. M. Lamble and D. L. Sparks, The kinetics of mixed Ni-Al hydroxide formation on clay and aluminum oxide minerals: A time-resolved XAFS study, *Geochim. Cosmochim. Acta*, 1998, **62**, 2233–2245.
- 40 A. M. Scheidegger, G. M. Lamble and D. L. Sparks, Investigation of Ni sorption on pyrophyllite: An XAFS study, *Environ. Sci. Technol.*, 1996, **30**, 548–554.
- 41 A. C. Scheinost and D. L. Sparks, Formation of layered single- and double-metal hydroxide precipitates at the mineral/water interface: A multiple-scattering XAFS analysis, *J. Colloid Interface Sci.*, 2000, **223**, 167–178.
- 42 W. Li, K. J. T. Livi, W. Q. Xu, M. G. Siebecker, Y. J. Wang, B. L. Phillips and D. L. Sparks, Formation of crystalline Zn-Al layered double hydroxide precipitates on gamma-alumina: The role of mineral dissolution, *Environ. Sci. Technol.*, 2012, **46**, 11670–11677.
- 43 A. N. Starcher, E. J. Elzinga and D. L. Sparks, Formation of a mixed Fe(II)-Zn-Al layered hydroxide: Effects of Zn co-sorption on Fe(II) layered hydroxide formation and kinetics, *Chem. Geol.*, 2017, **464**, 46–56.
- 44 A. N. Starcher, W. Li, R. K. Kukkadapu, E. J. Elzinga and D. L. Sparks, Fe(II) sorption on pyrophyllite: Effect of structural Fe(III) (impurity) in pyrophyllite on nature of layered double hydroxide (LDH) secondary mineral formation, *Chem. Geol.*, 2016, **439**, 152–160.
- 45 J. M. R. Génin, G. Bourrie, F. Trolard, M. Abdelmoula, A. Jaffrezic, P. Refait, V. Maitre, B. Humbert and A. Herbillon, Thermodynamic equilibria in aqueous suspensions of synthetic and natural Fe(II)-Fe(III) green rusts: Occurrences of the mineral in hydromorphic soils, *Environ. Sci. Technol.*, 1998, **32**, 1058–1068.
- 46 P. Refait and J. M. R. Génin, Mechanisms of oxidation of Ni(II)-Fe(II) hydroxides in chloride-containing aqueous media: Role of the pyroaurite-type Ni-Fe hydroxychlorides, *Clay Miner.*, 1997, **32**, 597–613.
- 47 S. C. F. Auyeung, G. Denes, J. E. Greedan, D. R. Eaton and T. Birchall, A novel synthetic route to iron trihydroxide, $\text{Fe}(\text{OH})_3$ - characterization and magnetic-properties, *Inorg. Chem.*, 1984, **23**, 1513–1517.
- 48 J. Klausen, S. P. Trober, S. B. Haderlein and R. P. Schwarzenbach, Reduction of substituted nitrobenzenes by Fe(II) in aqueous mineral suspensions, *Environ. Sci. Technol.*, 1995, **29**, 2396–2404.
- 49 A. Neumann, T. B. Hofstetter, M. Skarpeli-Liati and R. P. Schwarzenbach, Reduction of polychlorinated ethanes and carbon tetrachloride by structural Fe(II) in smectites, *Environ. Sci. Technol.*, 2009, **43**, 4082–4089.
- 50 T. B. Hofstetter, A. Neumann and R. P. Schwarzenbach, Reduction of nitroaromatic compounds by Fe(II) species associated with iron-rich smectites, *Environ. Sci. Technol.*, 2006, **40**, 235–242.
- 51 D. M. C. Huminicki and F. C. Hawthorne, The crystal structure of nikischerite, $\text{Na Fe}^{2+}_6 \text{Al}_3(\text{SO}_4)_2 (\text{OH})_{18}$

- (H₂O)₁₂, a mineral of the shigaite group, *Can. Mineral.*, 2003, **41**, 79–82.
- 52 A. Voegelin, A. C. Scheinost, K. Buhmann, K. Barmettler and R. Kretzschmar, Slow formation and dissolution of Zn precipitates in soil - A combined column-transport and XAFS study, *Environ. Sci. Technol.*, 2002, **36**, 3749–3754.
- 53 O. Jacquat, A. Voegelin, A. Villard, M. A. Marcus and R. Kretzschmar, Formation of Zn-rich phyllosilicate, Zn-layered double hydroxide and hydrozincite in contaminated calcareous soils, *Geochim. Cosmochim. Acta*, 2008, **72**, 5037–5054.
- 54 L. Bhattacharya and E. J. Elzinga, A comparison of the solubility products of layered Me(II)-Al(III) hydroxides based on sorption studies with Ni(II), Zn(II), Co(II), Fe(II), and Mn(II), *Soil Syst.*, 2018, **2**, 20.
- 55 S. D. Kelly, B. Hesterberg and B. Ravel, in *Methods of Soil Analysis Part 5—Mineralogical Methods*, SSSA Book Ser. 5.5. SSSA, ed. L. R. D. A. L. Ulery, Madison, WI, 2008, pp. 387–463, DOI: 10.2136/sssabookser5.5.c14.
- 56 F. Trolard, G. Bourrie, M. Abdelmoula, P. Refait and F. Feder, Fougerite, a new mineral of the pyroaurite-iowaite group: Description and crystal structure, *Clays Clay Miner.*, 2007, **55**, 323–334.
- 57 H. Shirozu and S. W. Bailey, Chlorite Polytypism .III. crystal structure of an orthohexagonal iron chlorite, *Am. Mineral.*, 1965, **50**, 868–885.
- 58 R. Dähn, B. Baeyens and M. H. Bradbury, Investigation of the different binding edge sites for Zn on montmorillonite using P-EXAFS - The strong/weak site concept in the 2SPNE SC/CE sorption model, *Geochim. Cosmochim. Acta*, 2011, **75**, 5154–5168.
- 59 R. Dähn, A. M. Scheidegger, A. Manceau, M. L. Schlegel, B. Baeyens, M. H. Bradbury and M. Morales, Neoformation of Ni phyllosilicate upon Ni uptake on montmorillonite: A kinetics study by powder and polarized extended X-ray absorption fine structure spectroscopy, *Geochim. Cosmochim. Acta*, 2002, **66**, 2335–2347.
- 60 A. Manceau, D. Chateigner and W. P. Gates, Polarized EXAFS, distance-valence least-squares modeling (DVLS), and quantitative texture analysis approaches to the structural refinement of Garfield nontronite, *Phys. Chem. Miner.*, 1998, **25**, 347–365.
- 61 A. Manceau and M. L. Schlegel, Texture effect on polarized EXAFS amplitude, *Phys. Chem. Miner.*, 2001, **28**, 52–56.
- 62 A. Manceau, D. Bonnin, P. Kaiser and C. Fretigny, Polarized EXAFS of Biotite and Chlorite, *Phys. Chem. Miner.*, 1988, **16**, 180–185.
- 63 A. Manceau, Critical evaluation of the revised akdalaite model for ferrihydrite, *Am. Mineral.*, 2011, **96**, 521–533.
- 64 A. P. Zhukhlistov, Crystal structure of lepidocrocite FeO(OH) from the electron-diffractometry data, *Crystallogr. Rep.*, 2001, **46**, 730–733.
- 65 L. L. Baker and D. G. Strawn, Fe K-edge XAFS spectra of phyllosilicates of varying crystallinity, *Phys. Chem. Miner.*, 2012, **39**, 675–684.
- 66 A. Voegelin, R. Kaegi, J. Frommer, D. Vantelon and S. J. Hug, Effect of phosphate, silicate, and Ca on Fe(III)-precipitates formed in aerated Fe(II)- and As(III)-containing water studied by X-ray absorption spectroscopy, *Geochim. Cosmochim. Acta*, 2010, **74**, 164–186.
- 67 D. E. Latta, A. Neumann, W. A. P. J. Premaratne and M. M. Scherer, Fe(II)-Fe(III) electron transfer in a clay mineral with low Fe content, *ACS Earth and Space Chem*, 2017, **1**, 197–208.
- 68 A. C. Senn, R. Kaegi, S. J. Hug, J. G. Hering, S. Mangold and A. Voegelin, Composition and structure of Fe(III)-precipitates formed by Fe(II) oxidation in water at near-neutral pH: Interdependent effects of phosphate, silicate and Ca, *Geochim. Cosmochim. Acta*, 2015, **162**, 220–246.
- 69 R. M. Cornell and U. Schwertmann, *The iron oxides : structure, properties, reactions, occurrences, and uses*, Wiley-VCH, Weinheim, New York, 2nd, completely rev. and extended edn, 2003.
- 70 P. Gütllich, E. Bill and A. X. Trautwein, *Mossbauer Spectroscopy and Transition Metal Chemistry: Fundamentals and Applications*, Springer, 2011.
- 71 L. T. Kuhn, K. Lefmann, C. R. H. Bahl, S. N. Ancona, P. A. Lindgard, C. Frandsen, D. E. Madsen and S. Morup, Neutron study of magnetic excitations in 8-nm alpha-Fe₂O₃ nanoparticles, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 2006, **74**, 184406.
- 72 C. M. Chen, R. Kukkadapu and D. L. Sparks, Influence of coprecipitated organic matter on Fe-(aq)(2+)-catalyzed transformation of ferrihydrite: implications for carbon dynamics, *Environ. Sci. Technol.*, 2015, **49**, 10927–10936.
- 73 A. M. Jones, R. N. Collins, J. Rose and T. D. Waite, The effect of silica and natural organic matter on the Fe(II)-catalysed transformation and reactivity of Fe(III) minerals, *Geochim. Cosmochim. Acta*, 2009, **73**, 4409–4422.
- 74 C. M. Hansel, S. G. Benner and S. Fendorf, Competing Fe(II)-induced mineralization pathways of ferrihydrite, *Environ. Sci. Technol.*, 2005, **39**, 7147–7153.
- 75 C. M. Hansel, S. G. Benner, J. Neiss, A. Dohnalkova, R. K. Kukkadapu and S. Fendorf, Secondary mineralization pathways induced by dissimilatory iron reduction of ferrihydrite under advective flow, *Geochim. Cosmochim. Acta*, 2003, **67**, 2977–2992.
- 76 J. W. Gruner, The composition and structure of minnesotaite, a common iron silicate in iron formations, *Am. Mineral.*, 1944, **29**, 363–372.
- 77 C. Mikutta, X-ray absorption spectroscopy study on the effect of hydroxybenzoic acids on the formation and structure of ferrihydrite, *Geochim. Cosmochim. Acta*, 2011, **75**, 5122–5139.

