


Photochemical & Photobiological Sciences

An international journal

rsc.li/pps

Volume 19
Number 8
August 2020
Pages 989-1106



ISSN 1474-905X





Cite this: *Photochem. Photobiol. Sci.*, 2020, **19**, 1001

Anti- and pro-oxidative mechanisms comparing the macular carotenoids zeaxanthin and lutein with other dietary carotenoids – a singlet oxygen, free-radical *in vitro* and *ex vivo* study

Fritz Boehm,^a Ruth Edge ^b and T. George Truscott^{*c}

The interactions of dietary carotenoids, and particularly the xanthophylls in the macula, with singlet oxygen and three different oxy-radicals, (hydroxyl radical, nitrogen dioxide and the superoxide radical anion) are compared using pulsed laser and γ -techniques. The results give possible molecular mechanisms for the switch from anti-oxidant (protection) by carotenoids to pro-oxidant (damage) by carotenoids. The participation of oxygen in radical mechanisms in the presence of different carotenoids is compared for the different radicals. It is shown that the mechanistic role of oxygen differs very significantly for anti-/pro-oxidation by hydroxyl radicals when compared to nitrogen dioxide. Lutein was found to be an extremely good cell protector against hydroxyl radicals at all oxygen concentrations, including under physiological conditions.

Received 30th March 2020,
Accepted 17th June 2020

DOI: 10.1039/d0pp00120a

rsc.li/pps

Introduction

As well as being important anti-oxidants in the eye, carotenoids (CAR) are of significance in photosynthesis, the skin, and in several other aspects of human health. The anti-oxidant processes mainly concern quenching of singlet oxygen ($^1\text{O}_2$) and/or free radicals. The quenching of $^1\text{O}_2$ by dietary carotenoids in 'simple' solvents and heterogeneous environments from micelles and liposomes to cells has already been extensively reported,¹ and it is well established most quench $^1\text{O}_2$ extremely efficiently, near the diffusion limit, (except lutein (LUT), with 10 conjugated double bonds, being typically 50% less efficient). However, some of this work deserves further comment, as follows: Ogilby and co-workers have thrown doubt on the relevance of such quenching – they were able to detect $^1\text{O}_2$ in cells² using individual HeLa mammalian cells and microscope-based time-resolved 1275 nm luminescence, but they could observe no quenching of the $^1\text{O}_2$ by β -carotene (β -CAR). However, Telfer and co-workers show carotenoids quench some of the $^1\text{O}_2$ in cells, but not all, due to the distance of the carotenoids from the source of the $^1\text{O}_2$.³ This work, in photosynthetic systems, suggests that in a specific

biological environment, quenching will only be efficient when the carotenoid is sufficiently close to the source of the $^1\text{O}_2$. Virtually all studies have concerned the $^1\text{O}_2$ quenching by carotenoids as non-aggregated, monomeric species. However, when aggregated, quenching by the xanthophylls (LUT and zeaxanthin (ZEA)) and probably all carotenoids is markedly reduced. The carotenoid concentration in the fovea of the eye approaches 1 mM, and the ratio of LUT to ZEA to *meso*-ZEA is 1:1:1.⁴ So, the extension of singlet oxygen quenching to aggregated carotenoids should also be further considered. Clearly, there is more to learn about the interaction of β -CAR with $^1\text{O}_2$ in the *ex vivo* and *in vivo* situation.

While the overall process of $^1\text{O}_2$ quenching simply converts the excess energy to heat *via* the carotenoids lowest triplet state, the reaction of carotenoids, with free radicals is much more complex, partly because individual radicals behave differently from each other and also because of a likely role of the oxygen concentration – this will also depend on the type of radical.

In this work, we review and give new results showing how three distinct radicals, the hydroxyl radical, nitrogen dioxide and the superoxide radical anion (OH^\cdot , NO_2^\cdot and $\text{O}_2^{\cdot-}$ respectively) exhibit totally different chemistry in their interaction with carotenoids. Often this is 'over simplified' by just referring to carotenoid quenching of Reactive Oxygen Species (ROS). Here, we compare OH^\cdot , NO_2^\cdot and $\text{O}_2^{\cdot-}$ reactivity with LUT, ZEA and other carotenoids, especially lycopene (LYC) – there are claims that LYC has a role in macular protection despite not accumulating in the macula.^{5,6}

^aPhotobiology Research, Internationales Handelszentrum (IHZ), Friedrichstraße 95, 10117 Berlin, Germany

^bDalton Cumbrian Facility, Westlakes Science Park, The University of Manchester, Cumbria, CA24 3HA, UK

^cSchool of Chemical and Physical Sciences, Keele University, Staffordshire, ST5 5BG, UK. E-mail: t.g.truscott@keele.ac.uk




Experimental

Radiation processes

The radiation processes are well established.⁷ The $\text{NO}_2\cdot$ radical is generated, in aqueous solutions, using 355 nm pulsed laser excitation of 1-nitronaphthalene (NN). This produces the NN triplet state (^3NN) which can subsequently react with sodium nitrite, to yield the radical anion of nitronaphthalene ($\text{NN}^{\cdot-}$) and the nitrogen dioxide radical ($\text{NO}_2\cdot$).^{8–10}

The steady-state γ -radiolysis was performed in a Model 812 cobalt-60 irradiator (Foss Therapy Services Inc.) using a turntable system rotating at 12 rpm with an average absorbed dose rate of around 90–120 Gy min^{-1} . All irradiations were carried out at room temperature and solutions were saturated, as appropriate, with research-grade nitrous oxide, argon, oxygen or mixtures. Dose rates were determined using a Radcal Corporation Accu-Dose+ base unit equipped with a 10×6 –0.18 ion chamber (calibrated annually by PHE to traceable national standards) and were checked against standard Fricke dosimetry and found to be accurate to $\pm 5\%$.

The radiolysis products of water⁷ can be simplified to the generation of main radical products $\text{OH}\cdot$ and the solvated electron, e_{aq}^-



(the $\text{H}\cdot$ being a minor product).

At all oxygen concentrations used the e_{aq}^- is totally converted to $\text{O}_2^{\cdot-}$



so, the concentration of $\text{O}_2^{\cdot-}$ produced is independent of oxygen concentration.

The cell types used for the *ex vivo* studies have also been described previously.^{8–10}

The individual carotenoids were gifts from Hoffmann-la-Roche (better than 99.9% *via* HPLC). For the *ex vivo* studies, dietary supplements were used, and the claimed amount checked *via* absorption spectroscopy. The duration of the diet (between 4 days and 2 weeks) was varied but maintaining a constant total amount of carotenoid, and this had no significant effect on our results.


The SOD (bovine) was supplied by Sigma-Aldrich as 75 000 units and was used at 30 000, units in a cell suspension volume maintained at 700 μl .

The sources of all other chemicals have been described previously.⁸

While there may be a link between dietary intake and serum levels of specific carotenoids this is complex and varies from one individual to another. We simply used a high dose of each supplement to ensure maximum uptake for each carotenoid, as described previously for lycopene.¹⁰ No attempt was made to determine the amount of carotenoid nor the distribution in each cell. We examined the effect of changing the rate of dietary intake – *i.e.* shorter times at higher concentrations. This made no difference to the results obtained.

Results and discussion

Free radicals, of course, are characterized by an unpaired electron. When a free radical interacts with a carotenoid, several possible modes of reaction can occur, and these depend mainly on the nature of the free radical. The situation is, therefore, much more complex than the quenching of $^1\text{O}_2$ by carotenoids. Many studies of the generation and reactions of carotenoid radical cations, and anions, have been reported in several reviews.^{11–13} One of the most well studied species is the carotenoid radical cation obtained *via* abstraction of an electron from the carotenoid by a strongly oxidising free radical.



Typical examples of such oxidising free radicals that lead to electron transfer are chlorinated peroxy radicals, such as $\text{CCl}_3\text{O}_2\cdot$, nitrogen dioxide ($\text{NO}_2\cdot$), arylperoxy radicals, dibromide radical anion ($\text{Br}_2^{\cdot-}$) and sulfonyl radicals ($\text{RSO}_2\cdot$). However, as we will discuss below, the hydroxyl radical ($\text{OH}\cdot$), even though it is a strongly oxidising radical, mainly adds to the carotenoid rather than producing the carotenoid radical cation. Furthermore, radicals, such as $\text{CCl}_3\text{O}_2\cdot$, and possibly $\text{NO}_2\cdot$, may give both the radical cation and also add to the carotenoid. Though, following extensive studies, only electron transfer has been previously observed for the reaction of β -CAR with $\text{NO}_2\cdot$.¹⁴

Two significant results from pulse radiolysis and laser flash photolysis show (i) the carotenoid radicals are themselves strongly oxidising radicals which can cause damage by oxidising bio-substrates and (ii) ascorbic acid (and other reducing species) can remove (quench) such carotenoid radical cations and possibly ameliorate the potential deleterious problems associated with the carotenoid radical cation production. See, for example ref. 15 and 16.

It has been shown that, for heavy smokers, a high concentration of β -CAR can have a surprisingly damaging effect for lung cancer,¹⁷ and our speculation was that this may, in part, be due to smoke-based free radicals (*e.g.* $\text{NO}_2\cdot$) reacting with β -CAR to generate the β -CAR radical cation in the lung, which can then damage biomolecules.¹⁰ Smokers have low levels of ascorbic acid,¹⁸ and so there is insufficient to efficiently quench any β -CAR radical cation generated *via* $\text{NO}_2\cdot$.

Results from Skibsted and co-workers^{19,20} are consistent with the above, showing reactions of β -CAR radical cation with tyrosine and tryptophan, regenerating the parent β -CAR but oxidising the amino acids. Interestingly, Skibsted used pH conditions where the redox potentials (the standard reduction potentials) were the same for tyrosine and tryptophan (the reduction potential for β -CAR radical cation is independent of pH in the region studied). These workers found that tyrosine reacted an order of magnitude faster than tryptophan and speculate that this may account for tyrosine, rather than tryptophan, as the protein moiety reacting with β -CAR in the protective mechanism, which operates in the photosynthetic reaction center. As Skibsted points out, the driving force in these reac-



tions depends on the “local” pH and in proteins the reverse reaction between a tyrosine radical and β -CAR may also be important.

Carotenoids and NO_2^{\cdot}

In early work we showed a significant protection of human cells against pulsed-laser generated NO_2^{\cdot} by LYC.⁸ In this work we compared the percentage of dead cells (obtained *via* eosin staining techniques) with and without LYC supplementation under atmospheric conditions. The ratio was reported as a protection factor with values between 8.2 and 8.6. A somewhat lower protection was observed for β -CAR supplementation (3.0–3.5). However, these early studies involved the use of ‘water-solubilised’ carotenoids (rather than the carotenoid itself) and an *in vitro* procedure in which the ‘soluble’ carotenoids were simply added to the blood lymphocyte suspensions prior to the exposure to the NO_2^{\cdot} . That is, no dietary route to carotenoid supplementation was used. Nevertheless, these studies did show that LYC can be an efficient protective species against NO_2^{\cdot} , a major toxic component of polluted air.

Subsequently,⁹ we reported a synergistic cell protection against NO_2^{\cdot} -induced cell kill when the actual diet was supplemented with β -CAR and with and without dietary vitamin C and E (typically 1000 mg day⁻¹ and 800 mg day⁻¹ respectively). The protection factor increasing from 2.0, 1.8 and 1.2 for β -CAR, vitamin E and vitamin C individually to 10 when all 3 were co-supplemented.

We now report similar measurements with LYC instead of β -CAR. Once again, control experiments with supplementation with only vitamin C and only vitamin E showed protection factors of 1.8 and 1.2 while with LYC (30 mg day⁻¹ for 3 weeks) plus these two vitamins the protection factor was greatly increased to 21. Essentially, the same synergistic observations, in a non-biological system, have been reported by Liu *et al.*²¹ We have also previously reported similar synergism when the specific amounts of lycopene were replaced by either 2 weeks supplementation with tomato soup or with pre-boiled tomato juice.¹⁰

To understand the mechanism of such synergism it is noteworthy that the beneficial quenching of the damaging NO_2^{\cdot} radical by a dietary carotenoid also generates the radical cation of the carotenoid, as described above, and this is a strongly oxidising species itself which can oxidise biological moieties such as tryptophan, tyrosine and cysteine^{19,20} leading to deleterious membrane protein damage and subsequent disease. However, we have previously shown in both methanol and micellar environments, that vitamin C quenches carotenoid radical cations^{15,16} and we suggest that such a process between vitamin C and the carotenoid radical cation could account for the observed synergistic effects. Furthermore, it could be related to the much discussed and surprising results showing β -CAR enhances lung cancer incidence in smokers.¹⁷ Of course, smokers are very low in vitamin C in the lungs!

The above does depend on the NO_2^{\cdot} reacting with carotenoids substantially *via* electron transfer, as has been reported

by Everett *et al.*¹⁴ However, no such information is available for LYC.

In new work we have studied the cell killing by NO_2^{\cdot} as a function of the oxygen concentration. As discussed below in detail, we would expect no effect on carotenoid radical cation/vitamin C/E processes of oxygen concentration. If radical addition processes arise then addition of oxygen (to give, possibly, a damaging peroxy radical) will be significant and the above mechanisms involving radical cations would not apply.

These new results show a cell kill of 53.7%, 54.2% and 57.8% at 0%, 20% and 100% oxygen, respectively, for the unprotected cells (*i.e.* no LYC, vitamin C or E supplementation).

For the protected cells (with the anti-oxidant supplement) the corresponding cell kill was 3.2%, 3.1% and 6.2%.

These lead to protection factors of 16.8 and 17.5 for 0% and 21% oxygen – that is no effect of the oxygen. For the 100% oxygen the protection factor is somewhat reduced to 9.3.

Our overall conclusion is that oxygen has played no role in normal physiological conditions and hence the value of LYC (and possibly β -CAR²²) is worthwhile considering as a means of reduction damage (possibly lung damage) due the environmental NO_2^{\cdot} in non-smokers. Of course, other processes involving other smoke-related radicals can also be related to the increased lung cancer in the presence of high β -CAR diets¹⁷ so such a beneficial value (for all dietary carotenoids) needs careful assessment.

Carotenoids and OH^{\cdot} – the role of oxygen

Here we compare the macular carotenoids, LUT and ZEA with previous data on LYC (with re-interpretation of some of those results), β -CAR and astaxanthin (ASTA). In this series of experiments, we used commercial carotenoid supplements. We avoided commercial supplement with carotenoid mixtures even though many people use such mixtures (*e.g.* LUT plus ZEA for AMD). We do not know the concentration of a carotenoid in each cell nor its position (near the aqueous interface or totally embedded in the membrane for example). Instead, our aim was to compare the different carotenoids where the cell loading is as high as can be achieved in normal use by the public – *i.e.* using amount of carotenoids well above that normally used (typically 70–100 mg day⁻¹ for 2 weeks or variations on this as mentioned above – all combinations studied gave the same results) so that effectively all cells have the maximum carotenoid loading under normal conditions. The numerical protection factors we report may not be of significance. However, the comparisons in these factors, as a function of oxygen concentration, for the 5 carotenoids studied are of interest. Indeed, our comparative data may suggest one possible explanation as to why the macular chooses ZEA and LUT for best protection.

The first carotenoid we choose to study the protection of human cells against damage by OH^{\cdot} (see eqn (1) and (2) above) generated *via* γ -radiation was LYC¹¹ – chosen partly because it has been established it is the carotenoid which is the strongest reducing agent (*i.e.* the most easily oxidised)¹



and also because it may be important with respect to protection against AMD despite not accumulating in the macular.^{5,6}

In air we reported a reduction in cell kill due to dietary LYC (typically about 100 mg day⁻¹ over 2 weeks, but other time/dose combinations gave the same results and comparisons with added vitamin C and E also showed no effect on LYC protection against the cell kill) from near 44% in the absence of lycopene to near only 9% with the supplementation – this is a protection factor (PF) of near 5 ± 1.3. While this is a significant protection, we saw a most dramatic effect of high oxygen concentration on the PF as shown in Fig. 1.

At 100% oxygen there was no protection at all while at 0% oxygen the PF was over 50. Thus, LYC gave near total protection against damage by OH[•] in the absence of oxygen.

Fig. 1 shows that in the presence of the LYC supplement, increasing the oxygen concentration reduces the protection of the cells. That is, despite the OH[•] concentration remaining constant between 0 and 100% oxygen, and the O₂^{•-} concentration between 5 and 100%, there is a huge change in the PF. Clearly, a key finding is that in the presence of the LYC supplement the PF is affected, to a large extent, by the concentration of oxygen itself rather than by the free radical concentrations.

It is instructive to compare the present results using γ -radiation with those we have previously published^{8–10} for nitrogen dioxide (NO₂[•]), discussed above. Using similar lymphoid cell procedures, we obtained a PF of about 17 in the laser studies – that is, considerably more protection than seen in the γ -radiation work (PF near 5 in air-saturated conditions). While the laser study generated virtually only oxidising NO₂[•] radicals the γ -radiation generates, under the conditions used in the presence of oxygen, both OH[•] radicals and the superoxide radical anion (discussed below). This comparison shows the LYC supplement protects against damage due to the oxidising NO₂[•] radicals markedly better than against the OH[•] radical (the superoxide reactivity is discussed below).






Fig. 1 Mean PF against oxygen percentage for lymphocyte cells irradiated with 2000 Gy dose; inset shows the percentage of dead cells under these conditions.

In a totally non-biological environment (non-cellular and non-aqueous) key work by Burton and Ingold²³ investigated the effect of oxygen concentration on processes involving lipid peroxidation leading to free radicals and oxygen radical addition reactions. In this study, adducts between the carotenoid and peroxy radicals were also invoked to explain a switch from anti- to pro-oxidant behaviour of β -CAR as the oxygen concentration increased in their system.

One mechanism discussed^{1,23} to explain the effect of O₂ concentration on the cell protection by the LYC supplement is related to the ability of O₂ to add to neutral radicals to produce reactive peroxy radicals. There is much evidence that the strongly oxidising OH[•] forms adducts with most substrates so that LYC, OH[•] neutral radical adduct (LYC-OH[•]) is likely to be a significant product in our cellular studies.^{24–26} Another possible product is the neutral LYC radical (LYC[•]) formed *via* hydrogen atom abstraction.²⁷ In the presence of oxygen, both radicals may add molecular oxygen to form reactive peroxy radicals leading to species such as LYC-OH-OO[•] as shown in eqn (4)–(6) below or LYC-OO[•] in the case of hydrogen abstraction followed by oxygen addition. We suggested, such peroxy radicals are formed in increasing concentration as the oxygen concentration increases and proposed these killed the lymphoid cells. Our cell study showed a near linear relationship between oxygen concentration and cell kill (Fig. 1 inset) and the following postulated mechanism:



is consistent with such a linear relationship.

However, other possible mechanisms involving oxygen concentration are possible.^{28,29} Our new results concern the carotenoids which accumulate in the macula – ZEA and LUT and comparative results are now presented on the protection against OH[•] by these xanthophylls. In addition, we present results on another xanthophyll – ASTA and some results on the ‘parent’ carotenoid, β -CAR.

Typical results are given in Fig. 2 showing comparative plots of cell kill and protection factors as a function of oxygen percentage.

The effect of increasing oxygen concentration reducing the efficiency of cell protection by LYC, discussed above, is also observed with the 4 new carotenoids studied. But less so with the xanthophylls and especially so for LUT, which is still an effective cell protector even at 100% oxygen.

The high protective ability of lutein compared to the 4 other carotenoids studied is also evident from Fig. 3 – the corresponding bar charts for air and oxygen.

Two significant trends due to the presence of oxygen are clear. Now the hydrocarbons (LYC and β -CAR) show the lowest PF compared to the xanthophylls (ASTA, ZEA and LUT) and, LUT shows the highest PF, especially at high oxygen concentrations. In fact, the efficient cell protection by lutein shows,






Fig. 2 (a) Mean PF for LYC (red), β-CAR (orange), ZEA (cyan), ASTA (blue) and lutein (green) carotenoids against oxygen percentage for lymphocyte cells irradiated with 2000 Gy dose. (b) Shows the corresponding percentages of dead cells under these conditions.




Fig. 3 (a) Dead cell count compared for 5 carotenoids against the control at 100% oxygen. (b) Dead cell count compared for 5 carotenoids against the control in air; insets shows the PFs under the same conditions.


more-or-less, rather little change from 0–100% oxygen (37; 33; 26; 27; and 25 for oxygen percentages of 0; 10; 21; 60; and 100%).

So, overall, our work with OH[•] in the presence of oxygen suggests that while lutein is the poorest protector against one ROS – singlet oxygen, it is significantly the best against radicals which may arise in oxygenated conditions (as in the eye).


Carotenoids and O₂^{•-} – the role of oxygen and SOD

The role and effectiveness of the first-line defense antioxidant, superoxide dismutase (SOD), is important and indispensable as an antioxidant, especially against O₂^{•-}, which is perpetually generated in normal body metabolism.

The product arising from carotenoids reacting with O₂^{•-} is not a primary concern of this work and is not well understood. Generation of the carotenoid radical cation has been suggested:³⁰



While others,³¹ using density function theory, show O₂^{•-} acts as an electron donor species to generate the carotenoid radical anions:



In our work on LYC protection⁹ we were able to calculate the theoretical concentration of OH[•], O₂^{•-} and the solvated electron (e⁻_{aq}) and this showed a significant cell kill by O₂^{•-} and OH[•] but rather little kill by the e⁻_{aq}. (5.6%, 8.75% and 2% respectively). As discussed above we report a significant and differing effect of oxygen concentration on the PFs of the 5 carotenoids studied and explained this in terms of the neutral radical adducts, CAR–OH[•], adding oxygen. Clearly, the effect of oxygen concentration on the cell kill by O₂^{•-} could be important.

Furthermore, an aspect of our early work⁹ which was not previously presented was that at 100% oxygen the LYC gives no cell protection whatsoever – that is, no protection against OH[•] nor O₂^{•-}. While at low oxygen levels (5%) there is only 1.3% cell kill in the presence of the LYC supplement. This implies LYC is protecting against OH[•] and O₂^{•-} at this low oxygen concentration.



In order to gain more information on the effects of $O_2^{\cdot-}$ we present results on the comparative effects, with and without superoxide dismutase (SOD) for LYC, ZEA and β -CAR. In particular, we report the results on protection factors with and without quenching of $O_2^{\cdot-}$ by SOD.

For LYC we used air, 60% and 100% oxygen, with previously reported PFs of 5, 2 and 1 (*i.e.* no protection at all at 100% oxygen). As can be easily seen from Fig. 4 there is no effect of oxygen concentration on the protection factor or cell kill count for just SOD, in the absence of LYC, (red plot).

As can be seen the PF observed for the SOD was constant at 2.6 for 21%, 60% and 100% oxygen – showing no effect of oxygen concentration on the protection of the cells by SOD (Fig. 4b). It is important to note the SOD was simply added to the cell suspension and so was predominantly localised outside the cells while the carotenoids (lycopene in this case) are present in the cells, *i.e.* in the membranes. So, a comparison between the individual protectors (SOD and lycopene here) is not likely to be useful. However, the same procedure was adopted for all the carotenoids studied so the comparison of the different carotenoids is worthwhile.




Fig. 4 (a) Mean percentages of dead cells for LYC with (green) and without (red) SOD, and for SOD alone (SOD control, blue), against oxygen percentage for lymphocyte cells irradiated with 2000 Gy dose. (b) PFs for SOD alone (red), LYC alone (black) and for both SOD and LYC compared to the main control (green) and also for both SOD and LYC compared to the SOD control (blue).

Of course, the combination of LYC and SOD gives the best protection – ranging from 14 to 3 for air to 100% oxygen. In good agreement with the protection for the individual antioxidants. The data also shows (compare the black and blue plots) no effect of the presence of SOD on the protection by LYC at any of the oxygen concentrations studied.

The corresponding results for ZEA are summarised in the Fig. 5.

We obtained similar results with the other carotenoids studied. Showing no effect of SOD on the carotenoid cell protection with respect to oxygen concentration.

We compare the PF for all carotenoids studied against $O_2^{\cdot-}$ and OH^{\cdot} (combined) to that of SOD (Table 1 below).




Fig. 5 (a) Mean percentages of dead cells for ZEA with (green) and without (red) SOD, and for SOD alone (SOD control, blue), against oxygen percentage for lymphocyte cells irradiated with 2000 Gy dose. (b) PFs for SOD alone (red), ZEA alone (black) and for both SOD and ZEA compared to the main control (green) and also for both SOD and ZEA compared to the SOD control (blue).

Table 1 Protection factors (PF \pm 10%) in air and 100% oxygen

Carotenoid	% O ₂	PF	% O ₂	PF
LYC	21	4.9	100	1.0
β -CAR	21	5.0	100	1.7
ASTA	21	10.8	100	4.8
ZEA	21	8.7	100	3.3
LUT	21	26.0	100	25.0
SOD	21	2.6	100	2.6



As can be seen under near physiological conditions LUT shows the highest PF by a significant amount, at all oxygen concentrations. As noted above, the important comparisons are between the individual carotenoids PFs against $O_2^{\cdot-} + OH^{\cdot}$ themselves rather than each with SOD. Possibly this reflects an important property of LUT – to protect efficiently against a wide range of oxidising radicals including OH^{\cdot} and $O_2^{\cdot-}$ at all oxygen concentrations, whereas SOD only protects efficiently against $O_2^{\cdot-}$. Possibly the outcome of the interaction of carotenoids, including LUT, with $O_2^{\cdot-}$ leads to other potentially damaging products such as carotenoid radicals,^{30,31} with no such potential damage from SOD.

Conclusions

While the role and relative efficiencies of the dietary carotenoids to quench singlet oxygen in solvents and some heterogeneous environments are well established this process is less clear in cells. In a single mammalian HeLa cell, 1O_2 was detected, but no quenching of the 1O_2 by β -CAR was observed, although such quenching has been reported in photosynthetic systems. A further complication, especially for the macular pigment, ZEA, is the marked reduction in 1O_2 quenching due to aggregation – much less effects of aggregation for LUT being reported.

For ROS other than 1O_2 we have shown the three radicals NO_2^{\cdot} , OH^{\cdot} and $O_2^{\cdot-}$ behave quite differently in reactivity with carotenoids.

The concentration of oxygen is shown to be pivotal in understanding the protection of cells against OH^{\cdot} radicals. Particularly the effectiveness of the macular pigment LUT (compared to other dietary carotenoids including ZEA) to protect human cells against OH^{\cdot} damage at physiological (and higher) oxygen concentrations is shown.

Neither NO_2^{\cdot} and $O_2^{\cdot-}$ showed any marked effects of oxygen concentration. However, the NO_2^{\cdot} radical was shown to be quenched by dietary carotenoids but this quenching leads to the production of carotenoid radical cations and pointed to the value of reducing agents, especially ascorbic acid, to prevent a switch to damaging pro-oxidation. The possible relevance of this to the increased lung cancer risk of heavy smokers by β -carotene was invoked and extended to a possible benefit for LYC/vitamin C mixtures.

Of course, SOD is a key *in vivo* protector against the $O_2^{\cdot-}$ radicals. However, our new results indicate that LUT protects against $O_2^{\cdot-}$ (by an order of magnitude) more effectively than SOD. While our *ex vivo* results are just a model system, the efficient human cell protection provided by LUT at all oxygen concentrations, not provided by other carotenoids, may indicate one reason why it is LUT that accumulates in the macula.

Overall, there is still much to learn about a possible anti-pro-oxidation switch for dietary carotenoid interactions with individual ROS components and this requirement is often hidden by ‘simple’ statements concerning dietary carotenoids quenching ROS.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We acknowledge the support of Keele University and The University of Manchester's Dalton Cumbrian Facility (DCF), a partner in the National Nuclear User Facility, the EPSRC UK National Ion Beam Centre and the Henry Royce Institute.

We would also like to acknowledge Dr Joan Roberts for the invitation to present this work at the World Congress on Light and Life (17th Congress of the International Union of Photobiology and 18th Congress of the European Society for Photobiology), 25-30 August 2019, Barcelona, Spain.

References

- 1 R. Edge and T. G. Truscott, *Antioxidants*, 2018, 7, 1.
- 2 G. N. Bosio, T. Breitenbach, J. Parisi, M. Reigosa, F. H. Blaikie, B. W. Pedersen, E. F. F. Silva, D. O. Martire and P. R. Ogilby, *J. Am. Chem. Soc.*, 2013, 135, 272.
- 3 A. Telfer, *Plant Cell Physiol.*, 2014, 55, 1216.
- 4 R. A. Bone, J. T. Landrum, L. M. Friedes, C. M. Gomez, M. D. Kilburn, E. Menendez, I. Vidal and W. Wang, *Exp. Eye Res.*, 1997, 64, 211.
- 5 N. Cardinault, J.-H. Abalain, B. Sairafi, C. Coudray, P. Grolier, M. Rambeau and J.-C. Carré, *Clin. Chim. Acta*, 2005, 357, 34.
- 6 J. A. Mares-Perlman, W. Brady, R. Klein, B. Klein, P. Bowen, M. Stacewiczapuntzakis, *et al.*, *Arch. Ophthalmol.*, 1995, 113, 1518.
- 7 R. V. Bensasson, E. J. Land and T. G. Truscott, *Excited states in biology and medicine*, Oxford University Press, UK, 1993, ch. 1-4.
- 8 F. Boehm, J. Tinkler and T. G. Truscott, *Nat. Med.*, 1995, 1, 98.
- 9 F. Boehm, R. Edge, D. J. McGarvey and T. G. Truscott, *FEBS Lett.*, 1998, 436, 387.
- 10 F. Böhm, R. Edge, M. Burke and T. G. Truscott, *J. Photochem. Photobiol., B*, 2001, 64, 176.
- 11 F. Boehm, R. Edge, T. G. Truscott and C. Witt, *FEBS Lett.*, 2016, 590, 1086.
- 12 F. Böhm, R. Edge and T. G. Truscott, *Mol. Nutr. Food Res.*, 2012, 56, 205.
- 13 R. Alvarez, B. Vaz, H. Gronemayer and A. R. de Lera, *Chem. Rev.*, 2014, 114, 1.
- 14 S. A. Everett, M. F. Dennis, K. B. Patel, S. Maddix, S. C. Kundu and R. L. Willson, *Am. Soc. Biochem. Mol. Biol.*, 1996, 271, 3988.
- 15 F. Böhm, R. Edge, E. J. Land, D. J. McGarvey and T. G. Truscott, *J. Am. Chem. Soc.*, 1997, 119, 621.
- 16 M. Burke, R. Edge, E. J. Land and T. G. Truscott, *J. Photochem. Photobiol., B*, 2001, 60, 1.



- 17 The α -Tocopherol, β -carotene Cancer Prevention Study Group, *N. Engl. J. Med.*, 1996, **330**, 1029.
- 18 X. D. Xang and R. M. Russell, *Nutr. Res.*, 1999, **57**, 263.
- 19 H. Cheng, R.-M. Han, M.-K. Lyu, J.-P. Zhang and L. H. Skibsted, *J. Phys. Chem. B*, 2015, **119**, 6603.
- 20 H.-T. Chang, H. Cheng, R.-M. Han, P. Wang, J.-P. Zhang and L. H. Skibsted, *J. Agric. Food Chem.*, 2017, **65**, 908.
- 21 D. Liu, J. Shi, A. C. Ibarra, Y. Kakuda and S. J. Xue, *Food Sci. Technol.*, 2008, **41**, 1344.
- 22 R. Góralczyk, *Nutr. Cancer*, 2009, **61**, 767.
- 23 G. W. Burton and K. U. Ingold, *Science*, 1984, **224**, 569.
- 24 S. Steenken, *Chem. Rev.*, 1989, **89**, 503–520.
- 25 C.-H. Chen, R.-H. Han, R. Liang, L.-M. Fu, P. Wang, A. Xi-Cheng, J.-P. Zhang and L. H. Skibsted, *J. Phys. Chem. B*, 2011, **115**, 2082.
- 26 A. Mortensen, L. H. Skibsted and T. G. Truscott, *Arch. Biochem. Biophys.*, 2001, **385**, 13.
- 27 A. El-Agamey and D. J. McGarvey, *Free Radical Res.*, 2007, **41**, 295.
- 28 A. El-Agamey, A. Cantrell, E. J. Land, D. J. McGarvey and T. G. Truscott, *Photochem. Photobiol. Sci.*, 2004, **3**, 802.
- 29 A. El-Agamey and D. J. McGarvey, *Free Radical Biol. Med.*, 2016, **90**, 75.
- 30 T. G. Truscott, *J. Photochem. Photobiol., B*, 1996, **35**, 233.
- 31 A. Galano, R. Vargast and A. Martinez, *Phys. Chem. Chem. Phys.*, 2010, **12**, 193.

