

Cite this: *J. Mater. Chem. C*,
2024, 12, 1640

Interaction strength in molecular junctions consisting of π -stacked antiaromatic molecules†

Shintaro Fujii,^a Ryoya Tomida,^b Aoshi Yamane,^a Kazuki Nabeyama,^a
Harunari Ohkura,^a Hiroshi Shinokubo^b and Tomoaki Nishino^a

Elucidation of intermolecular interactions is essential for understanding charge transport properties in organic materials and organic electronic devices. While interactions between aromatic molecules have been extensively studied, little is known about interactions between antiaromatic molecules. Theoretical considerations predict that when two antiaromatic porphyrin molecules are superimposed, the antiaromatic molecules should be stabilized by attractive intermolecular interactions. In this study, we used atomic force microscopy to evaluate intermolecular interactions originating from the π - π stacking of the two antiaromatic porphyrin moieties at the molecular level. We evaluated substantial interaction between the antiaromatic porphyrin moieties as an adhesion force of the antiaromatic porphyrin molecular pairs using force spectroscopy. The present result supports the formation of π - π stacking due to the attractive interaction between the two antiaromatic π -systems at the single-molecule level.

Received 13th November 2023,
Accepted 19th December 2023

DOI: 10.1039/d3tc04166b

rsc.li/materials-c

Introduction

The importance of intermolecular π - π stacking interactions in enhancing charge transport has been widely studied in organic electronics.^{1,2} Effective overlap of molecular orbitals along the π - π stacking has contributed to the enhancement of charge transport in organic electronic and bioelectronic materials. Recent developments in molecular electronics have made it possible to probe the π - π stacking at the single molecule level, revealing unique charge transport properties² such as quantum interference effects,^{3,4} length-independent charge transport,⁵ and high structural responsiveness to the external electric field.⁶ To understand the charge transport properties of π - π stacking, it is of crucial importance to clarify the interactions between π -stacked molecules. Indeed, extensive studies devoted to the interactions between aromatic molecules have shed light on their effects on electron transport. In terms of the π - π stacking interactions, recent studies have shown that antiaromatic compounds, characterized by planar rings of sp^2 carbon atoms sharing multiples of four π -electrons, form unique π - π stacking structures in solution and in the solid state that spatially delocalize the π -electrons of the two antiaromatic

systems and that the molecular pairs of antiaromatics exhibit stacked-ring aromaticity.^{7,8} Moreover, excellent charge transport properties were found for such antiaromatic compounds with a small HOMO-LUMO gap.⁹ Detailed knowledge of the π - π interactions is required to fully exploit the superior properties of the antiaromatics. However, unlike the aromatic counterpart, little is known about interactions between antiaromatic molecules.^{7,10,11} In this study, atomic force microscopy (AFM) was used to assess the intermolecular interactions between antiaromatic systems, norcorrole Ni(II) (denoted as Ni(nor)). We found substantial interaction between two antiaromatic Ni(nor) systems using force spectroscopy. This result supports the formation of π - π stacking due to the attractive interaction between the two antiaromatic π -systems. The present work reveals the intermolecular interactions of the antiaromatic systems at the single-molecule level and provides a better understanding of the interactions that make antiaromatic π -stacks suitable for building blocks in organic electronics.

Results and discussion

The antiaromatic intermolecular interaction was evaluated by measuring the interaction force between antiaromatic molecules using AFM. We used a molecular matrix of a hexanethiol (C6) monolayer as a host to support and isolate single dodecanethiol (C12) chains appended with an antiaromatic moiety of Ni(nor) as the tail group (Fig. 1a). Similar experiments were performed for C12 with an aromatic moiety of Ni(porph) as the tail group (Fig. 1a), which is expected to form the usual

^a Department of Chemistry, Graduate School of Science and Engineering, Tokyo Institute of Technology, 2-12-1 W4-10 Ookayama, Meguro-ku, Tokyo 152-8551, Japan. E-mail: fujii.s.af@m.titech.ac.jp, tnishino@chem.titech.ac.jp

^b Department of Molecular and Macromolecular Chemistry, Graduate School of Engineering, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, Aichi 464-8603, Japan. E-mail: hshino@chembio.nagoya-u.ac.jp

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d3tc04166b>

same order of rapture force reported for the hydrogen bonding between two 2-ureido-4[1*H*]-pyrimidinone (UPy) moieties (*ca.* 180 pN).¹⁵ Force measurements of C12–Ni(porph) aromatic pairs suggest that the π – π stacking interactions of aromatics are of the same order as that of antiaromatics (Fig. 3b and Fig. S1, ESI†) and that the forces for aromatic and antiaromatic interactions are in the order of 150–220 pN. The result is consistent with our previous observation based on ¹H NMR analysis.¹⁰ This argument was supported by density functional theory calculations.¹⁰

Face-to-face stacking of antiaromatics leads to energetic stabilization, which is referred to as stacked-ring aromaticity.^{8,16} This energetic stabilization is due to the small HOMO–LUMO gap of the antiaromatics and the interaction between the frontier orbitals when the antiaromatics are stacked face-to-face.¹⁷ Meanwhile, energy decomposition analyses of the π – π stacking structures of norcorrole and porphyrins demonstrate that the dispersion interaction is more favourable for the antiaromatic system than the aromatic one. The analyses also show stronger repulsive interaction for the former than the latter, due to large exchange repulsion in the stacked norcorrole with short stacking distance. The overall interactions in the stacking structures of the antiaromatic and aromatic compounds are counterbalanced by attractive and repulsive interactions. Consequently, the interaction energies and rapture forces of π -stacked aromatics and π -stacked antiaromatics can be comparable in the force spectroscopy measurements. Theoretical calculations predict interaction energies of 1.34 eV and 1.58 eV for aromatic and antiaromatic pairs, respectively (Fig. S2 and ESI†). To accurately determine the interaction force at the single molecule scale, it is necessary to clarify the surface density of antiaromatic molecules adsorbed on the probe surface, the actual contact area between the AFM probe and the sample, and the dynamics of the breaking process of the interacting molecular pairs, which are far beyond the current standard experimental level and will be addressed in the future. This study supports the formation of π -stacks through an attractive interaction between two antiaromatic π -systems at the single-molecule scale to demonstrate the potential of π -stacks of antiaromatics as building blocks for the construction of organic frameworks in molecular electronics.

Experimental

Materials and sample preparation

Nickel(II) antiaromatic and aromatic compounds (C12–Ni(nor) and C12–Ni(porph), Fig. 1a) were prepared according to previously reported procedures.¹⁸ Au(111) substrates were prepared by the thermal vacuum deposition of Au on mica substrates. STM tips were mechanically cut from an Au wire (diameter 0.3 mm, purity >99.9%; Nilaco). Samples are prepared by dipping the Au(111) substrate into a 1 mM ethanol solution of hexanethiol (C6) for 1 hour. Then, the substrate was immersed in 0.5–1 mg ml^{−1} toluene solution of C12–Ni(nor) or 12–Ni(porph) (Fig. 1a) for 10 min. After immersion, the substrate was washed thoroughly with toluene and dried under a flow of inert gas. For control experiments, samples were prepared by dipping the Au(111)

substrate into a 1 mM ethanol solution of C6 for 1 h. After immersion, the substrate was washed thoroughly with ethanol and dried under a flow of inert gas.

STM and AFM measurements

STM measurements were performed using an MS-10 STM (Bruker), controlled by a NanoScope V (Bruker) under ambient conditions. All the STM images presented were recorded in constant current mode using mechanically cut Au tips.

Force spectroscopy measurements were performed using a multimode AFM (Bruker) controlled by a NanoScope V (Bruker) under ambient conditions. Au-coated AFM cantilevers (PPP-NCSTAu) were purchased from NANOSENSORS. The nominal spring constant of the cantilever and radius of curvature of the cantilever probe are 7.6 N m^{−1} and 50 nm, respectively. Prior to the force spectroscopy measurements, the surface of the AFM cantilever tip was modified with C6 and C12–Ni(nor) or C6 and C12–Ni(porph) in the same manner as the sample preparation.

Conclusions

The interaction between the π -systems of norcorrole Ni(II) was investigated at the single molecule scale using scanning probe microscopy. The interaction was measured as the adhesive force acting between norcorrole molecules anchored to the probe surface and the sample surface. This study reveals substantial interactions in antiaromatic π -systems and provides a better understanding of the interactions that make π -stacks of antiaromatics suitable for building blocks in organic electronics.

Author contributions

All the authors contributed to writing the manuscript.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This research was supported in part by JSPS KAKENHI (No. JP22H04974, JP23K04517, and JP21H01959) and JST SICORP (JPMJSC22C2).

Notes and references

- V. Coropceanu, J. Cornil, D. A. da Silva Filho, Y. Olivier, R. Silbey and J.-L. Brédas, *Chem. Rev.*, 2007, **107**, 926–952.
- X. Li, W. Ge, S. Guo, J. Bai and W. Hong, *Angew. Chem., Int. Ed.*, 2023, **62**, e202216819.
- R. Frisenda, V. A. E. C. Janssen, F. C. Grozema, H. S. J. van der Zant and N. Renaud, *Nat. Chem.*, 2016, **8**, 1099–1104.
- M. Carlotti, A. Kovalchuk, T. Wächter, X. Qiu, M. Zharnikov and R. C. Chiechi, *Nat. Commun.*, 2016, **7**, 13904.

- 5 X. Li, Q. Wu, J. Bai, S. Hou, W. Jiang, C. Tang, H. Song, X. Huang, J. Zheng, Y. Yang, J. Liu, Y. Hu, J. Shi, Z. Liu, C. J. Lambert, D. Zhang and W. Hong, *Angew. Chem., Int. Ed.*, 2020, **59**, 3280–3286.
- 6 Y. Tang, Y. Zhou, D. Zhou, Y. Chen, Z. Xiao, J. Shi, J. Liu and W. Hong, *J. Am. Chem. Soc.*, 2020, **142**, 19101–19109.
- 7 R. Nozawa, H. Tanaka, W.-Y. Cha, Y. Hong, I. Hisaki, S. Shimizu, J.-Y. Shin, T. Kowalczyk, S. Irle, D. Kim and H. Shinokubo, *Nat. Commun.*, 2016, **7**, 13620.
- 8 J. Aihara, *J. Phys. Chem. A*, 2009, **113**, 7945–7952.
- 9 S. Fujii, S. Marques-Gonzalez, J.-Y. Shin, H. Shinokubo, T. Masuda, T. Nishino, N. P. Arasu, H. Vázquez and M. Kiguchi, *Nat. Commun.*, 2017, **8**, 15984.
- 10 H. Kawashima, S. Ukai, R. Nozawa, N. Fukui, G. Fitzsimmons, T. Kowalczyk, H. Fliegl and H. Shinokubo, *J. Am. Chem. Soc.*, 2021, **143**, 10676–10685.
- 11 R. Nozawa, J. Kim, J. Oh, A. Lamping, Y. Wang, S. Shimizu, I. Hisaki, T. Kowalczyk, H. Fliegl, D. Kim and H. Shinokubo, *Nat. Commun.*, 2019, **10**, 3576.
- 12 J. M. Tour, L. Jones, D. L. Pearson, J. J. S. Lamba, T. P. Burgin, G. M. Whitesides, D. L. Allara, A. N. Parikh and S. Atre, *J. Am. Chem. Soc.*, 1995, **117**, 9529–9534.
- 13 O. H. Olubowale, S. Biswas, G. Azom, B. L. Prather, S. D. Owoso, K. C. Rinee, K. Marroquin, K. A. Gates, M. B. Chambers, A. Xu and J. C. Garno, *ACS Omega*, 2021, **6**, 25860–25875.
- 14 <https://www.nanosensors.com/pointprobe-plus-non-contact-soft-tapping-mode-au-coating-afm-tip-PPP-NCSTAu>.
- 15 S. Zou, H. Schönherr and G. J. Vancso, *Angew. Chem., Int. Ed.*, 2005, **44**, 956–959.
- 16 Y. Tsuji, K. Okazawa and K. Yoshizawa, *J. Org. Chem.*, 2023, **88**, 14887–14898.
- 17 K. Okazawa, Y. Tsuji and K. Yoshizawa, *J. Phys. Chem. A*, 2023, **127**, 4780–4786.
- 18 (a) R. Nozawa, K. Yamamoto, J.-Y. Shin, S. Hiroto and H. Shinokubo, *Angew. Chem., Int. Ed.*, 2015, **54**, 8454–8457; (b) G.-Y. Gao, A. J. Colvin, Y. Chen and X. P. Zhang, *J. Org. Chem.*, 2004, **69**, 8886–8892.