

# Thermally stable coordinatively unsaturated alkyl complexes resistant to $\beta$ -hydride elimination: $\text{Tp}^{\text{iPr}}\text{M}-\text{CH}_2\text{CH}_3$ ( $\text{M} = \text{Co}, \text{Fe}$ )

Nobuhiko Shirasawa, Munetaka Akita,\* Shiro Hikichi and Yoshihiko Moro-oka\*

Research Laboratory of Resources Utilization, Tokyo Institute of Technology, 4259 Nagatsuta, Midori-ku, Yokohama 226-8503, Japan. E-mail: makita@res.titech.ac.jp

Received (in Cambridge, UK) 3rd December 1998, Accepted 27th January 1999


**Coordinatively unsaturated hydrocarbyl complexes bearing  $\beta$ -hydrogen atoms,  $\text{Tp}^{\text{iPr}}\text{M}-\text{CH}_2\text{CH}_3$  [ $\text{M} = \text{Fe}, \text{Co}$ ;  $\text{Tp}^{\text{iPr}} = \text{hydrotris}(3,5\text{-diisopropylpyrazolyl})\text{borate}$ ], **14e** and **15e** species, respectively, are prepared; they are resistant to  $\beta$ -hydride elimination.**

'18-Electron (EAN) rule' and ' $\beta$ -hydride elimination' are usually key principles in predicting stability of organometallic compounds.<sup>1</sup> In a previous paper, we reported a complex which does not obey the 18e rule: the coordinatively unsaturated **14e**  $\eta^1$ -allyliron complex supported by the  $\text{Tp}^{\text{iPr}}$  ligand [ $\text{Tp}^{\text{iPr}} = \text{hydrotris}(3,5\text{-diisopropylpyrazolyl})\text{borate}$ ],  $\text{Tp}^{\text{iPr}}\text{Fe}-\text{CH}_2\text{CH}=\text{CH}_2$ .<sup>2</sup> Although  $\eta^3$  ( $\pi$ )-coordination of the allyl ligand would give a 16e species, closer to the coordinatively saturated 18e configuration, X-ray crystallography of the allyliron complex revealed that the  $\eta^1$  ( $\sigma$ )-coordination of the allyl ligand to the iron center led to the highly coordinatively unsaturated **14e** species. Our communication was followed by a recent report by Parkin *et al.*, who disclosed the synthesis and unusual reactivity of the related methyliron complex,  $[\text{PhTp}^{\text{Bu}}]\text{Fe}-\text{Me}$  [ $\text{PhTp}^{\text{Bu}} = \text{PhB}(3\text{-Bu}^t\text{-pz})_3$ ,  $\text{pz} = \text{pyrazolyl}$ ],<sup>3</sup> and thus the study on the coordinatively unsaturated hydrocarbyl complexes of late transition metals has attracted increasing attention.<sup>4</sup> During the course of our study, we have succeeded in the synthesis and characterization of coordinatively unsaturated alkyl complexes resistant to  $\beta$ -hydride elimination.


Reaction of the Fe and Co chloride complexes **1** with ethylmagnesium bromide in THF afforded pale yellow (**2a**) and blue products (**2b**), respectively, after removal of the inorganic salts by filtration through a Celite pad followed by crystallization from pentane [eqn. (1)]. X-Ray crystallography of the Co

complex **2a** was also assigned to  $\text{Tp}^{\text{iPr}}\text{Fe}-\text{CH}_2\text{CH}_3$  with tetrahedral structure, on the basis of its cell parameters<sup>†</sup> similar to those of **2b**, but could not be refined satisfactorily due to the low quality of the crystals. In contrast to the Co and Fe complexes, the reaction of the Ni complex **1c** afforded an intractable mixture of products from which no characterizable product could be isolated.

In order to further confirm the presence of a M–C bond in **2**, the ethyl complexes **2a,b** were subjected to carbonylation. Stirring a toluene solution of **2a,b** for 1 h under CO atmosphere (1 atm) produced **3a,b**, respectively. Their IR spectra containing



complex **2b** [Fig. 1(a)]<sup>†</sup> revealed the formation of the ethyl complex with tetrahedral coordination geometry as judged by the similar N–Co distances and N–Co–CH<sub>2</sub> angles [Co1–N11



**Fig. 1** Molecular structures of **2b** (a) and **3c** (b) drawn at the 30% probability level.

$\nu(\text{CO})$  and  $\nu(\text{C}=\text{O})$  vibrations suggest formation of acyl complexes.<sup>‡</sup> The diamagnetic iron complex **3a** was characterized on the basis of its  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra indicating the  $(\text{CO})_2\text{Fe}-\text{C}(\text{=O})\text{CH}_2\text{CH}_3$  functional group in addition to the  $\kappa^3\text{-Tp}^{\text{IPr}}$  ligand, and its IR spectrum suggests the presence of two rotamers as observed for  $\text{TpFe}(\text{CO})_2-\text{C}(\text{=O})\text{CH}_3$ .<sup>5</sup> The assignment of the monocarbonyl structure of the cobalt complex **3b** is based on the single  $\nu(\text{CO})$  absorption. Interestingly, although the ethylnickel complex **2c** could not be isolated from the Grignard reaction mentioned above, the reaction under CO atmosphere produced the acyl complex **3c**, which was characterized spectroscopically and structurally [Fig. 1(b)].<sup>†</sup> The Ni center adopts a trigonal bipyramidal structure with the  $\text{Ni1}-\text{Ni1}-\text{C1}$  axis, where the  $\kappa^3\text{-Tp}^{\text{IPr}}$  ligand occupies the two basal and one apical coordination sites and the acyl and CO ligands occupy the remaining apical and coordination sites, respectively, as judged by the interligand angles [ $\text{Ni1}-\text{Ni1}-\text{C1}$  171.7(1) ( $\approx 180^\circ$ );  $\text{N21}-\text{Ni1}-\text{C1}$  89.8(2),  $\text{N31}-\text{Ni1}-\text{C1}$  87.7(1),  $\text{C1}-\text{Ni1}-\text{C4}$  87.2(2),  $\text{N11}-\text{Ni1}-\text{N21}$  85.1(1),  $\text{N11}-\text{Ni1}-\text{N31}$  86.6(1),  $\text{N11}-\text{Ni1}-\text{C4}$  101.1(2) $^\circ$  ( $\approx 90^\circ$ );  $\text{N21}-\text{Ni1}-\text{N31}$  98.9(1),  $\text{N21}-\text{Ni1}-\text{C4}$  131.0(2),  $\text{N31}-\text{Ni1}-\text{C4}$  129.8(2) $^\circ$  ( $\approx 120^\circ$ )]. In accord with this view, the axial  $\text{N}-\text{Ni}$  distance is slightly longer than the equatorial ones [ $\text{Ni1}-\text{N11}$  2.058(3) Å >  $\text{Ni1}-\text{N21}$  2.018(3),  $\text{Ni1}-\text{N31}$  2.034(3) Å]. As for the  $\text{Ni}(\text{CO})-\text{C}(\text{=O})\text{Et}$  part, the structure of the acyl part is normal [ $\text{Ni1}-\text{C1}$  1.947(4),  $\text{O1}-\text{C1}$  1.192(5) Å;  $\text{Ni1}-\text{C1}-\text{O1}$  124.2(3),  $\text{Ni1}-\text{C1}-\text{C2}$  115.7(3),  $\text{C1}-\text{C2}-\text{C3}$  112.2(4) $^\circ$ ], and the CO ligand is coordinated to the Ni center in a typical  $\eta^1$ -fashion [ $\text{Ni1}-\text{C4}$  1.767(5),  $\text{O2}-\text{C4}$  1.126(6) Å;  $\text{Ni1}-\text{C4}-\text{O2}$  177.0(4) $^\circ$ ]. Thus the successful characterization of the acyl complexes **3** supports the presence of the  $\text{M}-\text{CH}_2\text{CH}_3$  functional group in the starting complexes **2**. The formation of the acyl complexes **3** via addition and insertion of CO molecules is in contrast to the result of carbonylation of the related iron-alkyl complex,  $[\text{PhTp}^{\text{Bu}}]\text{Fe}-\text{Me}$ , giving the  $\text{Fe}(\text{I})$  carbonyl complex,  $[\text{PhTp}^{\text{Bu}}]\text{Fe}-\text{CO}$ .<sup>3</sup> The difference may arise from the highly sterically demanding  $\text{PhTp}^{\text{Bu}}$  ligand, which would hinder formation of penta- or hexa-coordinated species leading to the acyl structure.

When the ethyl complexes **2a,b** were heated in heptane for 5 h at  $110^\circ\text{C}$ , only small amounts of ethane [10 (**2a**) and 8% (**2b**) yields] were detected together with trace amounts of ethene by GLC analysis of the gas phase. In contrast, hydrogenolysis (2 atm; 16 h at room temperature) produced ethane in 60 (**2a**) and 91% (**2b**) yields and protonolysis with aqueous  $\text{HCl}$  afforded ethane in 66 (**2a**) and 96% (**2b**) yields. Thus complexes **2** bearing  $\beta$ -hydrogen atoms were found to be resistant to ' $\beta$ -hydride elimination'.

The ethyl-iron and -cobalt complexes prepared in the present study are highly coordinatively unsaturated 14e ( $\text{Fe}$ : **2a**) and 15e species ( $\text{Co}$ : **2b**), respectively, which are quite sensitive to air and moisture. It is remarkable that, despite their coordinatively unsaturated electronic structures, they are resistant to  $\beta$ -hydride elimination and are stable at room temperature. The magnetic moments [ $\mu = 5.0 \mu_{\text{B}}$  (**2a**),  $4.2 \mu_{\text{B}}$  (**2b**)] close to the calculated spin only magnetic moments [4.90 ( $S = 2$ ), 3.87 ( $S = 3/2$ )] and EHMO calculations indicate that the ethyl complexes **2** are high spin species and all their non-bonding d-orbitals are fully or half occupied. The stability of the ethyl complexes **2** may be interpreted in terms of the lack of a vacant

coordination site (a vacant non-bonding d-orbital)<sup>1</sup> and the high spin electronic configurations should be a result of the tetrahedral coordination geometry regulated by the tripodal  $\text{Tp}^{\text{IPr}}$  ligand. Their electronic structure, however, should be flexible. Upon interaction with an appropriate substrate the electronic configuration may be changed so as to form a vacant coordination site and accommodate a substrate into the coordination sphere as typically exemplified by carbonylation giving the acyl complexes **3**.

We are grateful to the Ministry of Education, Science, Sports, and Culture of the Japanese Government (Grants-in-Aid for Scientific Research: No. 08102006 and 09238105) for financial support of this research.

## Notes and references

<sup>†</sup> X-Ray diffraction measurements were made on a Rigaku RAXIS IV imaging plate area detector with graphite-monochromated  $\text{Mo-K}\alpha$  radiation. The structures were solved by using the teXsan crystal structure analysis package. *Crystal data*: **2b**:  $\text{C}_{26}\text{H}_{51}\text{N}_6\text{BCo}$ ,  $M = 553.5$ ,  $T = -60^\circ\text{C}$ , monoclinic, space group  $P2_1/c$ ,  $a = 10.309(3)$ ,  $b = 16.21(2)$ ,  $c = 19.207(5)$  Å,  $\beta = 100.02(2)^\circ$ ,  $V = 3160(4)$  Å<sup>3</sup>,  $Z = 4$ ,  $D_{\text{c}} = 1.16 \text{ g cm}^{-3}$ ,  $\mu = 5.7 \text{ cm}^{-1}$ ,  $R_1 = 0.083$  for the 3267 unique data with  $F_o > 4\sigma(F_o)$  and 347 parameters,  $wR_2 = 0.224$  for all 4201 unique data.

**3c**:  $\text{C}_{31}\text{H}_{51}\text{BN}_6\text{NiO}_2$ ,  $M = 609.3$ ,  $T = -60^\circ\text{C}$ , monoclinic, space group  $P2_1/n$ ,  $a = 9.968(5)$ ,  $b = 15.820(2)$ ,  $c = 21.570(2)$  Å,  $\beta = 97.02(2)^\circ$ ,  $V = 3376(1)$  Å<sup>3</sup>,  $Z = 4$ ,  $D_{\text{c}} = 1.20 \text{ g cm}^{-3}$ ,  $\mu = 6.1 \text{ cm}^{-1}$ ,  $R_1 = 0.084$  for the 5974 unique data with  $F_o > 4\sigma(F_o)$  and 431 parameters,  $wR_2 = 0.238$  for all 6368 unique data. The disordered C25, C26, C28, C29, C38 and C39 parts were refined taking into account the minor components. CCDC 182/1153. See <http://www.rsc.org/suppdata/cc/1999/417/> for crystallographic files in .cif format.

Cell parameters for **2a**:  $T = -60^\circ\text{C}$ , monoclinic, space group  $P2_1/c$ ,  $a = 10.465(4)$ ,  $b = 16.18(1)$ ,  $c = 19.200(8)$  Å,  $\beta = 100.87(1)^\circ$ ,  $V = 3193(3)$  Å<sup>3</sup>.

<sup>‡</sup> *Selected spectral data*: **2a**: 2543  $\text{cm}^{-1}$  ( $\nu_{\text{BH}}$ ). **2b**: 2541  $\text{cm}^{-1}$  ( $\nu_{\text{BH}}$ ). **3a**: 2553 ( $\nu_{\text{BH}}$ ), 2024, 2000, 1957, 1930 ( $\nu_{\text{CO}}$ ), 1668, 1620  $\text{cm}^{-1}$  ( $\nu_{\text{C=O}}$ ).  $\delta_{\text{H}}(\text{CD}_2\text{Cl}_2)$  6.39 (2H, 2, pz), 6.36 (1H, s, pz), 3.67 (q,  $J$  7.0 Hz,  $\text{CH}_2$ ).  $\delta_{\text{C}}(\text{CD}_2\text{Cl}_2)$  259.5 [s,  $\text{Fe}-\text{C}(\text{=O})$ ], 215.0 (s,  $\text{Fe}-\text{CO}$ ), 56.8 (t,  $J$  130 Hz,  $\text{CH}_2$ ), 10.3 (q,  $J$  128 Hz,  $\text{CH}_2\text{CH}_3$ ). **3b**: 2540 ( $\nu_{\text{BH}}$ ), 1999 ( $\nu_{\text{CO}}$ ), 1636  $\text{cm}^{-1}$  ( $\nu_{\text{C=O}}$ ). **3c**: 2547 ( $\nu_{\text{BH}}$ ), 2009 ( $\nu_{\text{CO}}$ ), 1682  $\text{cm}^{-1}$  ( $\nu_{\text{C=O}}$ ).  $\delta_{\text{H}}(\text{CD}_2\text{Cl}_2)$  5.84 (3H, s, pz), 1.88 (q,  $J$  7.4 Hz,  $\text{CH}_2$ ), 0.45 (t,  $J$  7.4 Hz,  $\text{CH}_2\text{CH}_3$ ).  $\delta_{\text{C}}(\text{CD}_2\text{Cl}_2)$  225.9 (s,  $\text{Ni}-\text{C}(\text{=O})$ ), 187.4 (s,  $\text{Ni}-\text{CO}$ ), 40.7 (t,  $J$  132 Hz,  $\text{CH}_2$ ), 11.1 (q,  $J$  127 Hz,  $\text{CH}_2\text{CH}_3$ ).

- 1 A. Yamamoto, *Organotransition Metal Chemistry*, Wiley-Interscience, New York, 1986; J. P. Collman, L. S. Hegeudus, J. R. Norton and R. G. Finke, *Principles and Applications of Organotransition Metal Chemistry*, University Science Books, Mill Valley, CA, 1987; R. H. Crabtree, *The Organometallic Chemistry of the Transition Metals*, Wiley-Interscience, New York, 2nd edn., 1994.
- 2 M. Akita, N. Shirasawa, S. Hikichi and Y. Moro-oka, *Chem. Commun.*, 1998, 973.
- 3 J. L. Kisko, T. Hascall and G. Parkin, *J. Am. Chem. Soc.*, 1998, **120**, 10561.
- 4 Several examples of coordinatively unsaturated hydrocarbyl complexes kinetically stabilized by bulky ligands are known. See, for example: B. K. Bower and H. G. Tennent, *J. Am. Chem. Soc.*, 1972, **94**, 2512; A. Kose, E. Solari, C. Floriani, A. Chiesi-Villa, C. Rizzoli and N. Re, *J. Am. Chem. Soc.*, 1994, **116**, 9123; R. C. Kerber, *Comprehensive Organometallic Chemistry II*, ed. E. W. Abel, F. G. A. Stone and G. Wilkinson, Pergamon, Oxford, 1995, vol. 7, ch. 2.3.1.
- 5 F. A. Cotton, B. A. Frenz and A. Shaver, *Inorg. Chim. Acta*, 1973, **7**, 161.

Communication 8/09441A