


Food & Function

Linking the chemistry and physics of food with health and nutrition

rsc.li/food-function

Volume 16
Number 8
21 April 2025
Pages 2917-3230



ISSN 2042-650X

PAPER

Víctor López et al.

Apple peels as an edible source of phenolic bioactive compounds with antidiabetic and antiglycation properties

Cite this: *Food Funct.*, 2025, 16, 2947

Apple peels as an edible source of phenolic bioactive compounds with antidiabetic and antiglycation properties

Javier Cano-Lou, ^{a,b} Adrián Millán-Laleona, ^{a,b} Rares Candrea,^a Francisco Les, ^{a,b} Ana Pina,^c Giovanni Caprioli ^d and Víctor López ^{*a,b}

Apples (*Malus domestica* Borkh.) are one of the most consumed fruits around the world with a high production of peels as wastes and by-products. In this work, peels from different commercial and local apple samples are explored as a source of phenolic bioactive compounds that could be directly related to the prevention of type 2 diabetes. Six different cultivars from local and commercial apple samples were processed to obtain the phenolic compounds by ultrasonication of the peels using methanol as the solvent. The phenolic content was explored using the Folin–Ciocalteu assay and the quantification of 37 individual phenolic compounds was carried out by high-performance liquid chromatography coupled with mass spectrometry (HPLC-MS/MS). Cellular viability was determined by performing the MTT assay in Caco-2 cell cultures exposed to the phenolic extracts. Subsequently, the capacity to inhibit α -glucosidase, α -amylase and pancreatic lipase enzymes, as well as antiglycation and antioxidant activities, was evaluated. These apple peel samples were considered a source of phenolic compounds with hyperoside, delphinidin 3,5-diglucoside, chlorogenic acid, phlorizin, epicatechin and procyanidin B2 as the main constituents. All samples neutralized the production of advanced glycation end-products and exhibited anti-radical activities in a dose-dependent manner; four samples (Amarilla de Octubre, Manzana Helada, Verde Doncella and Pinova) inhibited α -glucosidase but only the sample known as “Amarilla de Octubre” was successful in inhibiting pancreatic α -amylase. Cytotoxicity was discarded in Caco-2 cell cultures at physiological concentrations considering these extracts as a source of phenolic compounds with antidiabetic, antiglycation and antioxidant properties.

Received 26th October 2024,
Accepted 16th February 2025

DOI: 10.1039/d4fo05241b

rsc.li/food-function

1. Introduction

Diabetes mellitus is a chronic disease that develops when the pancreatic islet β -cells cannot produce insulin, or the body becomes resistant to insulin with a deficient action in sensitive tissues.¹ This contributes to high blood glucose levels, which induce a rise in reactive oxygen species (ROS) and inflammation, culminating in long-term damage to hyperglycaemia-sensitive cells in various organs and tissues.² The World Health Organization (WHO) classifies hyperglycaemic disorders into diabetes mellitus type 1, type 2 and gestational diabetes, with type-2 diabetes accounting for 90% of all diabetes

cases, which occurs due to an excess body weight, sedentary behaviour and dietary changes.³ The International Diabetes Federation (IDF) estimates 783.2 million diabetic people by 2045, which will be directly related to inevitable factors, such as population growth, major urbanisation, and an elderly society, combined with a better diabetes treatment that increases prevalence through decreased mortality.⁴ Some strategies on modifiable factors such as lifestyle behaviour and a healthy diet are required to stall the rise of developing diabetes and its complication.⁵ Plant-based diets or herbal supplements have traditionally had beneficial effects on chronic disease and have been associated with a decrease in the chance of developing type 2 diabetes.^{6,7} One of the most common components in plant-derived foods is polyphenols,⁸ which are phenolic compounds with a survival and antioxidant function as secondary metabolites in plants.⁹ These phenolic compounds might help in the prevention of metabolic and non-communicable diseases such as type 2 diabetes in part due to their antiradical and anti-inflammatory properties associated with specific antidiabetic properties such as enzyme inhibitory properties, anti-glycation activity, improve-

^aDepartment of Pharmacy, Faculty of Health Sciences, Universidad San Jorge, 50830 Zaragoza, Spain. E-mail: ilopez@usj.es

^bInstitute of Agri-food Research of Aragón, IA2, Zaragoza University-CITA, 50013 Zaragoza, Spain

^cUnidad de Hortofruticultura, Centro de Investigación y Tecnología Agroalimentaria de Aragón (CITA), Av Montañana 930, 50059 Zaragoza, Spain

^dChemistry Interdisciplinary Project (ChIP), School of Pharmacy, University of Camerino, Via Madonna delle Carceri, 62032 Camerino, Italy



ment of beta cell function and insulin resistance, as well as regulation of carbohydrate metabolism, among others.¹⁰ The most frequent classification of phenolic compounds includes five main classes: phenolic acids, flavonoids, tannins, stilbenes, lignans, and others.⁹ The flavonoid group is frequently found in plant foods, and some of the subgroups are flavonols, flavanols, flavones, isoflavones, anthocyanins and flavanones.¹¹ These phenolic compounds could be found in high amounts in plant residues, including food by-products and wastes.¹² Compared to other food-processing sectors, the agri-food industries in relation to fruits and vegetables generate increasing amounts of waste including 25–30% of peels, which are usually discarded.¹³ According to the Food and Agriculture Organization of the United Nations (FAO), 1.3 billion tons per year of food get lost or wasted in the world.¹⁴ In addition, the agri-food industries and distributors in Spain produced 0.0022 kg L⁻¹ of by-products and 0.0004 kg L⁻¹ of wastes for each kg L⁻¹ of end products.¹⁵ Phenolic compounds from this huge volume of plant residues could be extracted and transformed into health-promoting ingredients for the development of innovative food products such as nutraceuticals, food supplements, cosmetics or medicinal products¹⁶ and could provide a source of a great variety of phytochemicals as hypoglycaemic drugs to prevent diabetes type 2.^{17,18} Certain peels from fruits and vegetables are edible, but in many cases, they are also frequent residues found in the food industry. In this context, it is remarkable to delve into the added value from residues of apple-derived products which are usually processed as peeled fruits, producing a large quantity of peel as industrial waste.¹⁹ Furthermore, peels are considered as a great source of nutrients and phytochemicals with higher bioactivity than the pulp.^{20,21} Nevertheless, the quantitative distribution of phenolic compounds is variable depending not only on the plant type or the plant organ, but also on the genetics, maturity and climatic conditions.²² The main goal of this manuscript is to investigate whether edible peels from autochthonous and commercial apple cultivars can also be used as a natural matrix of phenolic bioactive compounds for different industrial applications.

2. Materials and methods

2.1 Apple peel samples

Six different varieties of fresh apples (Pinova and Verde Doncella as commercial samples; Esperiega de Ademuz, Manzana Helada, Amarilla de Octubre and Borau 01 as local autochthonous cultivars) were collected in 2022 from Garcipollera (Huesca, Aragon) between the months of August–October. After the apples were peeled, the pulp and the peel were completely separated in different parts. The peel tissue was frozen at –80 °C and lyophilized.

2.2 Extraction of phenolic compounds and determination of total phenolic content (TPC)

For this study, the extraction of phenolic compounds was achieved using methanol and the solid liquid ratio (SLR) was a

1 : 33 mixture. The method used for the extraction was treatment with intense ultrasonic waves for 20 minutes at room temperature, also known as ultrasound-assisted extraction (UAE). UAE was used as an effective extraction method of bioactive compounds, such as phenolic compounds.¹³ The liquid extract was centrifuged at 4000g for 10 minutes, the supernatant was filtered by clarification filtration with filter paper, and the solvent was evaporated using a rotary evaporator. The end product was a semi-solid mixture of each apple peel extract. TPC was determined using the Folin–Ciocalteu assay. A volume of 100 µL of the Folin–Ciocalteu reagent 10% (Sigma-Aldrich, Barcelona, Spain) was added to each well along with 20 µL of diluted extracts in methanol. After being kept for 3 minutes in the dark at ambient temperature, the reaction solution was mixed with 80 µL of Na₂CO₃ (10%).

The reaction was concluded after 30 minutes of incubation under the previous conditions, and the absorbance (760 nm) was measured on a Synergy H1 reader (BioTek® Instruments, Inc., Winooski, VT, USA). A standard curve of gallic acid was used to quantify the TPC, which was expressed as milligrams of gallic acid equivalent (GAE) per gram of the extract.

2.3 Analysis of phenolic compounds by HPLC-MS/MS

2.3.1 Reagents and standards. A total of 37 analytical standards have been used. Some of the standards, such as cyanidin-3-glucoside chloride, delphinidin-3,5-diglucoside chloride, delphinidin-3-galactoside chloride, petunidin-3-glucoside chloride, malvidin-3-galactoside chloride, quercetin-3-glucoside and kaempferol-3-glucoside were acquired from PhytoLab (Vestenbergsgreuth, Germany), while the rest of the standards were provided by Sigma-Aldrich (Milan, Italy). The isolated standards were dissolved in HPLC-grade methanol, and a specific solution of each standard was obtained at 1000 mg L⁻¹. All solvents and solutions were filtered through a 0.2 µm polyamide filter supplied by Sartorius Stedim (Goettingen, Germany). In addition, all samples were filtered prior to the HPLC 126 analysis, with a Phenex™ RC 4 mm 0.2 µm syringeless filter from Phenomenex (Castel Maggiore, BO, Italy).

2.3.2 HPLC-ESI-MS/MS instrument and analysis. The analysis of polyphenols was carried out using high performance liquid chromatography coupled with tandem mass spectrometry (HPLC-MS/MS) using a triple quadrupole detector following a method developed in our laboratory.²³

Briefly, the analytical separation of the analytes was carried out on a Synergi Polar-RP C18 column preceded by a pre-column. The solvents used as mobile phases were water (A) and methanol each with 0.1% of formic acid at a flow rate of 0.8 mL min⁻¹ in the gradient elution mode. The volume used for injection was 2 µL, and the temperature of the oven was 30 °C.

The source of ionization used was an electrospray ionization (ESI) operating in negative and positive modes and the detector was a triple quadrupole working in the dynamic MRM (multiple reaction monitoring) mode.



2.4 Cell viability assays

Caco-2 cells (epithelial cells isolated from colon tissue) were acquired from ATCC. The cell culture was treated with DMEM medium (Sigma Aldrich), 10% foetal bovine serum (FBS) (Sigma Aldrich) and 1% penicillin–streptomycin (Sigma Aldrich) in an incubator under a 5% CO₂ atmosphere and at 37 °C. For the MTT assay, firstly, Caco-2 cells were seeded in 96-well plates at 2×10^4 cells per well for 24 hours. After that period, certain cells were treated with different apple peel extracts from 62.5 to 1000 µg extract per mL in 1% FBS and DMEM, meanwhile other cells (control) were cultivated with 1% FBS and DMEM without treatment for 24 hours. Following that, 100 µL of MTT stock solution (0.4 mg mL⁻¹ in 10% FBS) was added in each well and the plate was collocated in an incubator under a 5% CO₂ atmosphere and at 37 °C, in the dark for 3 hours. Once the formazan crystals were formed, the MTT solution was removed and 100 µL of DMSO was added. Finally, the dissolution of the crystals was carried out by agitation, and a purple colour appeared. The absorbance was read at 550 nm in a Synergy H1 Hybrid Multi-Mode Reader (Biotek, Bad Friedrichshall, Germany). The viability was calculated using the following formula (eqn (1)):

$$\text{Cell viability (\%)} = [\text{Abs}_{\text{sample}}/\text{Abs}_{\text{control}}] \times 100 \quad (1)$$

2.5 Enzyme inhibitory assays of α -glucosidase, α -amylase and lipase

2.5.1 α -Glucosidase (from *Saccharomyces cerevisiae*) inhibition. The inhibition of α -glucosidase (Sigma-Aldrich, Barcelona, Spain) was analysed using a 96-microplate reader using α -glucosidase from *Saccharomyces cerevisiae*. A volume of 100 µL of α -glucosidase solution (1.0 U mL⁻¹) was preincubated with 50 µL of the diluted extracts in each well at ambient temperature for 10 min. After that period, the reaction occurred upon adding 50 µL of 3 mM pNPG (4-nitrophenyl α -D-glucopyranoside) (Sigma-Aldrich, Barcelona, Spain) to the mixture solution, and the absorbance was measured at a wavelength of 405 nm after 5 min of incubation at 37 °C. Phosphate buffer was used to dissolve the reagents and the extracts. The inhibitory activity was calculated using the equation below (2):

$$\text{Inhibition (\%)} = [\text{Abs}_{\text{control}} - \text{Abs}_{\text{sample}}/\text{Abs}_{\text{control}}] \times 100 \quad (2)$$

2.5.2 α -Amylase (from porcine pancreas) inhibition. The inhibition reaction of α -amylase (Sigma-Aldrich) was carried out in 1.5 mL microtubes. 100 µL of the extract sample and 100 µL of porcine pancreatic α -amylase (2 mg mL⁻¹ centrifuged) were preincubated in microtubes at 37 °C for 5 minutes. After that, 1% starch solution (Sigma-Aldrich) was added and incubated at 37 °C for another 10 minutes. Following that, 200 µL of 1% 3,5-dinitrosalicylic acid (DNS) (Sigma-Aldrich) color reagent and 50 µL of 1 M NaOH were incubated for 5 minutes at 100 °C. Once the reaction was finished, the solution changed the colour and the microtubes were cooled in a bath of cold water. Phosphate buffer was used to dissolve the reagents and the extracts. To perform the absor-

bance measurement at 540 nm, 200 µL of the solution from each microtube was transferred to a 96-well microplate. The inhibitory activity was determined by applying eqn (2).

2.5.3 Lipase (from porcine pancreas) inhibition. The inhibition of pancreatic lipase (Sigma-Aldrich, Barcelona, Spain) was assessed using a 96-microplate reader. A solution in each well was prepared from the mixture of 40 µL of diluted extracts and 40 µL of lipase type II from porcine pancreas (2.5 mg mL⁻¹ centrifuged), and a preincubation period of 15 min at ambient temperature was needed. After that period, 20 µL of 10 mM *p*-4-nitrophenyl butyrate (NPB) (Sigma Aldrich, Barcelona) was added to the mixture solution, and the absorbance was measured at a wavelength of 405 nm after 15 min of incubation at 37 °C. Phosphate buffer was used to dissolve the reagents and the extracts. The inhibitory activity was determined using eqn (2).

2.6 Antiglycation and antioxidant activities

2.6.1. Inhibition of advanced glycation end products (AGEs). Formation of AGEs was evaluated using a 96-well black microplate. 50 µL of sample extracts, 100 µL of bovine serum albumin solution (Santa Cruz, Barcelona, Spain) (10 mg mL⁻¹), and 100 µL of fructose (Sigma-Aldrich, Barcelona, Spain) (0.5 M) were added to each well. All solutions were prepared in phosphate buffer. After incubation for 24 hours at 37 °C in the dark, the fluorescence was analysed at an excitation wavelength of 355 nm and an emission wavelength of 460 nm. The inhibitory activity was determined using eqn (2).

2.6.2. Radical scavenging activity against 2,2-diphenyl-1-picrylhydrazyl (DPPH). The DPPH (Sigma-Aldrich, Barcelona, Spain) inhibition activity of the extracts was evaluated through spectrophotometric techniques. A total of 150 µL of diluted extracts in methanol were mixed with 150 µL of a DPPH-methanol solution (0.04 mg mL⁻¹) and incubated in the dark at ambient temperature for 30 min (absorbance read at 517 nm). The percentage of inhibition was calculated using eqn (2).

2.7 Statistical analysis

Each assay was carried out at least three times on different days and all results were expressed as the mean \pm standard error (SEM) of different assays. GraphPad Prism v.8.0 (GraphPad Software, San Diego, CA, USA) was applied to execute data analyses such as nonlinear regressions, a one-way analysis of variance (ANOVA) with Dunnett's *post hoc* test, Student's *t*-test, and Pearson correlations and statistics with a 95% confidence level and 5% of wise significance. The determination of linear correlation was performed for the total polyphenol content using the Folin–Ciocalteu assay and antidiabetic, anti-glycation or antioxidant properties.

2.8 Chemical structure generation

The chemical structures of most of the phenolic compounds found in the extract samples were generated using ChemDraw v. 21.0.0 software (PerkinElmer Informatics, Waltham, Massachusetts, USA).



3. Results

3.1 Phenolic content using the Folin–Ciocalteu assay and HPLC-MS/MS

According to the TPC obtained from the Folin–Ciocalteu assay, the best apple peel sample is Amarilla de Octubre (142.80 ± 26.61 mg GAE per g extract) (Table 1), revealing significant differences against the rest of the samples, either autochthonous or commercial samples. Nevertheless, the commercial sample known as Pinova (52.81 ± 13.70 mg GAE per g extract), and the autochthonous sample of Borau 01 (41.79 ± 5.82 mg GAE per g extract) showed the lowest polyphenol content, compared to the rest of the samples.

Considering HPLC-MS/MS analysis, 37 individual phenolic compounds of interest in six apple peel extracts from different apple cultivars were monitored (Table 2). The phenolic groups detected in the samples belong to flavonols, flavan-3-ols, anthocyanins, dihydrochalcones and cinnamic acid compounds. Nevertheless, none of the hydroxybenzoic acid, stilbene and flavanone compounds were identified. Amarilla de Octubre, as determined by HPLC-MS/MS, is the best sample as it contains 3 times higher phenolics ($24\,840.98 \mu\text{g g}^{-1}$) than Borau 01 ($8576.39 \mu\text{g g}^{-1}$). The highest number of different phenolic compounds analysed (16 compounds) were detected in Amarilla de Octubre, while 14 compounds were revealed in Borau 01, and 15 were detected in the other samples.

The most abundant phenolic compounds contained in the samples were hyperoside, delphinidin 3,5-diglucoside, epicatechin, procyanidin B2, phlorizin and chlorogenic acid (Fig. 1). However, hyperoside followed by delphinidin 3,5-diglucoside were found in high quantities in all the samples. Hyperoside belongs to the flavonol compounds as a glycosylated variety of quercetin, and its levels varied from $6336.00 \mu\text{g g}^{-1}$ (Pinova) to $1209.07 \mu\text{g g}^{-1}$ (Borau 01). However, delphinidin 3,5-diglucoside is included in the anthocyanin compounds as a glycosylated form of delphinidin, and its levels varied between $5456.33 \mu\text{g g}^{-1}$ (Pinova) and $948.11 \mu\text{g g}^{-1}$ (Borau 01).

It is noteworthy that the commercial samples are a better source of flavonols and anthocyanins but have a shortage of flavan-3-ols, dihydrochalcones and cinnamic acids (Fig. 2). The phenolic fraction of commercial samples of Pinova and Verde Doncella was made up of 69.58% and 64.94% flavonols, along with 28.19% and 23.73% anthocyanins, respectively,

with hyperoside, delphinidin 3,5-diglucoside and isoquercitrin being the most abundant compounds in both samples. The main differences between them were the absence of flavan-3-ol compounds in Pinova, compared to the presence of 7.77% flavan-3-ols in the Verde Doncella sample. Procyanidin B2 ($471.07 \mu\text{g g}^{-1}$), epicatechin ($296.31 \mu\text{g g}^{-1}$) and catechin ($87.73 \mu\text{g g}^{-1}$) were contained in Verde Doncella.

Nevertheless, for the autochthonous samples, a lower proportion of flavonols and anthocyanins was obtained in their TPC, with respect to the commercial samples. The decreased ratio of flavonols and anthocyanins is due to a notable increase in other types of phenolic compounds such as flavon-3-ols, dihydrochalcones and cinnamic acids (Fig. 2).

The sample of Amarilla de Octubre showed the most abundant quantities of flavonols and anthocyanins among all the autochthonous samples. Although, in comparison with the commercial samples, a lower percentage of 34.16% and 14.92% was obtained in their TPC, respectively. This is attributed to the great amounts of flavan-3-ol compounds existing as epicatechin ($5244.89 \mu\text{g g}^{-1}$) or its dimerized form called procyanidin B2 ($4297.35 \mu\text{g g}^{-1}$).

Similarly, it is worth noting that phlorizin, the glycosylated form of phloretin, which belongs to dihydrochalcone compounds, appears as an important phenolic compound in the autochthonous extracts of Esperiega de Ademuz ($3092.59 \mu\text{g g}^{-1}$) and Manzana Helada ($1894.48 \mu\text{g g}^{-1}$). Apart from that, the main difference between them is the fluctuation of the amount of cinnamic acids, flavonols and anthocyanins. Esperiega de Ademuz showed higher quantities of cinnamic acids as chlorogenic acid ($910.65 \mu\text{g g}^{-1}$). Nonetheless, Manzana Helada was made up of 44.88% flavonols, along with 15.60% anthocyanins, being the second autochthonous sample beyond Amarilla de Octubre with a rich amount of these types of phenolic compounds.

In addition, despite the low TPC of the autochthonous sample of Borau 01, chlorogenic acid from cinnamic acids, function as a significant compound ($2312.85 \mu\text{g g}^{-1}$). Therefore, being the sample with the largest amount of this type of phenolic compound, the sample of Borau 01 is considered as a good source of that substance.

3.2 Cell viability in Caco-2 cultures

The cytotoxicity of the apple peel phenolic extracts was evaluated in a wide range of concentrations, from lower and more physiological to very high and non-physiological concentrations. The MTT assay showed that Amarilla de Octubre, Manzana Helada and Esperiega de Ademuz, induced a slight decrease in the cell viability at high concentrations compared to the commercial samples (Verde Doncella and Pinova) (Fig. 3). The three autochthonous samples with the lowest viability, and the commercial sample of Pinova at $1000 \mu\text{g mL}^{-1}$ showed significant differences compared to the control cells without treatment, in terms of viability. Nonetheless, the cell viability was superior by 50% at the highest analysed concentration of all samples; considering that $1000 \mu\text{g mL}^{-1}$ is a very

Table 1 Total phenolic content (TPC) corresponding to the mean \pm SEM of a minimum of three independent experiments measured using the Folin–Ciocalteu assay (* $p < 0.05$, ** $p < 0.001$ versus Amarilla de Octubre)

Samples	Folin–Ciocalteu (mg GAE per g extract)
Amarilla de Octubre	142.80 ± 26.61
Esperiega de Ademuz	$69.21 \pm 8.98^*$
Verde Doncella	$67.43 \pm 11.89^*$
Manzana Helada	$61.94 \pm 11.18^*$
Pinova	$52.81 \pm 13.70^*$
Borau 01	$41.79 \pm 5.82^{**}$



Table 2 Individual phenolic compounds analysed by HPLC-MS/MS

Compounds	Amarilla de Octubre	Pinova	Esperiega de Ademuz	Manzana Helada	Verde Doncella	Borau 01
Hydroxybenzoic acids						
Gallic acid	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
3-Hydroxybenzoic acid	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
4-Hydroxybenzoic acid	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Cinnamic acids						
Chlorogenic acid	2105.95	154.30	910.65	333.76	20.77	2312.85
Neochlorogenic acid	39.05	n.d.	18.22	16.09	n.d.	n.d.
Caffeic acid	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Vanillic acid	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Syringic acid	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
<i>P</i> -Coumaric acid	n.d.	15.05	n.d.	n.d.	n.d.	10.94
Ferulic acid	n.d.	73.39	n.d.	n.d.	n.d.	n.d.
Ellagic acid	104.85	55.28	100.34	103.25	308.72	n.d.
<i>trans</i> -Cinnamic acid	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Stilbenes						
Resveratrol	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Dihydrochalcones						
Phloridzin	341.08	157.32	3092.59	1894.48	62.29	408.87
Phloretin	0.18	0.89	0.82	0.46	n.d.	0.14
Flavonols						
Quercetin	12.45	1387.82	6.60	5.53	10.34	33.07
Kaempferol	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Myricetin	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Isorhamnetin	n.d.	12.30	n.d.	n.d.	n.d.	n.d.
Kaempferol-3-glucoside	24.54	10.10	7.50	5.37	9.85	6.71
Quercitrin	925.76	1447.40	517.20	952.75	793.34	442.78
Rutin	302.41	541.97	74.68	148.06	666.50	34.83
Hyperoside	4246.61	6336.00	2766.28	2677.83	3495.94	1209.07
Isoquercitrin	2973.13	4517.45	1669.32	1943.33	2168.95	770.70
Flavan-3-ols						
Catechin	515.34	n.d.	181.13	113.11	87.73	108.17
Procyanidin A2	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Procyanidin B2	4297.35	n.d.	1160.39	1331.09	471.07	1086.90
Epicatechin	5244.89	n.d.	1977.79	1255.01	296.31	1186.63
Flavanones						
Naringenin	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Hesperidin	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Anthocyanin						
Delphinidin-3,5-diglucoside	3707.40	5456.33	1979.19	1992.60	2611.10	948.11
Delphinidin-3-galactoside	15.80	n.d.	n.d.	n.d.	n.d.	16.60
Cyanidin-3-glucoside	n.d.	303.24	n.d.	n.d.	n.d.	n.d.
Petunidin-3-glucoside	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Pelargonidin-3-glucoside	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Pelargonidin-3-rutinoside	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Malvidin-3-galactoside	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Total polyphenolic content ($\mu\text{g g}^{-1}$)	24 840.98	20 484.65	14 462.70	12 772.71	11 002.2	8576.39

n.d.: not detected.

high and non-physiologically relevant concentration, apple peel extracts can be considered as non-cytotoxic.

3.3 Evaluation of the antidiabetic activity against α -glucosidase, α -amylase and lipase

The substance used as reference in the α -glucosidase assay was the therapeutic agent of acarbose, with a half-maximal inhibitory concentration (IC_{50}) of $220 \pm 64.10 \mu\text{g mL}^{-1}$. Only four extracts (Amarilla de Octubre, Manzana Helada, Verde

Doncella and Pinova) were able to inhibit α -glucosidase as indicated in Fig. 4. Amarilla de Octubre showed the lowest IC_{50} value related to the highest activity. The most powerful sample is followed by Manzana Helada and the commercial sample of Verde Doncella, which had interesting results as well. The commercial sample of Pinova was less effective than the previous mentioned samples. Nevertheless, no activity was found in the inhibition of α -glucosidase in the autochthonous samples of Borau 01 and Esperiega de Ademuz.






Fig. 1 Chemical structures of the most abundant phenolic compounds of the apple peel extracts.

The substance used as reference in the α -amylase assay was gallic acid, a phenolic compound with an IC_{50} value of $909 \pm 15 \mu\text{g mL}^{-1}$. The autochthonous sample of Amarilla de Octubre was the unique sample with enough capacity to inhibit the α -amylase enzyme, showing promising antidiabetic activity.

Regarding the lipase inhibition assay, the substance used as reference in the lipase assay was the therapeutic agent of orlistat, with an IC_{50} value of $27.68 \pm 1.53 \mu\text{g mL}^{-1}$. None of the extracts were able to show activities, as they did not reach a concentration of $2000 \mu\text{g mL}^{-1}$.

In the set of the antidiabetic assays, none of the samples were able to achieve the reference pure substances (Table 3). However, the IC_{50} value of autochthonous samples of Amarilla de Octubre and Manzana Helada were very similar to the IC_{50} of reference substances used in the α -glucosidase assays, with no significant difference between them. Commercial samples of Verde Doncella showed an interesting result as well in the α -glucosidase inhibition assay, with no significant difference compared to the acarbose.

3.4 Evaluation of the inhibition of AGEs formation, and the antioxidant activity against radical DPPH

The substance used as reference in the AGEs inhibition assay was the investigational drug of aminoguanidine, with an IC_{50} value of $55.01 \pm 5.63 \mu\text{g mL}^{-1}$. All apple peel samples were able to inhibit AGEs formation. It should be emphasized that the high activity of the autochthonous sample of Amarilla de

Octubre was indicated by the lowest IC_{50} value, followed by Manzana Helada and the commercial sample of Verde Doncella. Nevertheless, some of the autochthonous samples (Borau 01, Esperiega de Ademuz) and the commercial sample of Pinova showed the lowest activity, with the commercial sample of Pinova being the least effective.

The substance used as reference in the DPPH inhibition assay was gallic acid from phenolic compounds. It revealed an outstanding profile as radical scavengers, showing an IC_{50} value of $0.471 \pm 0.0791 \mu\text{g mL}^{-1}$. All apple peel samples were able to scavenge DPPH radicals. It should be noted that the high activity of the autochthonous sample of Amarilla de Octubre was indicated by the lowest IC_{50} value. Other autochthonous and commercial samples such as Esperiega de Ademuz, Manzana Helada and Verde Doncella had optimal and similar results in terms of inhibitory activity. Nevertheless, the autochthonous and commercial samples of Borau 01 and Pinova showed the worst results.

None of the apple peel extracts (Table 3) were able to achieve IC_{50} values of the reference pure substances. However, the IC_{50} value of autochthonous sample of Amarilla was very comparable to the IC_{50} value of the reference substance used in AGE and DPPH assays, with no significant difference between them. The IC_{50} value of the other autochthonous sample of Manzana Helada was very similar to the IC_{50} value of aminoguanidine used in the AGE inhibition assay, with no significant difference as well.






Fig. 2 Phenolic fraction expressed as the percentage (%) of (A) commercial samples (Pinova, Verde Doncella) and (B) autochthonous samples (Amarilla de Octubre, Esperiega de Ademuz, Borau 01 and Manzana Helada).

3.5 Determination of linear correlation of TPC using the Folin–Ciocalteu assay and antidiabetic, antiglycative or antioxidant properties

Linear correlation analysis of TPC was obtained using the Folin–Ciocalteu assay and the IC_{50} values of the antidiabetic,

antiglycative and antioxidant assays using the Pearson correlation coefficient (Fig. 5). No correlation was found between the α -glucosidase and AGE assays and TPC, which suggests that TPC is not related to antidiabetic activity. However, regarding the antioxidant activity, a clear correlation was found between the DPPH assay and TPC.






Fig. 3 Cell viability percentage expressed as the mean \pm SEM of a minimum of three independent experiments at different concentrations in microgram per millilitre (from 62.5 to 1000 μg extract per ml) of the apple peel extracts in CaCo-2 cell cultures.




Fig. 4 Antidiabetic activity against (A) α -glucosidase (*Saccharomyces cerevisiae*) and (B) α -amylase (porcine pancreas).

4. Discussion

Apple peels are edible fruit parts but also a residue generated by the fruit and food industry all over the world. Considering that according to the WHO, 422 million people worldwide have diabetes, we here hypothesized whether these fruit peels could be an excellent source of phenolic compounds with bioactive properties such as antidiabetic, antiglycation or antioxidant properties and different food or health applications. The total extractable phenolic content was analysed using the Folin–Ciocalteu assay and ranges from 41 to 142 mg g^{-1} of apple peel extract. The concentration of TPC is greater in the apple peel than that in the apple pulp obtained in a previous research study from Aragón and Navarra apple cultivars,²⁴ and this result is consistent with other comparative studies between the peel and the pulp of the apple.²⁵

The Folin–Ciocalteu assay method is usually used as a normalized method for analysing the TPC and uses the Folin–Ciocalteu reagent to oxidise the phenolic compounds. These phenolic compounds analysed are the main objective of this study, and the amounts of these bioactive compounds in the apple peel extracts are relevant to revalorizing these types of wastes and by-products. However, the TPC assay used is based on a redox reaction, and other compounds than polyphenols such as ascorbic acid and sugars can reduce the Folin–Ciocalteu reagent, and an overestimated result of phenolics can be produced.²⁶ Nevertheless, the TPC can be very useful in comparing the amount of phenolic compounds between the different samples. In this assay, only Amarilla de Octubre has a significant difference among the other samples. Therefore, a complementary study of the antioxidant phytochemicals was needed, and HPLC-MS was selected as a reference method to analyse the overall phenolic compounds and 37 individual phenolic compounds of the samples. The TPC obtained by HPLC-MS was lower than that obtained by the Folin–Ciocalteu method. This finding is attributed to the fact that only 37 phenolic compounds were detected in this analysis and that many compounds such as polymerized polyphenols cannot be detected by HPLC-MS, such as proanthocyanidins (flavan-3-ol)

Table 3 Results of antidiabetic activities against α -glucosidase and α -amylase. AGE formation and antioxidant activity against DPPH. The results are presented as the mean \pm SEM of a minimum of three independent experiments. IC_{50} values were also calculated by nonlinear regression (ns = not significant, * $p < 0.05$, ** $p < 0.001$, *** $p < 0.0005$, **** $p < 0.0001$ versus the control)

Extracts	α -Glucosidase	α -Amylase	AGEs	DPPH
A. Octubre	625.51 \pm 52.23 ^{ns}	1796.51 \pm 40.95****	158.69 \pm 10.04 ^{ns}	23.78 \pm 3.69 ^{ns}
M. Helada	1074 \pm 77.53 ^{ns}	n.a	190.95 \pm 8.14 ^{ns}	51.68 \pm 5.13*
E. Ademuz	n.a	n.a	383.41 \pm 14.18***	56.09 \pm 8.23*
Pinova	2150 \pm 926.49*	n.a	1006.87 \pm 82.32****	92.43 \pm 10.80****
V. Doncella	1250 \pm 63.51 ^{ns}	n.a	221.61 \pm 14.12*	59.98 \pm 3.08*
Borau 01	n.a	n.a	378.33 \pm 14.90****	106.45 \pm 21.88****
Acarbose	220 \pm 64.10	—	—	—
Gallic acid	—	909 \pm 15	—	0.47 \pm 0.07
Aminoguanidine	—	—	55.01 \pm 5.63	—

n.a: no activity.






Fig. 5 Correlation statistical analysis of TPC using the Folin–Ciocalteu assay and IC_{50} values for α -glucosidase (A), AGEs (B) and DPPH (C) (ns = not significant, * $p < 0.05$).

and other oligomeric phenolics.²⁷ Nevertheless, the individual phenolic groups found in the extracts were flavonols, flavan-3-ol, anthocyanins, dihydrochalcones and cinnamic acids, which are similar to the results of other previous studies on apple peels.¹⁹

Among all samples, the autochthonous sample of Amarilla de Octubre was the singular sample with the highest amount of TPC, contrasted by the Folin–Ciocalteu assay and with the

identification of the major individual phenolic compounds by HPLC-MS. These results align with the best antidiabetic and antioxidant activities in all bioassays, showing no significant difference with all the reference substances, apart from the lipase and α -amylase assays.

In the other apple peel extracts, no significant differences were found in the TPC using the Folin–Ciocalteu assay, which



indicates a low variability in the total polyphenols between different apple peel varieties. These results are in contrast with the previous studies of the TPC using the Folin–Ciocalteu assay in the apple pulp from commercial and autochthonous varieties where a significant difference was found in the TPC between them.²⁴ Nevertheless, the ratio of the 37 individual phenolic compounds analysed by HPLC-MS showed a certain variability in the phenolic groups between the samples.

The autochthonous samples of Amarilla de Octubre followed by Manzana Helada showed the highest amount of flavonols and anthocyanins among all the autochthonous samples, combined with greater amounts of flavan-3-ols or dihydrochalcones compounds, respectively, compared to the commercial samples. These findings agree with the lowest IC₅₀ value in the inhibition of α -glucosidase, formation of AGEs and DPPH radical scavenging, among all extracts. Both extracts showed similar results to acarbose and aminoguanidine in the antidiabetic assays, but only Amarilla de Octubre exhibited a similar activity to gallic acid in the inhibition of DPPH and was able to inhibit the α -amylase enzyme. The commercial sample of Verde Doncella showed great results as well in the antidiabetic and antioxidant bioassays with similar activity to acarbose in the inhibition of α -glucosidase, and it is attributable to the high amounts of flavonols and anthocyanins, along with a significant amount of flavan-3-ols compared to the other commercial samples. Nevertheless, none of the extracts were able to achieve the inhibition of lipase, and none of the extracts apart from Amarilla de Octubre inhibited the α -amylase enzyme.

Correlations for the TPC using the Folin–Ciocalteu assay and the α -glucosidase or advanced glycation end-product assays were not found, which could mean that other phytochemicals might be acting as bioactive compounds in these types of properties. Previous studies suggest that triterpenes found in apples, such as oleanolic and ursolic acids, may act as α -glucosidase inhibitors.^{28,29} Therefore, the ratio of the individual phenolic compounds and their synergism can not only influence the antidiabetic activity, but also the antioxidant activity.³⁰ Some of these individual phenolic compounds found in high quantities in the samples were glycosylated flavonoids (hyperoside, delphinidin 3,5-diglucoside, and phlorizin), which modify the inhibitory activity of the α -amylase and α -glucosidase enzymes depending on the conjugation site and the type of sugar.³¹ In addition, the glycosylated chemical structure produces higher water solubility with higher bioavailability, which is a common issue of the phenolic compounds.³²

5. Conclusion

Six apple peel samples from local and commercial cultivars were explored in terms of phytochemicals and can be considered a rich source of phenolic compounds (particularly flavonols, anthocyanins, flavan-3-ols, dihydrochalcones, and cinnamic acids). The samples showed antidiabetic and anti-

oxidant properties without altering the cell viability at physiological concentrations. These bioassays indicated that the autochthonous local cultivar known as Amarilla de Octubre showed the highest phenolic content and the lowest IC₅₀ value in the bioassays, suggesting its potential use for the development of nutraceuticals or pharmaceutical products for the prevention or treatment of type-2 diabetes complications as well as for the extraction of certain phytochemicals. The most representative individual phenolic compounds found in the apple peel samples are glycosylated flavonols such as hyperoside and isoquercitrin, flavan-3-ols such as procyanidin B2 and epicatechin, or anthocyanins such as delphinidin 3,5-diglucoside.

Abbreviations

AGEs	Advanced glycation end products
DNS	Dinitrosalicylic acid
DPPH	2,2-Diphenyl-1-picrylhydrazyl
ESI	Electrospray ionization
FAO	Food and Agriculture Organization of the United Nations
FBS	Foetal bovine serum
GAE	Gallic acid equivalent
HPLC-MS/MS	High performance liquid chromatography coupled with mass spectrometry
IC ₅₀	Half-maximal inhibitory concentration
IDF	International Diabetes Federation
NPB	<i>p</i> -4-Nitrophenyl butyrate
ROS	Reactive oxygen species
SEM	Standard error of mean
SLR	Solid–liquid ratio
TPC	Total phenolic content
UAE	Ultrasound-assisted extraction
WHO	World Health Organization

Data availability

The authors confirm that the data supporting the findings of this study are available within the article. Raw data and derived data supporting the findings of this study will be available from the corresponding author upon request.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work is part of the APPLIEDIV Project (*Multidisciplinary approach for exploiting genetic diversity to increase the value of autochthonous apple: valorizing local apple by-products and wastes through bioactive compounds*; reference



PID2022-141847OR-C33) funded by the Research State Agency in the 2022 call for Projects of Generation Knowledge oriented towards societal challenges and the previous project APPLICUT (PID2019-108081RR-C21) founded in the 2019 call of the Research State Agency. The authors thank Teva Pharma S.L.U for the research PhD grant to Javier Cano-Lou, the Government of Aragon for financial support from the Phyto-pharm group (ref. B44_23R) and Universidad San Jorge for funding Internal Project 2425013 through the Call for Internal Research Projects (Academic Year 2024–2025).

References

- 1 Y. Zheng, S. Ley and F. Hu, Global aetiology and epidemiology of type 2 diabetes mellitus and its complications, *Nat. Rev. Endocrinol.*, 2018, **14**(2), 88–98.
- 2 E. Rendra, V. Riabov, D. M. Mossel, T. Sevastyanova, M. C. Harmsen and J. Kzhyshkowska, Reactive oxygen species (ROS) in macrophage activation and function in diabetes, *Immunobiology*, 2019, **224**(2), 242–253.
- 3 World Health Organization, WHO [Internet]. Health topics: Diabetes [cited 2024 Oct 15]. Available from: https://www.who.int/health-topics/diabetes#tab=tab_1.
- 4 H. Sun, P. Saeedi, S. Karuranga, M. Pinkepank, K. Ogurtsova, B. B. Duncan, *et al.*, IDF Diabetes Atlas: Global, regional and country-level diabetes prevalence estimates for 2021 and projections for 2045, *Diabetes Res. Clin.*, 2022, 183–109119.
- 5 T. Geng, K. Zhu, Q. Lu, Z. Wan, X. Chen, L. Liu, *et al.*, Healthy lifestyle behaviors, mediating biomarkers, and risk of microvascular complications among individuals with type 2 diabetes: A cohort study, *PLoS Med.*, 2023, **20**(1), e1004135.
- 6 V. K. Sullivan, H. Kim, L. E. Caulfield, L. M. Steffen, E. Selvin and C. M. Rebholz, Plant-Based Dietary Patterns and Incident Diabetes in the Atherosclerosis Risk in Communities (ARIC) Study, *Diabetes Care*, 2024, **47**(5), 803–809.
- 7 K. A. Amirehsani, J. Hu, D. C. Wallace and T. P. McCoy, Herbal/Plant Remedies and Supplements Used by Hispanics/Latinxs for Diabetes: Source of Functional Foods?, *Sci. Diabetes Self-Manage. Care.*, 2021, **47**(1), 94–104.
- 8 M. Guasch-Ferré, J. Merino, Q. Sun, M. Fitó and J. Salas-Salvadó, Dietary Polyphenols, Mediterranean Diet, Prediabetes, and Type 2 Diabetes: A Narrative Review of the Evidence, *Oxid. Med. Cell. Longevity*, 2017, **2017**, 6723931.
- 9 C. Manach, A. Scalbert, C. Morand, C. Rémésy and L. Jiménez, Polyphenols: food sources and bioavailability, *Am. J. Clin. Nutr.*, 2004, **79**(5), 727–747.
- 10 D. De Paulo Farias, F. F. de Araújo, I. A. Neri-Numa and G. M. Pastore, Antidiabetic potential of dietary polyphenols: A mechanistic review, *Food Res. Int.*, 2021, **145**, 110383.
- 11 J. J. C. Serina and P. C. M. F. Castilho, Using polyphenols as a relevant therapy to diabetes and its complications, a review, *Crit. Rev. Food Sci. Nutr.*, 2022, **62**(30), 8355–8387.
- 12 S. P. Sha, D. Modak, S. Sarkar, S. K. Roy, S. P. Sah, K. Ghatani, *et al.*, Fruit waste: a current perspective for the sustainable production of pharmacological, nutraceutical, and bioactive resources, *Front. Microbiol.*, 2023, **14**, 1260071.
- 13 N. P. Nirmal, A. C. Khanashyam, A. S. Mundanat, K. Shah, K. S. Babu, P. Thorakkattu, *et al.*, Valorization of Fruit Waste for Bioactive Compounds and Their Applications in the Food Industry, *Foods*, 2023, **12**(3), 556.
- 14 J. Gustavsson, C. Cederberg, U. Sonesson, R. van Otterdijk and A. Meybeck, *Global food losses and food waste - Extent, causes and prevention*, En: FAO. Save Food! Congress, Düsseldorf. Rome, 2011.
- 15 Ministerio de Agricultura, Pesca y Alimentación, *Informe del Desperdicio Alimentario en España-Octubre 2023*, Dirección General de Alimentación, Madrid, 2023.
- 16 R. De la Peña-Armada and I. Mateos-Aparicio, Sustainable Approaches Using Green Technologies for Apple By-Product Valorisation as A New Perspective into the History of the Apple, *Molecules*, 2022, **27**(20), 6937.
- 17 B. Dinda, M. Dinda, A. Roy and S. Dinda, Dietary plant flavonoids in prevention of obesity and diabetes, *Adv. Protein Chem. Struct. Biol.*, 2020, **120**, 159–235.
- 18 C. M. Weaver, Bioactive foods and ingredients for health, *Adv. Nutr.*, 2014, **5**(3), 306S–311S.
- 19 J. Popiolek-Kalisz and P. Glibowski, Apple Peel Supplementation Potential in Metabolic Syndrome Prevention, *Life*, 2023, **13**(3), 753.
- 20 M. Kalinowska, A. Bielawska, H. Lewandowska-Siwkiewicz, W. Priebe and W. Lewandowski, Apples: content of phenolic compounds vs. variety, part of apple and cultivation model, extraction of phenolic compounds, biological properties, *Plant Physiol. Biochem.*, 2014, **84**, 169–188.
- 21 B. Łata, A. Trampczynska and J. Paczesna, Cultivar variation in apple peel and whole fruit phenolic composition, *Sci. Hortic.*, 2009, **121**(2), 176–181.
- 22 R. C. Pimpão, T. Dew, P. B. Oliveira, G. Williamson, R. B. Ferreira and C. N. Santos, Analysis of phenolic compounds in Portuguese wild and commercial berries after multienzyme hydrolysis, *J. Agric. Food Chem.*, 2013, **61**(17), 4053–4062.
- 23 A. M. Mustafa, S. Angeloni, D. Abouelenein, L. Acquaticci, J. Xiao, G. Sagratini, *et al.*, A new HPLC-MS/MS method for the simultaneous determination of 36 polyphenols in blueberry, strawberry and their commercial products and determination of antioxidant activity, *Food Chem.*, 2022, **367**, 130743.
- 24 A. Millán-Laleona, F. J. Bielsa, E. Aranda-Cañada, C. Gómez-Rincón, P. Errea and V. López, Antioxidant, Antidiabetic, and Anti-Obesity Properties of Apple Pulp Extracts (*Malus domestica* Bork): A Comparative Study of 15 Local and Commercial Cultivars from Spain, *Biology*, 2023, **12**(7), 891.



- 25 K. Wolfe, X. Wu and R. H. Liu, Antioxidant activity of apple peels, *J. Agric. Food Chem.*, 2003, **51**(3), 609–614.
- 26 S. Sultana, K. Foster, T. Bates, M. L. Hossain, L. Y. Lim, K. Hammer, *et al.*, Determination of Physicochemical Characteristics, Phytochemical Profile and Antioxidant Activity of Various Clover Honeys, *Chem. Biodivers.*, 2024, **21**(5), e202301880.
- 27 O. Vidal-Casanella, J. Moreno-Merchan, M. Granados, O. Nuñez, J. Saurina and S. Sentellas, Total Polyphenol Content in Food Samples and Nutraceuticals: Antioxidant Indices versus High Performance Liquid Chromatography, *Antioxidants*, 2022, **11**(2), 324.
- 28 H. Hussain, I. R. Green, I. Ali, I. A. Khan, Z. Ali, A. M. Al-Sadi, *et al.*, Ursolic acid derivatives for pharmaceutical use: a patent review (2012–2016), *Expert Opin. Ther. Pat.*, 2017, **27**(9), 1061–1072.
- 29 S. T. Carginin and S. B. Gnoatto, Ursolic acid from apple pomace and traditional plants: A valuable triterpenoid with functional properties, *Food Chem.*, 2017, **220**, 477–489.
- 30 W. J. Yeh, S. M. Hsia, W. H. Lee and C. H. Wu, Polyphenols with antiglycation activity and mechanisms of action: A review of recent findings, *J. Food Drug Anal.*, 2017, **25**(1), 84–92.
- 31 H. Cao and X. Chen, Structures required of flavonoids for inhibiting digestive enzymes, *Anticancer Agents Med. Chem.*, 2012, **12**(8), 929–939.
- 32 M. Fujimori, K. Kadota, K. Kato, Y. Seto, S. Onoue, H. Sato, *et al.*, Low hygroscopic spray-dried powders with trans-glycosylated food additives enhance the solubility and oral bioavailability of ipriflavone, *Food Chem.*, 2016, **190**, 1050–1055.

