


NJC

New Journal of Chemistry

Accepted Manuscript

A journal for new directions in chemistry

This article can be cited before page numbers have been issued, to do this please use: M. Yanbaeva, T. Steinke, E. Engelage, R. Weiss and S. Huber, *New J. Chem.*, 2025, DOI: 10.1039/D5NJ03660G.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

COMMUNICATION

“Anti-Electrostatic” (Inter-Anion) Chalcogen Bonding Interactions

Margarita Yanbaeva^{[a]†}, Tim Steinke^{[a]†}, Elric Engelage^[a], Robert Weiss^{[b]*}, and Stefan M. Huber^{[a]*}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Dedicated to Professor Resnati, celebrating a career in fluorine and noncovalent chemistry on the occasion of his 70th birthday.

In this work, the ability of 1,2-bis(dicyanomethylene)-cyclopropanide substituents to enable inter-anion interactions has been expanded to chalcogen bonding. An anionic selenium-based chalcogen-bond donor featuring this ligand displays “anti-electrostatic” interactions in the solid state. Three polymorphic structures of associates with both mono- and biaxial binding modes are discussed.

Noncovalent interactions (NCIs) such as halogen¹ (XB), chalcogen² (ChB) and pnictogen³ (PnB) bonding have received increased interest in the last decades, and based on the anisotropic electrostatic potential of the interacting atom, they have sometimes collectively been termed “σ-hole interactions”.⁴ These NCIs have established themselves as a flexible tool in various fields like crystal engineering⁵, catalysis⁶, ion recognition⁷ and biochemistry⁸. Electrostatics are considered to play a crucial role in the formation of these bonds,⁹ however, components contributing to the interaction energies are not limited to simple electrostatic attraction.¹⁰ Several other factors must be taken into account, including dispersion, polarization or charge-transfer.^{11†} “Anti-electrostatic” interactions between ions of like charge represent one of the key examples demonstrating the importance of the latter contributions.

In fact, several studies on this counterintuitive phenomenon have been put forth in recent years,¹² and an even broader selection of examples emerges as previously reported data is re-evaluated with this concept in mind.^{13††} Nevertheless,

experimental findings are limited in scope, especially those involving the interactions between organic molecules. One of the first examples of “anti-electrostatic” halogen bonding (AEXB) between structurally distinct species was previously reported by our group.¹⁴ The anionic iodinated cyclopropanide XB donor **1** and cyclopropanide-substituted imidazole species **2** (see Figure 1, top) have displayed halide adduct formation and self-association, respectively.




Figure 1. Top: Previously reported cases of “anti-electrostatic” halogen bonded aggregates of 1,2-bis(dicyanomethylene)-cyclopropanide-containing synthons in the solid state: **A** – 2:1 anion binding and **B** – trimer formation. For **A**: $d_{I...I} = 85\% \Sigma r_w$, in **B**: $d_{I...N} = 79\text{--}81\% \Sigma r_w$. Cations omitted for clarity. Bottom: Known cyclopropanide- and cyclopropenium-substituted chalcogen compounds. Counterions omitted for clarity.

[a] M. Yanbaeva, Dr. T. Steinke, Dr. E. Engelage, Prof. Dr. S. M. Huber
Fakultät für Chemie und Biochemie, Ruhr-Universität Bochum
Universitätsstraße 150, 44801 Bochum (Germany)
E-Mail: stefan.m.huber@rub.de

[b] Prof. Dr. R. Weiss
Institut für Organische Chemie, Friedrich-Alexander-Universität Erlangen-
Nürnberg
Henkestraße 42, 91054 Erlangen (Germany)
E-Mail: robert.weiss@chemie-uni-erlangen.de

[†] These authors contributed equally to this work.

COMMUNICATION

Journal Name


Since halogen and chalcogen bonding share many similarities, the concept of “anti-electrostatic” bonding could potentially also be expanded to ChB (“AEChB”).¹⁵ Obvious candidate structures to realize these contacts are organochalcogen compounds bearing cyclopropanide moieties analogously to the reported halogen-bond donors.

Both cationic cyclopropenium- and anionic cyclopropanide-substituted selenium compounds (**3-5**) have previously been reported by Seitz and co-workers.¹⁶ The crystal structure of the mixed species **4** is literature-known; however, no further analysis of the intermolecular interactions in the solid state has been performed to date.

Following up on our previous investigations regarding “anti-electrostatic” noncovalent interactions, we herein present, to the best of our knowledge, the first example of inter-anion chalcogen bonding in the solid state.

Results and discussion

Anionic 1,2-bis(dicyanomethylene)cyclopropanide derivative **7** was prepared from commercially available trichloroethylene over four steps following previously established procedures.¹⁴ The chalcogen-bond donor **8** was subsequently synthesized from phenylselenenyl chloride with 85% yield. This reaction further underlines the autouploading¹⁷ properties of **7**, as the transformation likely occurs *via* a nucleophilic attack of the ambiphilic carbon of the cyclopropanide moiety on the selenium (s. Scheme 1).



During crystallization experiments with compound **8** without further additives, two different polymorphic structures of supramolecular aggregates could be identified. Crystals suitable

for analysis were obtained *via* vapor diffusion from THF with cyclohexane as antisolvent, or *via* slow evaporation from DCM, with the individual structures discussed below.

A summary of mean bond lengths of all unique instances of **8** as well as values obtained from gas phase geometry optimizations on different levels of theory are presented in Figure 2. Overall, high agreement between experimental and theoretical data can be observed. The C-C distances are representative of a highly conjugated structure, as the values are close to those present in benzene (1.390 Å).¹⁸ Notably, the C2-C3 bond is slightly longer at 1.411 Å, while the Se1-C1 distance is, in turn, shorter than a typical carbon-selenium single bond.¹⁹ The former observation is more in line with the right resonance structure of Scheme 1B, while the latter seems to indicate a contribution from the left one. The short Se-C distance could also be indicative of an n→σ* interaction of the anionic carbon into the Se-C_{Ph} σ* orbital, though.





Figure 2. Selected average bond lengths of **8**. Ellipsoids set at 50% probability. TDA cations omitted for clarity. Geometry optimization performed with B3LYP²⁰ and M06-2X²¹ functionals, applying def2TZVP(D)²² basis set.

The electronic nature of the chalcogen-bond donor was further investigated with the help of an electrostatic potential (ESP) map, which was calculated at B3LYP/def2TZVP(D) level of theory. Overall, a rather homogeneous distribution of the negative charge is observed, with no areas with positive potential present (see Figure 3). Nevertheless, a region with relatively lower electron density (blue) is seen opposite to the cyclopropanide substituent, while no such area is visible along the phenyl substituent's axis. The cyano groups feature the most negative electrostatic potential (displayed in red) and are thus capable of acting as chalcogen-bond acceptors.



Additionally, total energy profiles of the Cl...Se distance scans were plotted at the M06-2X/def2TZVP(D) level of theory to evaluate the thermodynamics of a potential non-isostructural interaction between the anion of **8** and chloride (Figure 3). The formation of the adduct in both gas phase and acetonitrile is endothermic; a result expected from an anion-anion interaction. Still, well depths (the differences between the energies of the complex at minima and extended distances) of -11.6 kJ/mol and -5.1 kJ/mol are observed in the gas phase and

in acetonitrile solution, respectively. These minima are located at 3.1 Å (85% vdW) and 3.2 Å (88% vdW); distances that are typical for a chalcogen-bonded adduct.

In an attempt to obtain chalcogen bonding complexes with halides, co-crystallization experiments with one equivalent of tris(dimethylamino)-cyclopropenyl cation (TDACl) were repeated under the same conditions used previously. These, however, only yielded an additional polymorph of **8**, as well as isolated TDACl single crystals.




Figure 4: Single crystal X-ray structure of **8**: **A** - Biaxial binding; **B**, **C** - monoaxial binding. The sum of van-der-Waals radii for selenium and nitrogen amounts to 3.45 Å. TDA cations omitted for clarity. Ellipsoids set at 50% probability. In both cases of monoaxial binding, the formation of the chalcogen-bond is observed in the elongation of the cyclopropanide-selenium bond. In **A**, suitable interactions are detected along both axes. The bond formed in the elongation of the cyclopropanide substituent is significantly shorter. A summary of observed bond lengths and angles can be found in Table 1.

Crystal structures of the three polymorphs of **8** are depicted in Figure 4. A major difference between ChB and neutral XB donors is the ability of the former to form non-covalent bonds along both substituents' axes, so-called biaxial binding. Still,

adduct formation is often only observed along one of these, usually opposite the more electronegative substituent.²³ Here, both modes are observed.

COMMUNICATION

Journal Name

From the conceptual standpoint, polymorph **A**, which was obtained from slow evaporation from DCM, provides the most information due to its biaxial binding motif. This tetraanionic aggregates presented in Figure 4A display the longest chain formed within the crystal structure. As seen from the atom labelling, the anions in the smallest asymmetrical unit represent half of that chain, and all the anionic moieties are a part of those superstructures (see Figures S7-S8 in Supporting Information). Due to packing effects, no chalcogen bond is formed in the elongation of both axes of the terminal Se atoms. Those molecules are bound to anions of neighbouring chains by weak N-H contacts (Figure S10 in Supporting Information). The corresponding cations are located above and below the cyclopropanide moieties, displaying π - π interactions and short contacts. For the inner moieties, a direct comparison between the chalcogen bond lengths along the different substituents' axes can be drawn. A shortening of about 6% of the sum of vdW radii is observed for the cyclopropanide axis compared to the phenyl one despite the net negative charge of the former. Additionally, this structure contains the shortest chalcogen bond amongst the herein reported, corresponding to 90% vdW radii (see Table 1 for a summary of relevant parameters). The monoaxial binding in structures **B** and **C** only occurs along the elongation of the selenium-cyclopropanide bond, which is in line with the differences of ChBs in **A** discussed above. It indicates that the cyclopropanide substituent, despite its anionic nature, creates the superior ChB axis. The dimeric structure of **B** was obtained from vapor diffusion experiments with THF/cyclohexane. All the selenium-containing moieties partake in the dimer formation, and no other interactions apart from these short contacts and alternating π stacking between the cations and the cyclopropanide part of the anions can be observed. The latter form offset columns (see Supporting Information, Figures S11-S12). The chalcogen bond here amounts to 95% vdW, comparably longer than the one observed in structure **A** for the cyclopropanide substituent. Additionally, although the orientation of the molecules would suggest a rather symmetric motif in structure **B** with both selenium atoms forming a bond with the corresponding nitrogens, a slight distortion is present. The distance of the second Se-N pair amounts to 3.507 Å, 101% vdW, at or beyond the border of a chalcogen bond. A slight offset between the neighboring planes results in a relatively low directionality of the chalcogen bonds.

Table 1. Selected geometric parameters of the chalcogen bonds in the crystal structures presented in Figure 3. Most favourable values marked in bold.

Bond	dSe...N, [Å]	% Σr_w	$\angle$ C-Se...N, [°]
A _a (Se2 ...N2)	3.110	90	168.7
A _b (Se2 ...N8)	3.318	96	170.4
B (Se1 ...N6)	3.265	95	154.6
C (Se2 ...N11)	3.282	95	172.4

The final polymorph **C** was obtained from gas diffusion experiments with THF/cyclohexane with 1 eq. of TDACl additive.

Remarkably, only one of the individual anion pairs in the smallest asymmetrical unit is displaying chalcogen bonding (see Figure S13). Here, a Se-N bond distance of 95% vdW is observed. This is the most directional interaction amongst the observed structures, with a C-Se-N angle of 172.4°. A third moiety, which does not display any intermolecular interactions besides short contacts towards the hydrogens of the adjacent phenyl groups of other anions and π stacking with the TDA cations, is present. Similarly to the other structures, an alternating pattern in the π -stacking between the positively and negatively charged fragments is observed.

While no experimental data on "anti-electrostatic" chalcogen bonding is reported to the best of our knowledge, numerous studies involving neutral and cationic donors in the solid state have been presented over the years, with individual results,²⁴ as well as overarching data bank surveys²⁵ being available. The values observed in the anionic systems presented above are comparable with those literature-known parameters - despite their unfavourable electrostatic contribution.

Conclusions

In conclusion, we present a novel chalcogen-bond donor based on the 1,2-bis(dicyanomethylene)cyclopropanide moiety, which is capable of inter-anion bonding. Three polymorphs featuring varying self-aggregation patterns could be described. The interaction parameters are comparable to previously reported complexes of dicationic selenium-based donors with neutral and cationic Lewis bases. Since selenium is known to form weaker chalcogen bonds compared to its heavier homologue, we anticipate that the preparation of cyclopropanide-substituted tellurium donors would allow to fully realize the potential of AEChB. These studies are currently ongoing.

Conflicts of interest

There are no conflicts to declare.

Data availability

Additional data on experimental procedures, spectra and computational details is presented in the supporting information. Crystal structures are available for download in CCDC, under repository numbers 2477888, 2477889 and 2477887.

Notes and references

* While we prefer to call the contribution arising from the overlap of orbitals „charge-transfer“, it has also been postulated that its effect can solely be explained by polarization.²⁶ This discussion is outside the scope of this study.

** For an early example of an inter-cation AEChB, see: R. Laitinen, R. Steudel, R. Weiss, *J. Chem. Soc., Dalton Trans.* **1986**, 1095.

- 1 P. R. Varadwaj, A. Varadwaj, H. M. Marques and K. Yamashita, *Crystal Growth & Design*, 2024, **24**, 5494–5525.
- 2 C. B. Aakeroy, D. L. Bryce, G. R. Desiraju, A. Frontera, A. C. Legon, F. Nicotra, K. Rissanen, S. Scheiner, G. Terraneo, P. Metrangolo and G. Resnati, *Pure and Applied Chemistry*, 2019, **91**, 1889–1892.
- 3 G. Resnati, D. L. Bryce, G. R. Desiraju, A. Frontera, I. Krossing, A. C. Legon, P. Metrangolo, F. Nicotra, K. Rissanen, S. Scheiner and G. Terraneo, *Pure and Applied Chemistry*, 2024, **96**, 135–145.
- 4 T. Clark, M. Hennemann, J. S. Murray and P. Politzer, *Journal of Molecular Modeling*, 2007, **13**, 291–296.
- 5 a) P. Metrangolo and G. Resnati, *Crystal Growth & Design*, 2012, **12**, 5835–5838; b) R. Siddiqui, J. Rani, H. M. Titi and R. Patra, *Coordination Chemistry Reviews*, 2024, **517**, 215994;
- 6 D. Jovanovic, M. Poliyodath Mohanan and S. M. Huber, *Angewandte Chemie International Edition*, 2024, **63**, e202404823.
- 7 G. Berger, P. Frangville and F. Meyer, *Chemical Communications*, 2020, **56**, 4970–4981.
- 8 M. G. Walker, C. G. Mendez and P. S. Ho, *Chemistry – An Asian Journal*, 2023, **18**, e202300026.
- 9 P. Politzer, J. S. Murray and T. Clark, *Physical Chemistry Chemical Physics*, 2010, **12**, 7748–7757.
- 10 a) P. Politzer, P. Lane, M. C. Concha, Y. Ma and J. S. Murray, *Journal of Molecular Modeling*, 2007, **13**, 305–311; b) P. Politzer, J. S. Murray and M. C. Concha, *Journal of Molecular Modeling*, 2007, **13**, 643–650; c) L. de Azevedo Santos, T. C. Ramalho, T. A. Hamlin and F. M. Bickelhaupt, *Chemistry – A European Journal*, 2023, **29**, e202203791; d) A. Lara, A. Sterling, A. Aldossary, M. Head-Gordon, 2024, doi:10.26434/chemrxiv-2024-t26hd;
- 11 a) H. A. Bent, *Chemical Reviews*, 1968, **68**, 587–648; b) A. E. Reed, F. Weinhold, R. Weiss and J. Macheleid, *The Journal of Physical Chemistry*, 1985, **89**, 2688–2694;
- 12 a) C. Wang, Y. Fu, L. Zhang, D. Danovich, S. Shaik and Y. Mo, *Journal of Computational Chemistry*, 2018, **39**, 481–487; b) Z. Zhu, G. Wang, Z. Xu, Z. Chen, J. Wang, J. Shi and W. Zhu, *Physical Chemistry Chemical Physics*, 2019, **21**, 15106–15119;
- 13 J. M. Holthoff, R. Weiss, S. V. Rosokha and S. M. Huber, *Chemistry – A European Journal*, 2021, **27**, 16530–16542.
- 14 J. M. Holthoff, E. Engelage, R. Weiss and S. M. Huber, *Angewandte Chemie International Edition*, 2020, **59**, 11150–11157.
- 15 D. Fan, L. Chen, C. Wang, S. Yin and Y. Mo, *The Journal of Chemical Physics*, 2021, **155**, 234302.
- 16 R. Allmann, F.-J. Kaiser, M. Krestel and G. Seitz, *Angewandte Chemie International Edition in English*, 1986, **25**, 183–184.
- 17 R. Weiss, S. Reichel, M. Handke and F. Hampel, *Angewandte Chemie International Edition*, 1998, **37**, 344–347.
- 18 I. Heo, J. C. Lee, B. R. Özer and T. Schultz, *The Journal of Physical Chemistry Letters*, 2022, **13**, 8278–8283.
- 19 P. Wonner, L. Vogel, M. Düser, L. Gomes, F. Kniep, B. Mallick, D. B. Werz and S. M. Huber, *Angewandte Chemie International Edition*, 2017, **56**, 12009–12012.
- 20 A. D. Becke, *The Journal of Chemical Physics*, 1993, **98**, 5648–5652.
- 21 Y. Zhao and D. G. Truhlar, *Theoretical Chemistry Accounts*, 2008, **120**, 215–241.
- 22 P. J. Stephens, F. J. Devlin, C. F. Chabalowski and M. J. Frisch, *The Journal of Physical Chemistry*, 1994, **98**, 11623–11627.
- 23 S. Scheiner, *ChemPhysChem*, 2023, **24**, e202200936.
- 24 a) T. Steinke, E. Engelage and S. M. Huber, *Acta Crystallographica Section C*, 2023, **79**, 26–35; b) P. Wonner, L. Vogel, M. Düser, L. Gomes, F. Kniep, B. Mallick, D. B. Werz and S. M. Huber, *Angewandte Chemie International Edition*, 2017, **56**, 12009–12012;
- 25 a) A. Frontera and A. Bauzá, *Crystals*, 2021, **11**; b) E. R. Tiekink, *Coordination Chemistry Reviews*, 2021, **427**, 213586;
- 26 T. Clark and A. Heßelmann, *Physical Chemistry Chemical Physics*, 2018, **20**, 22849–22855.

The data supporting this article have been included as part of the Supplementary Information.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Open Access Article. Published on 18 October 2025. Downloaded on 10/24/2025 12:06:31 AM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

