Energy & Environmental Science



View Article Online

CORRECTION

Check for updates

Cite this: Energy Environ. Sci., 2022, 15, 3097

Correction: Local reaction environment for selective electroreduction of carbon monoxide

Ming Ma,*^{ab} Wanyu Deng,^b Aoni Xu,^b Degenhart Hochfilzer,^b Yu Qiao,^b Karen Chan,*^b Ib Chorkendorff^b and Brian Seger*^b

DOI: 10.1039/d2ee90030k

Correction for 'Local reaction environment for selective electroreduction of carbon monoxide' by Ming Ma et al., Energy Environ. Sci., 2022, https://doi.org/10.1039/d1ee03838a.

rsc.li/ees

In our work we discussed that the key step for acetate formation should occur near the cathode surface, which was based on ref. 10. During the analysis we became aware of the theoretical effort by Heenen *et al.*¹ that hypothesized this key step was a solution reaction in Scheme S1 (ESI, ref. 7). While we cited this work in the supplementary information, we failed to acknowledge the work from Heenen *et al.* in the main text and feel this needs to be corrected.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

References

1 H. H. Heenen, H. Shin, G. Kastlunger, S. Overa, J. A. Gauthier, F. Jiao and K. Chan, Mechanism for acetate formation in electrochemical CO(2) reduction on Cu: Selectivity with potential, pH, and nanostructuring, *ChemRxiv*, 2022, preprint, DOI: 10.26434/chemrxiv-2021-p3d4s-v3.

^a School of Chemical Engineering and Technology, Xi'an Jiaotong University, Xi'an 710049, People's Republic of China. E-mail: mingma@xjtu.edu.cn

^b Department of Physics, Technical University of Denmark, 2800 Kgs. Lyngby, Denmark. E-mail: kchan@fysik.dtu.dk