

Cite this: *Chem. Sci.*, 2024, 15, 19936

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 19th June 2024
Accepted 2nd November 2024

DOI: 10.1039/d4sc04038d

rsc.li/chemical-science

Dual ligand-enabled iron and halogen-containing carboxylate-based photocatalysis for chloro/fluoro-polyhaloalkylation of alkenes†

Wanru Han,^{‡a} Zhenyan Zhao,^{‡a} Kui Jiang,^a Yu Lan,^{‡abc} Xuehan Yu,^a Xiaoyu Jiang,^a Wei Yang,^a Donghui Wei,^{‡ab} Shi-Jun Li^{‡ab} and Linbin Niu^{*ab}

Herein, we demonstrate a practical dual ligand-enabled iron photocatalysis paradigm—converting all kinds of halogen-containing carboxylates ($C_nX_mCOO^-$, X: F, Cl, Br) into C_nX_m radicals for the valuable chloro/fluoro-polyhaloalkylation of non-activated alkenes with easily available trichloroacetonitrile/Selectfluor as the electrophilic halogenation reagent. The modular *in situ* assembly of the effective iron and $C_nX_mCOO^-$ -based light-harvesting species using the two ligands—OMe/CF₃-substituted bipyridine and acetonitrile/trichloroacetonitrile is evidenced by detailed mechanistic studies. The late-stage modification, low loading amount of iron (TON: 257) and feasible gram-scale synthesis show the utility of this protocol. We thus anticipate that the dual ligand-enabled iron photocatalysis paradigm may facilitate activation and transformation of inert bulk chemicals.

Introduction

Using abundant bulk chemicals to synthesize fine chemicals through a sustainable catalytic strategy is significant in the field of synthetic chemistry.¹ Regarding the efficient activation of inert bulk chemicals, general considerations or solutions are to increase the energy input such as resorting to high temperatures,² noble metal catalysts,³ strong or special oxidants/reductants/acids/bases,⁴ UV light,⁵ etc.⁶ Due to the urgent demand for green and sustainable chemistry,⁷ balancing the requirement of reducing the energy input/consumption and the growing interest of chemists in the activation of inert chemicals is what chemists should imminently address (Fig. 1a).

The nontoxicity and abundance of iron endow iron-catalyzed synthetic chemistry with strong sustainability and potential application prospects in bio-medicine and industrial manufacturing fields.⁸ The low cost and easily available iron(III) salts have been playing versatile roles in the synthetic chemistry community owing to their remarkable Lewis acidity⁹ and redox reactivity. With respect to the redox character of iron(III), it has exhibited the ability to oxidise some active radical species¹⁰ or

electron rich compounds such as anilines¹¹ and phosphines.¹² To achieve iron-mediated oxidative activation of inert compounds (alkane as an example), strong stoichiometric oxidants are usually required to yield high valent iron(IV or V)oxo species that can dominate the activation and transformation of inert alkanes for the generation of value-added alcohols/ketones, which is a representative mode for iron-catalyzed functionalization of inert compounds (Fig. 1b).¹³ By contrast, the oxidative ability of iron(III) is relatively moderate, hence direct iron(III)-dominated oxidative activation of inert compounds is still rare.

The input of light energy into iron(III) enhances the redox activity to a certain degree.¹⁴ Photo-induced iron ligand-to-metal charge transfer (LMCT) possesses excellent ability to produce radical species from abundant chemical feedstocks like alkyl carboxylates (^tBuCOO[−] as a representative, $E_{1/2}^{ox} = +1.26$ V *versus* SCE),^{15–18} in which the necessary prerequisite is the suitable coordination of the substrate to the iron center. It means that activating those inert compounds with exceedingly weak coordination capacity in this fashion is helpless and doubtful. Very recently, through cooperating iron photocatalysis and redox-active thiol catalysis, West and Xia elegantly reported direct decarboxylation of F-containing carboxylic acids for hydro-fluoroalkylation of alkenes.¹⁹ The extremely high oxidative potential of TFA (CF₃COO[−], $E_{1/2}^{ox} > +2.4$ V *versus* SCE) and the strong acid system make the corresponding activation mechanism of this special method complicated and confusing (Fig. 1c).

To develop a general strategy for the activation and transformation of inert and weakly coordinating compounds *via* metal-based photocatalysis, we hope to choose abundant and alkaline halogen-containing carboxylates ($C_nX_mCOO^-$, X: F, Cl, Br) as the

^aCollege of Chemistry, Pingyuan Laboratory, Zhengzhou University, 100 Science Avenue, Zhengzhou 450001, Henan, China. E-mail: lanyu@cqu.edu.cn; donghuiwei@zhu.edu.cn; lishijunzong@zhu.edu.cn; nlb@zhu.edu.cn

^bState Key Laboratory of Antiviral Drugs, Pingyuan Laboratory, Henan Normal University, Xixiang 453007, Henan, China

^cSchool of Chemistry and Chemical Engineering, Chongqing Key Laboratory of Chemical Theory and Mechanism, Chongqing University, Chongqing 401331, China

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4sc04038d>

‡ These authors contributed equally to this work.




Fig. 1 Paradigms for activation of inert compounds. (a) Pursuing the activation of inert chemicals under milder conditions. (b) The redox character of iron(III). (c) Decarboxylation of C_nF_mCOOH promoted by purple light. (d) This work: dual ligand-enabled iron and halogen-containing carboxylate-based photocatalysis for chloro/fluoro-polyhaloalkylation of alkenes.

investigational object to circumvent the strong acid conditions,²⁰ which is conducive to bringing the effect of the external ligand into full play.²¹ Given the fact that the role of the external ligand in 3d metal-based LMCT chemistry is usually intricate but significant, herein, we adopted the dual ligand-enabled strategy to efficiently assemble iron and $C_nX_mCOO^-$ -based light-harvesting species for C_nX_m radical production under visible light conditions,²² thus successfully achieving chloro/fluoro-polyhaloalkylation of non-activated alkenes with easily available trichloroacetonitrile/Selectfluor as the electrophilic halogenation reagent (Fig. 1d). OMe/ CF_3 -substituted bipyridine (bpy) and acetonitrile/trichloroacetonitrile indeed serve as the crucial dual ligands to achieve the desired LMCT between $C_nX_mCOO^-$ and the iron(III) center and furnish the decarboxylation of $C_nX_mCOO^-$ under visible light rather than UV light.

Results and discussion

To quickly obtain photochemical insights into iron(III) and $C_nX_mCOO^-$ -based light-harvesting species, stoichiometric

photolysis experiments of $Fe(CF_3COO)_3$ were first carried out. As we expected, the high energy light ($\lambda_{max} = 365 \text{ nm}$) irradiation could induce the decomposition of $Fe(CF_3COO)_3$ to the CF_3 radical, which was captured by a non-activated alkene **1**, and the generated radical adduct was further quenched through the halogen atom transfer (XAT) with Cl_3CCN , delivering the chloro-trifluoromethylation product of alkene (Fig. 2a, entry 1).²³ This process failed under visible light ($\lambda_{max} = 450 \text{ nm}$) or darkness conditions (entries 2 and 3). To reduce the energy consumption in the activation of CF_3COO^- , the external ligands were investigated to achieve the release of the CF_3 radical under visible light conditions. Bpy **L1** was incapable of controlling the effective decarboxylation of CF_3COO^- to access the chloro-trifluoromethylation of alkene, whereas its derivatives functionalized with electron-donating/withdrawing functional groups (**L2** and **L3**) both showed the potential of being a competent ligand (entries 4–6).

Based on the preliminary results of the above stoichiometric experiments, a hypothetical catalytic cycle is illustrated in Fig. 2b. We proposed that a suitable electrophilic halogenation reagent (**VIII** or **IX**) should not only participate in the halogen-atom transfer (XAT) with the radical adduct (**VII**) that was generated from the C_nX_m radical (**V**) and alkene (**VI**) for the designed chloro/fluoro-polyhaloalkylation (**XIV** or **XV**), but also its concomitant electrophilic carbon/nitrogen radical species (**X** or **XI**) generated from the XAT would be responsible for the oxidation of $[Fe^{II}/L]$ species (**III**) to regenerate the $[Fe^{III}/L]$ complex (**I**). Notably, the additional effect of the ligand also possibly affected the process of $[Fe^{II}/L]$ to $[Fe^{III}/L]$, indicating that the choosing of the competent ligand is first and foremost for investigating conditions. After our detailed investigations (Fig. S14 and S15†), it was found that the chloro-trifluoromethylation and fluoro-chlorodifluoromethylation of alkenes *via* visible light-induced iron catalysis could be accomplished in good yields and regioselectivity under the regulation of **L2** and **L4**, respectively (Fig. 2c). Regarding the choosing of the halogenation reagent, it was necessary to mention the result that commonly used chlorine sources such as trichloromethane, ethyl chloroacetate analogue, NCS (*N*-chlorosuccinimide), *etc.* could not yield the desired chloro-trifluoromethylation product (Fig. S17†), and bromo-trifluoromethylation of alkene by this method was infeasible (Fig. S18†).

With the optimized conditions in hand, the scope of this protocol was explored in terms of various non-activated alkenes' chloro/fluoro-polyhaloalkylation (Fig. 3). We first examined the reactivity of benzoate-containing terminal olefins. The different alkyl carbon chain structures of olefins did not disturb the reactivity and both provided good yields (**4** and **5**). Electron-donating or electron-withdrawing substituents in benzoate groups were compatible (**6–8**), and the substituents installed at *ortho* and *meta* positions also provided satisfactory yields (**9–11**). The alkene assembled with valuable benzotriazole successfully introduced the Cl and CF_3 groups in 74% yield under our standard conditions (**12**). Moreover, sulfonate/phenoxy-modified alkenes, simple hexadecane and α -methyl olefin were all well tolerated (**13–17**). Besides the above-mentioned terminal






Fig. 2 Investigations of chloro/fluoro-polyhaloalkylation of alkenes. (a) Photo-induced iron mediated chloro-trifluoromethylation of an alkene. (b) Proposed mechanism of photo-induced iron catalyzed chloro/fluoro-polyhaloalkylation of alkenes. (c) Optimal conditions of the chloro/fluoro-polyhaloalkylation of alkenes.

olefins, the successful chloro-trifluoromethylation of the internal olefin enriched the diversity of the product (18). Significantly, the synthetic utility of this method in achieving fluoro-polyhaloalkylation²⁴ of non-activated alkenes was elaborated for the divergent synthesis of complex fluorine-containing molecules (19–25). Unfortunately, conjugated alkenes were not suitable for this reaction (Fig. S12†).

Furthermore, a wide variety of alkenes derived from pharmaceutical molecules were examined for the late-stage functionalization to extend this practical synthetic platform. The medicine oxaprozin used for rheumatoid arthritis and post-operative analgesia could be modified with halogen-containing alkyl in moderate yield (26). It is noteworthy that the performance of the derivative of piperonylic acid was amazing with 83% yield, whose sensitive ternary ring structure was well preserved (27). Likewise, ciprofibrate as an oral medicine for treating endogenous hypercholesterolemia and hypertriglyceridemia in adults, its derivative could also be achieved chloro-trifluoromethylation smoothly (28). The modification based on amino acid *N*-Boc-D-proline (29) increased the opportunities to discover late-model biomolecules. Of special note, eugenol possesses antioxidant, anticancer, anti-inflammatory, and antimicrobial activities, and its analogue gave moderate yield (30), despite the existence of the sensitive phenolic hydroxyl group. Besides, the late-stage chloro-trifluoromethylation, chloro-chlorodifluoromethylation, chloro-bromodifluoromethylation and chloro-pentafluoroethylation of loxoprofen/febuxostat/flurbiprofen derivatives demonstrated

intriguing synthesis potential for new drug research and development (31–40). Importantly, the utility and sustainability of this method were further highlighted by the gram-scale synthesis and low loading amount of the iron catalyst (TON = 257).

Based on the encouraging results of this ligand-enabled iron photocatalysis under visible light conditions, we turned our attention to the mechanistic studies. When stoichiometric radical scavenger (TEMPO/BHT) was subjected to the standard conditions (Fig. S30†), the desired chloro-trifluoromethylation process was severely inhibited, in which the radical adduct of the CF_3 radical and BHT was detected by electrospray ionization high-resolution mass spectrometry (ESI-HRMS, Fig. S31†). The existence of CF_3 radical species was also evidenced by the designed radical probe experiment (Fig. S32†). Theoretically, the production of the C_nX_m radical required the excitation of iron and $\text{C}_n\text{X}_m\text{COO}^-$ -based light-harvesting species, meaning that continuous blue light irradiation should be necessary. The light on/off experiments verified this viewpoint (Fig. 4a). According to the fact that the combination of $\text{Fe}(\text{OTf})_3$ and L2 showed first-order dependence during the kinetic studies (Fig. 4b), we realized that how to assemble iron salt/L2, $\text{C}_n\text{X}_m\text{COO}^-$ and other potential ligands in this system into iron-based light-harvesting species for C_nX_m radical release is the most crucial factor for the whole catalytic cycle. Because of the weaker coordination ability of OTf^- and NO_3^- than that of $\text{C}_n\text{X}_m\text{COO}^-$, easily dissociated iron salts like $\text{Fe}(\text{OTf})_3$ and $\text{Fe}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$ whose coordination anions prefer to be






Fig. 3 Chloro/fluoro-polyhaloalkylation of alkenes via iron photocatalysis. ^a 24 h reaction time. ^b 0.5 mmol Selectfluor was used. Isolated yields.

exchanged by excess $\text{C}_n\text{X}_m\text{COO}^-$, showed good catalytic performance. In contrast, the efficiency of $\text{Fe}(\text{acac})_3$ and $\text{Fe}(\text{OAc})_3 \cdot n\text{H}_2\text{O}$ was unsatisfactory, probably resulting from the stagnant ligand exchange between $\text{C}_n\text{X}_m\text{COO}^-$ and $\text{acac}^-/\text{OAc}^-$ (Fig. 4c). Moreover, the fact that only when $\text{Fe}(\text{OTf})_3$, **L2** and $\text{C}_n\text{X}_m\text{COO}^-$ are all present under the irradiation of blue light, can the iron(II) intermediate be observed, was revealed by UV-Vis experiments (Fig. 4d and S29†).

To quickly determine the possible structure of iron(III) and $\text{C}_n\text{X}_m\text{COO}^-$ -based light-harvesting species, DFT calculations were carried out (Tables S1 and S2†). As shown in Fig. 5a, the suitable coordination of **L2** and CH_3CN to $\text{Fe}(\text{CF}_3\text{COO})_3$ (**IntA**) to form **IntB** is thermodynamically favorable ($\Delta G = -20.8 \text{ kcal mol}^{-1}$), which also shows an obvious absorption band at 454 nm and an effective LMCT between CF_3COO^- and the iron(III) center in its excited state (Fig. S21–S24†). This result revealed






Fig. 4 Mechanistic studies. (a) Necessity of continuous blue light irradiation. (b) The first-order rate dependency on $\text{Fe}(\text{OTf})_3/\text{L2}$ of the reaction. (c) The comparison of various iron(III) salts. (d) UV-Vis studies on the iron photocatalysis.

that CH_3CN served as the second ligand to construct **IntB**. The ESI-HRMS experiment also identified the real existence of **IntB** under the standard conditions (Fig. S34†). Given that some alkenes were equipped with the ester scaffold whose coordination capability in comparison to $\text{C}_n\text{X}_m\text{COO}^-$ and nitriles should not be ignored, **IntC** as a possible LMCT-light-harvesting species could not be excluded. In contrast, the electron withdrawing inductive effect of the Cl group weakens the structural stability of **IntD**, potentially causing the low efficiency of producing C_nX_m radicals. Furthermore, we compared the performance of alkene 1 and 43 in different solvents such as CH_3CN , dichloromethane (DCM), acetone and CCl_3CN to prove the necessity of the second ligand for the activation of inert $\text{C}_n\text{X}_m\text{COO}^-$ (Fig. 5b). Due to the fact that there is no

coordination effect between the iron(III) catalyst and DCM/43, the chloro-trifluoromethylation of 43 under the condition of DCM as solvent was inefficient (entry 1). By contrast, replacing DCM with acetone obviously enhanced the yield of 16 (entry 2). The similar reactivities of 43 in CH_3CN and CCl_3CN solvent explained the possibility that **IntD** truly served as alternative iron and $\text{C}_n\text{X}_m\text{COO}^-$ -based light-harvesting species in the absence of acetonitrile (entries 3 and 4). We also studied the relationship between the final yields of 16 and the amounts of adsorbed CH_3CN (Fig. 5c). With the increase of the equivalent of CH_3CN , the desired reactivity for chloro-trifluoromethylation gradually improved, confirming that CH_3CN is the more efficient second ligand than CCl_3CN (Fig. S33†).




Fig. 5 Identification of iron and $C_nX_mCOO^-$ -based light-harvesting species. (a) DFT calculations on iron and $C_nX_mCOO^-$ -based light-harvesting species. (b) Experimental evidence for CH_3CN/CCl_3CN as the second ligand. (c) CH_3CN as the second ligand to promote the reaction.

Conclusion

In conclusion, we have developed a general and practical method in which dual ligand-enabled iron photocatalysis was leveraged for the activation of halogen-containing carboxylates under blue light conditions, thus achieving the chloro/fluoro-polyhaloalkylation of non-activated alkenes with broad functional group compatibility. Mechanistic studies revealed that the most likely real iron and $C_nX_mCOO^-$ -based light-harvesting species is assembled by the external dual ligand—OMe/ CF_3 -substituted bipyridine and acetonitrile/trichloroacetonitrile. Extending this strategy to the activation of more inert bulk chemicals is underway in our laboratory.

Data availability

The data supporting this article have been included as part of the ESI.†

Author contributions

Conceptualization: W. H., Z. Z., K. J., Y. L., D. W., S. L., and L. N.; methodology: W. H., Z. Z., K. J., Y. L., X. Y., D. W., S. L., and L. N.; investigation: W. H., Z. Z., K. J., Y. L., X. Y., D. W., S. L., and L. N.; supervision: Y. L., D. W., S. L., and L. N.; writing—original draft: W. H., Z. Z., K. J., Y. L., D. W., S. L., and L. N.; writing—review & editing: W. H., Z. Z., K. J., Y. L., X. Y., X. J., W. Y., Y. L., D. W., S. L., and L. N.

Conflicts of interest

The authors declare that they have no conflict of interest.

Acknowledgements

This project was supported by the National Natural Science Foundation of China (Grant No. 22101265, 21903071 and



21822303); Promotion Projects for Key Research & Development in Henan Province, China (Grant No. 222102310042); China Postdoctoral Science Foundation (Grant No. 2022M712866 and 2023M733212); Joint Fund of Key Technologies Research & Development Program of Henan Province (Grant No. 222301420006); The Ministry of Science and Technology of the People's Republic of China. The authors are also grateful for the support from Center of Advanced Analysis & Gene Sequencing of Zhengzhou University.

Notes and references

- (a) C. K. Prier, D. A. Rankic and D. W. MacMillan, *Chem. Rev.*, 2013, **113**, 5322–5363; (b) P. Bellotti, H.-M. Huang, T. Faber and F. Glorius, *Chem. Rev.*, 2023, **123**, 4237–4352; (c) J. F. Hartwig and M. A. Larsen, *ACS Cent. Sci.*, 2016, **2**, 281–292; (d) H. Jiang and A. Studer, *Chem. Soc. Rev.*, 2020, **49**, 1790–1811; (e) J. M. Narayanam and C. R. Stephenson, *Chem. Soc. Rev.*, 2011, **40**, 102–113; (f) D. Ravelli, S. Protti and M. Fagnoni, *Chem. Rev.*, 2016, **116**, 9850–9913; (g) N. Holmberg-Douglas and D. A. Nicewicz, *Chem. Rev.*, 2022, **122**, 1925–2016.
- (a) M. Chen and S. L. Buchwald, *Angew. Chem., Int. Ed.*, 2013, **52**, 11628–11631; (b) K. Matsui, E. Tobita, M. Ando and K. Kondo, *Chem. Lett.*, 1981, **10**, 1719–1720.
- (a) E. J. Cho, T. D. Senecal, T. Kinzel, Y. Zhang, D. A. Watson and S. L. Buchwald, *Science*, 2010, **328**, 1679–1681; (b) N. Rodríguez and L. J. Goossen, *Chem. Soc. Rev.*, 2011, **40**, 5030–5048; (c) J. D. Weaver, A. Recio, A. J. Grenning and J. A. Tunge, *Chem. Rev.*, 2011, **111**, 1846–1913; (d) Y. Wei, P. Hu, M. Zhang and W. Su, *Chem. Rev.*, 2017, **117**, 8864–8907; (e) X. Li, T. Yang, J. Li, X. Li, P. Chen, Z. Lin and G. Liu, *Nat. Chem.*, 2023, **15**, 862–871.
- (a) Y. Fujiwara, J. A. Dixon, F. O'Hara, E. D. Funder, D. D. Dixon, R. A. Rodriguez, R. D. Baxter, B. Herlé, N. Sach, M. R. Collins, Y. Ishihara and P. S. Baran, *Nature*, 2012, **492**, 95–99; (b) D. Yin, D. Su and J. Jin, *Cell Rep. Phys. Sci.*, 2020, **1**, 100141; (c) M. Tang, F. Draper, L. N. Pham, C. C. Ho, H. Huang, J. Sun, S. C. Thickett, M. L. Coote, T. U. Connell and A. C. Bissember, *J. Org. Chem.*, 2024, **89**, 2683–2690.
- (a) C. Lai and T. E. Mallouk, *J. Chem. Soc., Chem. Commun.*, 1993, 1359–1361; (b) W. Liu, X. Yang, Z.-Z. Zhou and C.-J. Li, *Chem*, 2017, **2**, 688–702; (c) T.-T. Yuan, J. Chen, L. N. Pham, S. Paul, L. V. White, J. Li, P. Lan, M. L. Coote, M. G. Banwell and Y.-T. He, *Org. Chem. Front.*, 2023, **10**, 4649–4657.
- (a) C. Depecker, H. Marzouk, S. Trevin and J. Devynck, *New J. Chem.*, 1999, **23**, 739–742; (b) H. Morimoto, T. Tsubogo, N. D. Litvinas and J. F. Hartwig, *Angew. Chem., Int. Ed.*, 2011, **50**, 3793–3798.
- (a) C.-J. Li, *Green Chem.*, 2016, **18**, 1836–1838; (b) P.-Z. Wang, W.-J. Xiao and J.-R. Chen, *Nat. Rev. Chem.*, 2023, **7**, 35–50; (c) J. Xuan, Z.-G. Zhang and W.-J. Xiao, *Angew. Chem., Int. Ed.*, 2015, **54**, 15632–15641; (d) S. Das, C. Zhu, D. Demirbas, E. Bill, C. De and B. List, *Science*, 2023, **379**, 494–499.
- (a) A. Guérinot and J. Cossy, *Top. Curr. Chem.*, 2016, **374**, 49; (b) A. Fürstner, *ACS Cent. Sci.*, 2016, **2**, 778–789.
- (a) Z. Huang, L. Jin, Y. Feng, P. Peng, H. Yi and A. Lei, *Angew. Chem., Int. Ed.*, 2013, **52**, 7151–7155; (b) M.-Q. Tian, Z.-Y. Shen, X. Zhao, P. J. Walsh and X.-H. Hu, *Angew. Chem., Int. Ed.*, 2021, **60**, 9706–9711.
- (a) W. Jian, L. Ge, Y. Jiao, B. Qian and H. Bao, *Angew. Chem., Int. Ed.*, 2017, **56**, 3650–3654; (b) F. Du, S.-J. Li, K. Jiang, R. Zeng, X.-C. Pan, Y. Lan, Y.-C. Chen and Y. Wei, *Angew. Chem., Int. Ed.*, 2020, **59**, 23755–23762.
- (a) T. Taniguchi, Y. Sugiura, H. Zaimoku and H. Ishibashi, *Angew. Chem., Int. Ed.*, 2010, **49**, 10154–10157; (b) J. Y. Hwang, A. Y. Ji, S. H. Lee and E. J. Kang, *Org. Lett.*, 2020, **22**, 16–21.
- (a) B. Liu, F. Jin, T. Wang, X. Yuan and W. Han, *Angew. Chem., Int. Ed.*, 2017, **56**, 12712–12717; (b) T. Tanaka, R. Yazaki and T. Ohshima, *J. Am. Chem. Soc.*, 2020, **142**, 4517–4524.
- (a) T. J. Collins and A. D. Ryabov, *Chem. Rev.*, 2017, **117**, 9140–9162; (b) X. Huang and J. T. Groves, *Chem. Rev.*, 2018, **118**, 2491–2553; (c) P. R. Montellano, *Chem. Rev.*, 2010, **110**, 932–948; (d) P. Xu, W. Fan, P. Chen and G. Liu, *J. Am. Chem. Soc.*, 2022, **144**, 13468–13474.
- L. H. Groot, A. Ilic, J. Schwarz and K. Wärnmark, *J. Am. Chem. Soc.*, 2023, **145**, 9369–9388.
- (a) Z. Li, X. Wang, S. Xia and J. Jin, *Org. Lett.*, 2019, **21**, 4259–4265; (b) G. Feng, X. Wang and J. Jin, *Eur. J. Org. Chem.*, 2019, 6728–6732; (c) S.-C. Kao, K.-J. Bian, X.-W. Chen, Y. Chen, A. A. Martí and J. G. West, *Chem. Catal.*, 2023, **3**, 100603; (d) L. M. Denkler, M. A. Shekar, T. S. Ngan, L. Wylie, D. Abdullin, T. Hett, F. H. Pilz, B. Kirchner, O. Schiemann, P. Kielb, A. Bunesco, M. Engeser and G. Schnakenburg, *Angew. Chem., Int. Ed.*, 2024, e202403292.
- (a) F. Juliá, *ChemCatChem*, 2022, **14**, e202200916; (b) Q. Zhang, S. Liu, J. Lei, Y. Zhang, C. Meng, C. Duan and Y. Jin, *Org. Lett.*, 2022, **24**, 1901–1906; (c) B. Niu, K. Sachidanandan, M. V. Cooke, T. E. Casey and S. Laulhé, *Org. Lett.*, 2022, **24**, 4524–4529; (d) S. Oh and E. E. Stache, *J. Am. Chem. Soc.*, 2022, **144**, 5745–5749; (e) D. Gygi, M. I. Gonzalez, S. J. Hwang, K. T. Xia, Y. Qin, E. J. Johnson, F. Gygi, Y.-S. Chen and D. G. Nocera, *J. Am. Chem. Soc.*, 2021, **143**, 6060–6064; (f) M. I. Gonzalez, D. Gygi, Y. Qin, Q. Zhu, E. J. Johnson, Y. S. Chen and D. G. Nocera, *J. Am. Chem. Soc.*, 2022, **144**, 1464–1472; (g) Y. Jin, Q. Zhang, L. Wang, X. Wang, C. Meng and C. Duan, *Green Chem.*, 2021, **23**, 6984–6989; (h) Z.-Y. Dai, S.-Q. Zhang, X. Hong, P.-S. Wang and L.-Z. Gong, *Chem. Catal.*, 2022, **2**, 1211–1222; (i) Y. Jin, L. Wang, Q. Zhang, Y. Zhang, Q. Liao and C. Duan, *Green Chem.*, 2021, **23**, 9406–9411; (j) G. Zhang, Z. Zhang and R. Zeng, *Chin. J. Chem.*, 2021, **39**, 3225–3230; (k) Y. C. Kang, S. M. Treacy and T. Rovis, *ACS Catal.*, 2021, **11**, 7442–7449.
- (a) W. Liu, Q. Wu, M. Wang, Y. Huang and P. Hu, *Org. Lett.*, 2021, **23**, 8413–8418; (b) N. Xiong, Y. Li and R. Zeng, *Org. Lett.*, 2021, **23**, 8968–8972; (c) T. Xue, Z. Zhang and R. Zeng, *Org. Lett.*, 2022, **24**, 977–982.



- 18 (a) M. Zhang, J. Zhang, Q. Li and Y. Shi, *Nat. Commun.*, 2022, **13**, 7880; (b) K.-J. Bian, S.-C. Kao, D. Nemoto Jr, X.-W. Chen and J. G. West, *Nat. Commun.*, 2022, **13**, 7881; (c) H. Lindner, W. M. Amberg and E. M. Carreira, *J. Am. Chem. Soc.*, 2023, **145**, 22347–22353; (d) W. Zhang, T. Liu, H. T. Ang, P. Luo, Z. Lei, X. Luo, M. J. Koh and J. Wu, *Angew. Chem., Int. Ed.*, 2023, **62**, e202310978.
- 19 (a) K.-J. Bian, Y.-C. Lu, D. Nemoto Jr, S.-C. Kao, X. Chen and J. G. West, *Nat. Chem.*, 2023, **15**, 1683–1692; (b) J. Qi, J. Xu, H. T. Ang, B. Wang, N. K. Gupta, S. R. Dubbaka, P. O'Neill, X. Mao, Y. Lum and J. Wu, *J. Am. Chem. Soc.*, 2023, **145**, 24965–24971; (c) X.-K. Qi, L.-J. Yao, M.-J. Zheng, L. Zhao, C. Yang, L. Guo and W. Xia, *ACS Catal.*, 2024, **14**, 1300–1310.
- 20 X.-Y. Jiang, Y. Lan, Y. Hao, K. Jiang, J. He, J. Zhu, S. Jia, J. Song, S.-J. Li and L. Niu, *Nat. Commun.*, 2024, **15**, 6115.
- 21 During the preparation of this manuscript, Juliá-Hernández *et al.* published the photo-induced iron catalyzed decarboxylation of trifluoroacetates: S. Fernández-García, V. O. Chantzakou and F. Juliá-Hernández, *Angew. Chem., Int. Ed.*, 2023, **63**, e202311984.
- 22 A. J. Fernandes, R. Giri, K. N. Houk and D. Katayev, *Angew. Chem., Int. Ed.*, 2024, e202318377.
- 23 (a) S. Kamijo, S. Yokosaka and M. Inoue, *Tetrahedron*, 2012, **68**, 5290–5296; (b) X.-J. Tang and W. R. Dolbier Jr, *Angew. Chem., Int. Ed.*, 2015, **54**, 4246–4249; (c) D. B. Bagal, G. Kachkovskiy, M. Knorn, T. Rawner, B. M. Bhanage and O. Reiser, *Angew. Chem., Int. Ed.*, 2015, **54**, 6999–7002; (d) J. Fang, Z.-K. Wang, S.-W. Wu, W.-G. Shen, G.-Z. Ao and F. Liu, *Chem. Commun.*, 2017, **53**, 7638–7641; (e) K.-Y. Ye, G. Pombar, N. Fu, G. Sauer, I. Keresztes and S. Lin, *J. Am. Chem. Soc.*, 2018, **140**, 2438–2441; (f) M. Alkan-Zambada and X. Hu, *Organometallics*, 2018, **37**, 3928–3935; (g) X. Sun, H.-X. Ma, T.-S. Mei, P. Fang and Y. Hu, *Org. Lett.*, 2019, **21**, 3167–3171; (h) H. Sun, G. Cui, H. Shang and B. Cui, *J. Org. Chem.*, 2020, **85**, 15241–15255; (i) P. Lian, W. Long, J. Li, Y. Zheng and X. Wan, *Angew. Chem., Int. Ed.*, 2020, **59**, 23603–23608; (j) V. S. Kostromitin, A. A. Zemtsov, V. V. Levin and A. D. Dilman, *Adv. Synth. Catal.*, 2021, **363**, 5336–5340; (k) P. Basnet, G. Hong, C. E. Hendrick and M. C. Kozlowski, *Org. Lett.*, 2021, **23**, 433–437; (l) X.-H. Li, G.-C. Yang, J.-F. Gong and M.-P. Song, *Org. Biomol. Chem.*, 2022, **20**, 4815–4825; (m) D. Wu, W. Fan, L. Wu, P. Chen and G. Liu, *ACS Catal.*, 2022, **12**, 5284–5291; (n) C. Wu, X. Hui, D. Zhang, M. Zhang, Y. Zhu and S. Wang, *Green Chem.*, 2022, **24**, 1103–1108; (o) M. Zhang, J. Zhang and M. Oestreich, *Nat. Synth.*, 2023, **2**, 439–447; (p) X.-J. Ren, P.-W. Liao, H. Sheng, Z.-X. Wang and X.-Y. Chen, *Org. Lett.*, 2023, **25**, 6189–6194; (q) Y. Xie, Z. Zhang, B. Zhang, N. He, M. Peng, S. Song, B. Wang and F. Yu, *J. Org. Chem.*, 2024, **89**, 8521–8530; (r) J. Elfert, N.-L. Frye, I. Rempel, C. G. Daniliuc and A. Studer, *Nat. Commun.*, 2024, **15**, 7230; (s) K.-J. Bian, D. Nemoto Jr, X.-W. Chen, S.-C. Kao, J. Hoosona and J. G. West, *Chem. Sci.*, 2024, **15**, 124–133.
- 24 (a) S. Kindt and M. R. Heninrich, *Chem.-Eur. J.*, 2014, **20**, 15344–15348; (b) W. Yu, X.-H. Xu and F.-L. Qing, *Adv. Synth. Catal.*, 2015, **357**, 2039–2044; (c) Z. Liu, H. Chen, Y. Lv, X. Tan, H. Shen, H.-Z. Yu and C. Li, *J. Am. Chem. Soc.*, 2018, **140**, 6169–6175; (d) A. S. Porzer, E.-M. Alvarez, H. Friedrich and M. R. Heinrich, *Chem.-Eur. J.*, 2019, **25**, 2786–2792; (e) Q. Wang, Y. Qu, H. Tian, Y. Liu, H. Song and Q. Wang, *Chem.-Eur. J.*, 2019, **25**, 8686–8690; (f) J.-L. Li, Y.-Q. Liu, W.-L. Zou, R. Zeng, X. Zhang, Y. Liu, B. Han, Y. He, H.-J. Leng and Q.-Z. Li, *Angew. Chem., Int. Ed.*, 2020, **59**, 1863–1870; (g) S. Ma, F. Li, G. Zhang, L. Shi and X. Wang, *ACS Catal.*, 2021, **11**, 14848–14853; (h) A. Granados, R. K. Dhungana, M. Sharique, J. Majhi and G. A. Molander, *Org. Lett.*, 2022, **24**, 4750–4755; (i) V. S. Kostromitin, A. O. Sorokin, V. V. Levin and A. D. Dilman, *Org. Lett.*, 2023, **25**, 6598–6602; (j) Y. Li, Y. Dong, X. Wang, G. Li, H. Xue, W. Xin, Q. Zhang, W. Guan and J. Fu, *ACS Catal.*, 2023, **13**, 2410–2421; (k) V. S. Kostromitin, V. V. Levin and A. D. Dilman, *J. Org. Chem.*, 2023, **88**, 6252–6262; (l) R.-X. Liu, W. Wang and L. Yang, *Adv. Synth. Catal.*, 2024, **366**, 1643–1648.

