

inal alkenes.¹⁷ Moreover, the (P^CNN)Co catalyst effects the hydrosilylation of internal alkynes, and intriguingly, the reactions of arylalkyl disubstituted alkynes produce vinylsilanes containing the silyl unit at the alkyl-substituted olefinic carbon, rather than the aryl-substituted one. The (Z)-β-vinylsilane products can undergo Pd-catalyzed Hiyama-Denmark cross-couplings to form (Z)-disubstituted alkenes.

Results and discussion

We commenced our studies by examining the P^CNN ligand-supported Fe and Co catalysts for the hydrosilylation of a simple aliphatic alkyne, 1-hexyne (3a), with Ph₂SiH₂ as the silicon source. The results are summarized in Table 1. Using NaHBET₃H (2 mol%) as the activator, both *t*Bu- and *i*Pr-substituted Fe complexes, ((^{*t*Bu}P^CNN^{*i*Pr}))FeCl₂ (**1a**, 1 mol%) and ((^{*i*Pr}P^CNN^{*i*Pr}))FeCl₂ (**1b**, 1 mol%), were inactive for the hydrosilylation (entries 1 and 2). However, 1-hexyne (1.2 equiv.) reacted with Ph₂SiH₂ to give (Z)-β-vinylsilane **4a** and (E)-β-vinylsilane **5a** when the Co analogues were employed. Using the most sterically hindered complex **2a** (1 mol%) bearing a *t*Bu-substituted phosphino group and *i*Pr substituents at the 2,6-positions of the *N*-aryl ring, the hydrosilylation in THF at 25 °C gave 13% **4a** and 18% **5a** after 24 h (entry 3). Reducing the steric hindrance of the P^CNN ligands resulted in a significant improvement of the activity and selectivity for the desired

(Z)-β-product. The reactions using complexes ((^{*i*Pr}P^CNN^{*i*Pr}))CoCl₂ **2b** and ((^{*i*Pr}P^CNN^{Me}))CoCl₂ **2c** containing *i*Pr-substituted phosphino groups formed **4a** in 85% and 83% yield; α-vinylsilane was not observed in both cases (entries 4 and 5).

Further optimizations were conducted to improve the selectivity for *cis*-vinylsilane. The best Z-selectivity was obtained when the reaction occurred under solvent-free conditions using a 1.5 : 1 ratio of **3a** : Ph₂SiH₂ in the presence of 1 mol% of **2b** and 2 mol% of NaHBET₃; the run at 25 °C for 24 h produced (Z)-vinylsilane **4a** in 95% yield with the Z/E stereoselectivity of 95 : 1 (entry 6). A control experiment without the precatalyst **2b**, but with NaHBET₃H, gave no hydrosilylation product (entry 7), and another control without NaHBET₃ indicated that the addition of the activator was necessary for the catalysis (entry 8). Finally, we explored the catalytic performance of this Co catalyst for hydrosilylations with primary and tertiary hydrosilanes. Tertiary silanes (EtO)₃SiH, MD'M (1,1,1,3,5,5,5-heptamethyltrisiloxane), Et₃SiH and PhMe₂SiH were unreactive under the reaction conditions (entries 9–12). Although the reaction with PhSiH₃ gave the *anti*-Markovnikov products in high yield (95% totally), the stereoselectivity was low (Z/E = ~1 : 1) (entry 13).

To evaluate the scope of the Co-catalyzed alkyne hydrosilylation, the reactions with a variety of functionalized terminal alkynes and a number of internal alkynes were carried out using **2b** as the precatalyst. The results are summarized in Table 2. All reactions with Ph₂SiH₂ proceeded to completion within 24 h at room temperature under solvent-free conditions. For terminal alkyne substrates, most reactions were highly selective for the formation of the (Z)-β-vinylsilane products (Z/E > 95 : 5). Linear terminal alkynes gave the corresponding products **4a–d** in moderate-to-high isolated yield with excellent selectivity. 3-Cyclohexyl-1-propyne **3e** and 3-cyclopentyl-1-propyne **3f** underwent hydrosilylations in useful yields and high selectivities. The catalyst is compatible with terminal alkynes bearing various functional groups, including chloride (**4g**), nitrile (**4h**), tertiary amine (**4i**, **4j**), amide (**4k**) and even ester (**4o**). Aliphatic alkynes containing protecting groups, such as silyl ether (**4l**, **4m**) and tosylate (**4n**, 88%), afforded the desired (Z)-β-vinylsilanes in high yields and high selectivities. Aryl terminal alkynes are unreactive under the catalytic conditions, as demonstrated by the full recovery of the alkyne in the reaction of phenylacetylene with Ph₂SiH₂.

Internal alkynes underwent the hydrosilylation reactions under the catalytic conditions to form the products in good-to-high yields. The hydrosilylation of symmetric dialkyl-substituted alkynes (**3p**, **3q**) occurred to form the *syn*-addition products in high yields (**4p**, **4q**). Next, we turned our attention to the more challenging unsymmetrical internal alkynes. The unsymmetrical dialkyl-substituted alkynes with different steric properties, such as 1-*tert*-butyl-1-propyne (**3r**) and 4-methylpent-2-yne (**3s**), were suitable for the selective hydrosilylation, producing the *syn*-addition products **4r** and **4s** with the incorporation of the silyl group into the less sterically demanding sp-hybridised carbon. The reaction of arylalkyl disubstituted alkynes (**3t–v**) gave the products in 91–94% yields (**4t–v**).

Table 1 (P^CNN)Fe and (P^CNN)Co complex-catalyzed hydrosilylation of 1-hexyne with hydrosilanes^a

Entry	Cat.	Silane	Solvent	3a/[Si]	Yield [%]	
					4a	5a
1	1a	Ph ₂ SiH ₂	THF	1.2	0	0
2	1b	Ph ₂ SiH ₂	THF	1.2	0	0
3	2a	Ph ₂ SiH ₂	THF	1.2	13	18
4	2b	Ph ₂ SiH ₂	THF	1.2	85	3
5	2c	Ph ₂ SiH ₂	THF	1.2	83	3
6	2b	Ph ₂ SiH ₂	None	1.5	95	1
7 ^b	2b	Ph ₂ SiH ₂	None	1.5	0	0
8 ^c	2b	Ph ₂ SiH ₂	None	1.5	0	0
9	2b	(EtO) ₃ SiH	None	1.5	<5	<5
10	2b	MD'M	None	1.5	0	0
11	2b	Et ₃ SiH	None	1.5	0	0
12	2b	PhMe ₂ SiH	None	1.5	0	0
13	2b	PhSiH ₃	None	1.5	50	45

^a Reaction conditions: **3a** (0.72–0.9 mmol) and silane (0.6 mol) in THF (1 mL) or under neat conditions at 25 °C. The yields were determined by ¹H NMR spectroscopy. ^b Without precat. ^c Without NaHBET₃.



Table 2 Cobalt-catalyzed hydrosilylation of various terminal and internal alkynes with Ph_2SiH_2 ^a

$\begin{array}{c} \text{R}-\text{C}\equiv\text{C}-\text{H} \\ \text{or} \\ \text{R}-\text{C}\equiv\text{C}-\text{R}' \end{array}$		$\xrightarrow[\text{neat, rt, 24 h}]{\begin{array}{c} 1 \text{ mol\% } \mathbf{2b} \\ 2 \text{ mol\% } \text{NaBEt}_3\text{H} \\ 100 \text{ mol\% } \text{Ph}_2\text{SiH}_2 \end{array}}$	$\begin{array}{c} \text{R}-\text{CH}=\text{CH}-\text{SiHPh}_2 \\ \text{or} \\ \text{R}-\text{CH}=\text{CH}-\text{SiHPh}_2 \end{array}$
3a-w			4a-w
	4a , 88%, 99:1		4b , 88%, 98:2
	4c , 91%, 99:1		4d , 61%, 93:7
	4e , 83%, 91:9		4f , 58%, 90:10
	4g , 83%, 98:2		4h , 68%, 84:16 ^b
	4i , 92%, 98:2		4j , 88%, 98:2
	4k , 85%, 97:3		4l , 90%, 98:2
	4m , 93%, 98:2		4n , 88%, 99:1
	4o , 80%, 98:2 ^c		4p , 84%, 99:1
	4q , 71%, 99:1		4r , 82%, 99:1 ^d
	4s , 80%, 90:10 ^d		4t , 94%, 91:9 ^d
	4u , 93%, 88:12 ^d		4v , 91%, 91:9 ^d

^a Reaction conditions: Ph_2SiH_2 (0.6 mmol) and alkyne (0.9 mmol) at 25 °C. Yield of isolated products. *Z/E* ratios of isolated products were determined by ^1H NMR spectroscopy. ^b With 1.2 mmol **3h**. A 84:16 *Z/E* ratio was determined by ^1H NMR spectroscopy. ^c With 2 mol% **2b** and 4 mol% NaBEt_3H . ^d Regioselectivities determined by ^1H NMR spectroscopy.

Noteworthy, the processes exhibited good regioselectivities, furnishing tri-substituted olefins with the Ph_2HSi unit selectively added to the alkyl-substituted olefinic carbon, which is in contrast to previous reports on Co-catalyzed hydrosilylations of arylalkyl disubstituted alkynes that yielded vinylsilanes with the Si atom adjacent to the aryl group.^{9e,11,13}

Due to their high stability, non-toxicity, and ease of handling, vinylsilanes have become valuable intermediates for Pd-catalyzed Hiyama–Denmark cross-couplings.¹⁸ However, the silane reagents used in the coupling reactions often require heteroatom substituent(s) (e.g., O or F) to make the Si atom feasible toward activation with a base or fluoride source.¹⁹ To this end, it is of interest to develop conditions for cross-couplings of our (*Z*)-vinylsilane products bearing the Ph_2HSi moiety.²⁰


After a screening of various mono- and bisphosphine ligands (see the ESI†), we identified the Pd catalyst generated from $\text{Pd}(\text{OAc})_2$ (10 mol%) and 1,4-bis(diphenylphosphino)butane (dppb, 12 mol%) was effective for stereoselective couplings of (*Z*)-vinylsilane (**4**) with aryl iodides using TBAF as the activator (2 equiv.). Most reactions occurred at 30 °C to form the *cis*-1,2-disubstituted alkenes in high yields and high stereoselectivities (Table 3). The electrophiles **6** bearing both electron-donating and -withdrawing groups were suitable coupling partners. Functionalized groups, such as ester (**7a**), ketone (**7b**), tertiary amine (**7d**) and nitrile (**7j**) could be tolerated. The reaction of **6c** containing a nitro group provided the coupling product **7c** in high yield, albeit with relatively low *Z*-selectivity (75:25). Aryl iodide **6i** with a *meta*-MeO group gave the product with somewhat decreased selectivity (**7i**, 88:12) compared to the reaction with 4-iodoanisole (**7h**, 97:3). Aryl bromide is suitable for the coupling reaction as demonstrated by the formation of the desired product **7h** in useful yield (62%) with 4-bromoanisole as the reagent. Heteroaromatic iodide, such as 2-iodothiophene and 2-acetyl-5-iodothiophene,

Table 3 Palladium-catalyzed Hiyama–Denmark cross-couplings of (*Z*)-vinylsilane with various aryl halides^a

$\begin{array}{c} \text{R}-\text{CH}=\text{CH}-\text{SiHPh}_2 \\ \mathbf{4} \end{array}$	$\begin{array}{c} \text{R}'-\text{C}_6\text{H}_4-\text{I} \\ \mathbf{6} \end{array}$	$\xrightarrow[\text{THF, 30 }^\circ\text{C, 12 h}]{\begin{array}{c} \text{Pd}(\text{OAc})_2 (10 \text{ mol\%}) \\ \text{dppb} (12 \text{ mol\%}) \\ \text{TBAF} (200 \text{ mol\%}) \end{array}}$	$\begin{array}{c} \text{R}'-\text{C}_6\text{H}_4-\text{CH}=\text{CH}-\text{SiHPh}_2 \\ \mathbf{7a-n} \end{array}$
	7a , 89%, 96:4		7b , 77%, 98:2
	7c , 86%, 75:25 ^b		7d , 62%, 94:6
	7e , 91%, 94:6		7f , 78%, 97:3
	7g , 83%, 93:7		7h , 84%, 97:3 62%, 92:8 ^c
	7i , 70%, 88:12		7j , 55%, 91:9
	7k , 66%, 90:10		7l , 64%, 79:21
	7m , 96%, 94:6		7n , 94%, 98:2

^a Reaction conditions: **4c** (0.24 mmol), aryl iodide (0.2 mmol), TBAF (0.4 mmol) in THF (0.4 mL). Yield of isolated products, unless otherwise noted. *Z/E* ratios of isolated products were determined by GC. ^b Without dppb, 24 h. ^c Use 4-bromoanisole. Yield determined by ^1H NMR spectroscopy.





Scheme 2 Proposed mechanism involving the Ojima–Crabtree type rearrangement for Co-catalyzed (Z)-selective hydrosilylation of terminal alkynes.

gave the corresponding products **7k** and **7l** in 66 and 64% isolated yields, respectively. Functional Z-vinylsilanes such as **4g** and **4o** reacted with 4-iodo-1,1'-biphenyl under standard conditions and gave the corresponding Z-alkenes **7m** and **7n** in excellent yields and selectivities.

A mechanism accounting for the formation of (Z)-β-vinylsilane is presented in Scheme 2. On the basis of the precedents of Co-catalyzed hydrosilylation processes,^{11,16,17,21} we propose that the catalytic cycle involves a low-valent Co(I) silyl intermediate. The insertion of the coordinated terminal alkyne into the Co–Si bond forms a vinyl Co intermediate, (Z)-l-silyl-l-alken-2-yl-cobalt (**C**). Due to the significant steric repulsion between the bulky silyl group and the metal moiety, complex **C** undergoes isomerization to form a sterically less demanding complex **F**, (E)-l-silyl-l-alken-2-yl-cobalt, through the Ojima–Crabtree type rearrangement.^{4h,5g} **E** then reacts with Ph₂SiH₂ to form (Z)-vinylsilane and regenerate the Co(I) silyl species **A**.

Conclusions


In conclusion, we have developed a highly efficient cobalt-catalyzed regio- and stereoselective hydrosilylation of terminal alkynes with Ph₂SiH₂ to generate (Z)-β-vinylsilanes with high functional group tolerance. The silyl groups introduced into the vinylsilane products offer flexible synthetic manipulations as demonstrated by the Hiyama–Denmark cross-couplings with aryl halides, allowing a convenient, stereoselective synthesis of *Cis*-disubstituted alkenes.

Acknowledgements

We gratefully acknowledge the financial support from the National Natural Science Foundation of China (No. 21272255, 21121062).

Notes and references

- (a) H. Brunner, *Angew. Chem., Int. Ed.*, 2004, **43**, 2749; (b) B. M. Trost and Z. T. Ball, *Synthesis*, 2005, 853; (c) S. J. Clarson, *Silicon*, 2009, **1**, 57; (d) D. S. W. Lim and E. A. Anderson, *Synthesis*, 2012, 983.
- (a) T. H. Chan and I. Fleming, *Synthesis*, 1979, 761; (b) T. A. Blumenkopf and L. E. Overman, *Chem. Rev.*, 1986, **86**, 857; (c) E. Langkopf and D. Schinzer, *Chem. Rev.*, 1995, **95**, 1375; (d) I. Fleming, A. Barbero and D. Walter, *Chem. Rev.*, 1997, **97**, 2063.
- (a) H. Aneetha, W. Wu and J. G. Verkade, *Organometallics*, 2005, **24**, 2590; (b) G. De Bo, G. Berthon-Gelloz, B. Tinant and I. E. Markó, *Organometallics*, 2006, **25**, 1881; (c) Y. Nakao, H. Imanaka, J. Chen, A. Yada and T. Hiyama, *J. Organomet. Chem.*, 2007, **692**, 585; (d) G. Berthon-Gelloz, J.-M. Schumers, G. De Bo and I. E. Markó, *J. Org. Chem.*, 2008, **73**, 4190; (e) A. Hamze, O. Provot, J.-D. Brion and M. Alami, *J. Organomet. Chem.*, 2008, **693**, 2789; (f) M. Blug, X.-F. Le Goff, N. Mézailles and P. Le Floch, *Organometallics*, 2009, **28**, 2360; (g) R. Cano, M. Yus and D. J. Ramón, *ACS Catal.*, 2012, **2**, 1070; (h) K. Igawa, D. Yoshihiro, N. Ichikawa, N. Kokan and K. Tomooka, *Angew. Chem., Int. Ed.*, 2012, **51**, 12745; (i) S. Dierick, E. Vercruysse, G. Berthon-Gelloz and I. E. Markó, *Chem. – Eur. J.*, 2015, **21**, 17073; (j) D. Munz, C. Allolio, D. Meyer, M. Micksch, L. Roessner and T. Strassner, *J. Organomet. Chem.*, 2015, **794**, 330; (k) L. Ortega-Moreno, R. Peloso, C. Maya, A. Suarez and E. Carmona, *Chem. Commun.*, 2015, **51**, 17008.
- (a) M. Atsunori, T. Eisuke, K. Hiroshi, H. Kazunori, N. Yasushi and H. Tamejiro, *Chem. Lett.*, 1998, **27**, 443; (b) J. W. Faller and D. G. D'Alliessi, *Organometallics*, 2002, **21**, 1743; (c) A. Mori, E. Takahisa, Y. Yamamura, T. Kato, A. P. Mudalige, H. Kajiro, K. Hirabayashi, Y. Nishihara and T. Hiyama, *Organometallics*, 2004, **23**, 1755; (d) M. V. Jiménez, J. J. Pérez-Torrente, M. I. Bartolomé, V. Gierz, F. J. Lahoz and L. A. Oro, *Organometallics*, 2008, **27**, 224; (e) J. L. McBee, J. Escalada and T. D. Tilley, *J. Am. Chem. Soc.*, 2009, **131**, 12703; (f) M. Iglesias, M. Perez-Nicolas, P. J. S. Miguel, V. Polo, F. J. Fernandez-Alvarez, J. J. Perez-Torrente and L. A. Oro, *Chem. Commun.*, 2012, **48**, 9480; (g) M. Iglesias, P. J. Sanz Miguel, V. Polo, F. J. Fernández-Alvarez, J. J. Pérez-Torrente and L. A. Oro, *Chem. – Eur. J.*, 2013, **19**, 17559; (h) I. Ojima, N. Clos, R. J. Donovan and P. Ingallina, *Organometallics*, 1990, **9**, 3127.
- (a) C.-H. Jun and R. H. Crabtree, *J. Organomet. Chem.*, 1993, **447**, 177; (b) M. Viciano, E. Mas-Marzá, M. Sanaú and E. Peris, *Organometallics*, 2006, **25**, 3063; (c) M. Yoshihiro, I. Eigo and I. Masahiko, *Chem. Lett.*, 2006, **35**, 836; (d) V. S. Sridevi, W. Y. Fan and W. K. Leong, *Organometallics*, 2007, **26**, 1157; (e) A. Zanardi, E. Peris and J. A. Mata, *New J. Chem.*, 2008, **32**, 120; (f) J. J. Pérez-Torrente, D. H. Nguyen, M. V. Jiménez, F. J. Modrego, R. Puerta-Oteo, D. Gómez-Bautista, M. Iglesias and



- L. A. Oro, *Organometallics*, 2016, **35**, 2410; (g) R. S. Tanke and R. H. Crabtree, *J. Am. Chem. Soc.*, 1990, **112**, 7984.
- 6 (a) M. A. Esteruelas, J. Herrero and L. A. Oro, *Organometallics*, 1993, **12**, 2377; (b) H. Katayama, K. Taniguchi, M. Kobayashi, T. Sagawa, T. Minami and F. Ozawa, *J. Organomet. Chem.*, 2002, **645**, 192; (c) C. S. Arico and L. R. Cox, *Org. Biomol. Chem.*, 2004, **2**, 2558; (d) S. V. Maifeld, M. N. Tran and D. Lee, *Tetrahedron Lett.*, 2005, **46**, 105; (e) C. Menozzi, P. I. Dalko and J. Cossy, *J. Org. Chem.*, 2005, **70**, 10717; (f) M. Nagao, K. Asano, K. Umeda, H. Katayama and F. Ozawa, *J. Org. Chem.*, 2005, **70**, 10511; (g) R. Gao, D. R. Pahls, T. R. Cundari and C. S. Yi, *Organometallics*, 2014, **33**, 6937; (h) C. Conifer, C. Gunanathan, T. Rinesch, M. Hölscher and W. Leitner, *Eur. J. Inorg. Chem.*, 2015, 333; (i) Y. Na and S. Chang, *Org. Lett.*, 2000, **2**, 1887.
- 7 (a) K. Junge, K. Schroder and M. Beller, *Chem. Commun.*, 2011, **47**, 4849; (b) B. D. Sherry and A. Fürstner, *Acc. Chem. Res.*, 2008, **41**, 1500; (c) I. Bauer and H.-J. Knölker, *Chem. Rev.*, 2015, **115**, 3170; (d) M. D. Greenhalgh, A. S. Jones and S. P. Thomas, *ChemCatChem*, 2015, **7**, 190; (e) J. Sun and L. Deng, *ACS Catal.*, 2016, **6**, 290; (f) X. Du and Z. Huang, *ACS Catal.*, 2017, 1227.
- 8 (a) S. C. Bart, E. Lobkovsky and P. J. Chirik, *J. Am. Chem. Soc.*, 2004, **126**, 13794; (b) C. Belger and B. Plietker, *Chem. Commun.*, 2012, **48**, 5419; (c) M. D. Greenhalgh, D. J. Frank and S. P. Thomas, *Adv. Synth. Catal.*, 2014, **356**, 584; (d) A. J. Challinor, M. Calin, G. S. Nichol, N. B. Carter and S. P. Thomas, *Adv. Synth. Catal.*, 2016, **358**, 2404.
- 9 (a) M. Isobe, R. Nishizawa, T. Nishikawa and K. Yoza, *Tetrahedron Lett.*, 1999, **40**, 6927; (b) L. Yong, K. Kirleis and H. Butenschön, *Adv. Synth. Catal.*, 2006, **348**, 833; (c) T. Konno, K.-i. Taku, S. Yamada, K. Moriyasu and T. Ishihara, *Org. Biomol. Chem.*, 2009, **7**, 1167; (d) K.-H. Huang and M. Isobe, *Eur. J. Org. Chem.*, 2014, 4733; (e) A. Rivera-Hernández, B. J. Fallon, S. Ventre, C. Simon, M.-H. Tremblay, G. Gontard, E. Derat, M. Amatore, C. Aubert and M. Petit, *Org. Lett.*, 2016, **18**, 4242.
- 10 (a) T. Bartik, G. Nagy, P. Kvintovics and B. Happ, *J. Organomet. Chem.*, 1993, **453**, 29; (b) A. Tillack, S. Pulst, W. Baumann, H. Baudisch, K. Kortus and U. Rosenthal, *J. Organomet. Chem.*, 1997, **532**, 117; (c) M. R. Chaulagain, G. M. Mahandru and J. Montgomery, *Tetrahedron*, 2006, **62**, 7560; (d) J. Berding, J. A. van Paridon, V. H. S. van Rixel and E. Bouwman, *Eur. J. Inorg. Chem.*, 2011, 2450.
- 11 Z. Mo, J. Xiao, Y. Gao and L. Deng, *J. Am. Chem. Soc.*, 2014, **136**, 17414.
- 12 J. H. Docherty, J. Peng, A. P. Dominey and S. P. Thomas, *Nat. Chem.*, 2017, DOI: 10.1038/nchem.2697.
- 13 (a) J. Guo and Z. Lu, *Angew. Chem., Int. Ed.*, 2016, **55**, 10835; (b) J. Guo, X. Shen and Z. Lu, *Angew. Chem., Int. Ed.*, 2017, **56**, 615.
- 14 Z. Zuo, J. Yang and Z. Huang, *Angew. Chem., Int. Ed.*, 2016, **55**, 10839.
- 15 Note that (Z)- β -selective hydroboration of terminal alkynes has been previously reported by Chirik *et al.* See: J. V. Obligacion, J. M. Neely, A. N. Yazdani, I. Pappas and P. J. Chirik, *J. Am. Chem. Soc.*, 2015, **137**, 5855.
- 16 W. J. Teo, C. Wang, Y. W. Tan and S. Ge, *Angew. Chem., Int. Ed.*, 2017, **56**, 4328.
- 17 X. Du, Y. Zhang, D. Peng and Z. Huang, *Angew. Chem., Int. Ed.*, 2016, **55**, 6671.
- 18 (a) S. E. Denmark and J. H. C. Liu, *Angew. Chem., Int. Ed.*, 2010, **49**, 2978; (b) Y. Nakao and T. Hiyama, *Chem. Soc. Rev.*, 2011, **40**, 4893; (c) H. F. Sore, W. R. J. D. Galloway and D. R. Spring, *Chem. Soc. Rev.*, 2012, **41**, 1845; (d) S. E. Denmark and A. Ambrosi, *Org. Process Res. Dev.*, 2015, **19**, 982; (e) T. Komiyama, Y. Minami and T. Hiyama, *ACS Catal.*, 2017, **7**, 631.
- 19 C. Cheng, E. M. Simmons and J. F. Hartwig, *Angew. Chem., Int. Ed.*, 2013, **52**, 8984.
- 20 Lu reported cross-couplings of two diphenyl-substituted α -vinylsilanes with aryl iodide using $\text{Pd}_2(\text{dba})_3$ as the catalyst. See ref. 13.
- 21 C. C. H. Atienza, T. Diao, K. J. Weller, S. A. Nye, K. M. Lewis, J. G. P. Delis, J. L. Boyer, A. K. Roy and P. J. Chirik, *J. Am. Chem. Soc.*, 2014, **136**, 12108.

