

Silver-catalysed trifluoromethylation of arenes at room temperature†

Sangwon Seo,^a John B. Taylor^b and Michael F. Greaney^{*a}Cite this: *Chem. Commun.*, 2013, **49**, 6385Received 11th March 2013,
Accepted 16th May 2013

DOI: 10.1039/c3cc41829d

www.rsc.org/chemcomm

A variety of heteroarenes and electron rich arenes can be trifluoromethylated at room temperature with TMSCF_3 , catalytic silver and PhI(OAc)_2 .

The trifluoromethyl group is valued for its ability to modulate the properties of diverse materials such as pharmaceuticals, agrochemicals and polymers. Aryl CF_3 groups are electron-withdrawing, hydrophobic and generally very stable, all properties that can be harnessed in the design of biologically active molecules and functional materials. Synthetic methods for aryl and heteroaryl trifluoromethylation are thus critical to the discovery and production of new molecules of high value to society.¹ Recent developments in metal-mediated trifluoromethylation have produced significant advances in this area,² with common functional groups such as aryl boronic acids and halides undergoing efficient trifluoromethylation under palladium and copper catalysis.³ Metal-catalysed trifluoromethylation of unactivated C–H positions, by contrast, is significantly less developed and has great potential for accelerating medicinal and agrochemical syntheses. Despite some recent ground-breaking developments in this area,⁴ there is still great demand for the development of catalytic C–H trifluoromethylation methods that function under mild and simple conditions.

We were interested in developing a catalytic trifluoromethylation based on silver; in contrast to its Group 11 neighbour copper there have been few reports on silver-mediated trifluoromethylation^{4g,k,5} and none that we are aware of using silver catalysis. The redox catalysis of silver, comprising one electron steps between 0, +1, +2 and +3 oxidation states, has been scarcely exploited in synthesis relative to other late TMs^{6,7} and could offer productive catalytic pathways for trifluoromethylation. We have recently developed silver-catalysed decarboxylative C–H cross-coupling under oxidative radical conditions,⁸ and were keen to see if a similar approach was viable for C–H trifluoromethylation.

We started with a screen of reaction conditions based around TMSCF_3 as the trifluoromethylating agent.⁹ The groups of Sanford, Bräse and Wang have recently demonstrated the compatibility of this reagent with stoichiometric silver salts,^{4g,k,5c} encouraging us that it could form the basis of a catalytic system. Using 1,4-dimethoxybenzene (**1a**) as the substrate, we conducted an initial solvent screen using combinations of AgF , TMSCF_3 and PhI(OAc)_2 (Table 1). We worked at room temperature under air throughout, with the aim of developing a mild reaction with as broad a functional group tolerance as possible. The reaction proved sensitive to solvent choice with initially only MeCN from a selection of common organic solvents producing any reaction (entries 1 and 2). DMSO proved more effective still, affording the trifluoromethylated compound **2a** in 51% conversion (entry 3). Fluoride was not a requirement, with Ag_2CO_3 being similarly effective at promoting reaction (entry 4). Alternative oxidants did not improve on PhI(OAc)_2 (entries 5 and 6), and the use of a

Table 1 Ag-catalysed trifluoromethylation: reaction optimisation

Entry ^a	Catalyst (equiv.)	Oxidant	Solvent	Yield ^b (%)
1	AgF (1)	PhI(OAc)_2	Solvent ^c	0
2	AgF (1)	PhI(OAc)_2	MeCN	26
3	AgF (1)	PhI(OAc)_2	DMSO	51
4	Ag_2CO_3 (0.5)	PhI(OAc)_2	DMSO	48
5	AgF (1)	PhI(TFA)_2	DMSO	5
6	AgF (1)	$\text{K}_2\text{S}_2\text{O}_8$	DMSO	20
7 ^d	AgF (1)	PhI(OAc)_2	DMSO	35
8 ^e	AgF (0.25)	PhI(OAc)_2	DMSO	55
9 ^{e,f}	AgF (0.25)	PhI(OAc)_2	DMSO	55 (58 ^g)
10 ^{e,h}	AgF (0.25)	PhI(OAc)_2	DMSO	60

^a 1,4-Dimethoxybenzene **1a** (0.3 mmol), TMSCF_3 (1.2 mmol), oxidant (0.6 mmol), F^- source or base, solvent (1.0 mL), room temperature, 20 h. ^b Yields determined by ^{19}F NMR using 4-fluoroanisole as the internal standard. ^c THF, 1,4-dioxane, MeOH, $(\text{CF}_3)_2\text{CHOH}$, DCE, DCM.

^d Under N_2 . ^e Slow addition of AgF to the stirring mixture of **1a**, TMSCF_3 and PhI(OAc)_2 in DMSO. ^f 2 equiv. of TMSCF_3 . ^g Isolated yield.

^h TESCF_3 (2 equiv.) instead of TMSCF_3 .

^a School of Chemistry, University of Manchester, Oxford Rd, Manchester, M13 9PL, UK. E-mail: michael.greaney@manchester.ac.uk

^b Syngenta, Jealott's Hill International Research Centre, Bracknell, RG42 6EY, UK

† Electronic supplementary information (ESI) available: Synthesis and characterisation data for all new compounds. See DOI: 10.1039/c3cc41829d



nitrogen atmosphere led to a reduction in yield (entry 7). Crucially, sub-stoichiometric amounts of silver salts proved equally effective (entries 8–10), indicating that a catalytic reaction was feasible. We settled on conditions of AgF (25 mol%) with TMSCF₃ (2 equiv.) and PhI(OAc)₂ (2 equiv.), at room temperature (entry 9) to take forward. The use of the more stable (and expensive) TESCF₃ reagent gave only marginal improvement (entry 10), so we continued with the cheaper TMSCF₃ reagent.

Substrate scope investigations established that the procedure was effective for a variety of electron rich arenes with broad substrate scope tolerance (Table 2). For unsymmetrical substrates isomeric mixtures were generally observed, with regioselectivities consistent with radical S_{Ar}H addition (*vide infra*). Importantly, the reaction was compatible with halogen groups, illustrating an orthogonal reactivity to conventional C–X trifluoromethylations whereby neighbouring C–H bonds undergo preferential reaction. The useful building blocks 2f, 2g, 2h and 2i were prepared in this fashion. Electron-withdrawing groups such as aldehyde (2j), ketone (2k) and ester (2l) were likewise tolerated without problem. Importantly, dialkylanilines could be trifluoromethylated, a key class of building block that has rarely featured in C–H trifluoromethylation reports.^{4i,10} A slight preference for *ortho* over *para* selectivity was observed for simple dimethylamine (2m), with bromo substitution also tolerated (2n) along with N-acylation (2o). We were pleased to observe that the reaction was also effective for un-activated arenes (2p, 2q and 2r), although these substrates did require an excess of the arene and the reaction temperature raised to 70 °C.

The reaction could be extended to heteroarenes with N–Me pyrroles in particular being excellent substrates (2s, 2t). Electron-withdrawing groups on the heteroarene nucleus were well-tolerated (2t), but on nitrogen less so (N–Boc, 2u). Furans (2v), thiophenes (2w, 2x) and indoles (2y) were all productive, indicating that the method is viable for the major classes of π-excessive heterocycle. π-Deficient heteroarenes, by contrast, were not generally effective in the reaction but could be efficiently captured by masking the azine nucleus with electron-donating groups (2aa).


We next turned to the trifluoromethylation of more complex, biologically active molecules – a major driver for the development of new methods in this area. Introduction of the CF₃ group at unactivated C–H positions represents a very versatile approach to fluorine incorporation for modulation of biological activity,^{4c,d,j} demanding mild reaction conditions that are tolerant of functional groups and reasonable stoichiometries with respect to the (often valuable) C–H substrate. Accordingly, we extended the reaction to trifluoromethylate some more complex molecules in the agrochemistry field, an area where the CF₃ group is particularly prevalent. We could successfully incorporate the CF₃ group into the commercial herbicides pyriflatalid¹¹ and napropamide¹² (Scheme 1). The functional group tolerance of the reaction was illustrated by sulfide, lactone and α-hydroxyamide functionality all being stable to the reaction conditions (Scheme 1, 3 and 4).

A radical mechanism is implicated for the trifluoromethylation reaction,¹³ as radical quenching reactions using TEMPO and galvinoxyl radical both shut down the reaction, with the TEMPO–CF₃ adduct being clearly observed in the crude ¹⁹F NMR. The electrophilic CF₃ radical usually (but with some exceptions)^{4c,j}

Table 2 Ag-catalysed trifluoromethylation: substrate scope^{a,b}

Ar–H 1	TMSCF ₃ (2 equiv.) PhI(OAc) ₂ (2 equiv.) AgF (25 mol%) DMSO, rt, 20 h	Ar–CF ₃ 2


^a 1 (0.3 mmol), TMSCF₃ (0.6 mmol), PhI(OAc)₂ (0.6 mmol), AgF (0.075 mmol), DMSO (1.0 mL), room temperature, 20 h. ^b Isolated yields. For isomer mixtures, the minor regioisomeric position is labeled with *. ^c Yields determined by ¹⁹F NMR using 4-fluoroanisole as the internal standard. ^d Reaction conducted at 70 °C, 5–10 equiv. of arene.



Scheme 1 Agrochemical trifluoromethylation.

displays a marked preference for electron rich substrates, as seen here, underlining the likelihood of a radical pathway. A possible





Scheme 2 Silver-catalysed radical trifluoromethylation.

mechanism is shown in Scheme 2 whereby TMSCF₃ is oxidised to the CF₃ radical, followed by S_AH addition, then a second one electron oxidation and proton loss to give the product 2. Control experiments to investigate the role of silver in the first step of the proposed mechanism indicated that AgF alone was insufficiently oxidizing to generate CF₃[•] (mixing AgF with TMSCF₃ in the presence of TEMPO in DMSO at room temperature gave only trace quantities of TEMPO-CF₃). PhI(OAc)₂ alone was moderately effective (44% NMR yield of TEMPO-CF₃) and the combination of PhI(OAc)₂ and AgF highly effective (91% NMR yield).¹⁴ The background oxidizing activity of the hypervalent iodine reagent could be quantified in the trifluoromethylation of 1,4-dimethoxybenzene **1a** in the absence of any silver salt, producing a low conversion to the trifluoromethylated product **2a** (26% NMR yield).

Alternative mechanisms were investigated by treating dimethoxy-anisole **1a** with *in situ* prepared AgCF₃^{5b} in both MeCN and DMSO as solvents. No reaction could be observed in each case, suggesting organometallic AgCF₃ intermediates are not participating under our reaction conditions. A further control experiment with Togni's reagent^{4a} in DMSO at room temperature gave no reaction, ruling out simple S_EAr attack on an electrophilic CF₃ source. Finally, we considered the possibility of initial arene oxidation by PhI(OAc)₂, followed by CF₃ anion addition to a cationic arene intermediate. Extensive work by Kita has demonstrated the C–H functionalization of electron rich arenes using PhI(TFA)₂ in the presence of stoichiometric BF₃·OEt₂ and nucleophiles.¹⁵ It seems the present conditions are not sufficiently oxidizing to enable an analogous pathway, as a control reaction in the absence of TMSCF₃ gave no reaction, where some degree of homocoupling would be expected if this mechanism was in operation.

In conclusion, we have developed a silver-catalysed trifluoromethylation system for electron rich aromatic and heteroaromatic substrates. The reaction works at room temperature under air, does not require excessive stoichiometries of substrate or reagent, and is operationally simple to carry out. The application of this chemistry to new trifluoromethylation substrates will be the subject of future work in our laboratory.

We thank Syngenta, the University of Manchester and the EPSRC for funding (Leadership Fellowship to M.F.G.), and the EPSRC mass spectrometry service at the University of Swansea.

Notes and references

- (a) A. Studer, *Angew. Chem., Int. Ed.*, 2012, **51**, 8950; (b) O. A. Tomashenko and V. V. Grushin, *Chem. Rev.*, 2011, **111**, 4475; (c) T. Furuya, A. S. Kamlet and T. Ritter, *Nature*, 2011, **473**, 470.
- Recent examples of stoichiometric metal-mediated trifluoromethylation: (a) K. A. McReynolds, R. S. Lewis, L. K. G. Ackerman, G. G. Dubinina, W. W. Brennessel and D. A. Vicic, *J. Fluorine Chem.*, 2010, **131**, 1108; (b) Y. Ye, N. D. Ball, J. W. Kampf and M. S. Sanford, *J. Am. Chem. Soc.*, 2010, **132**, 14682; (c) T. D. Senecal, A. T. Parsons and S. L. Buchwald, *J. Org. Chem.*, 2011, **76**, 1174; (d) N. D. Ball, J. B. Gary, Y. Ye and M. S. Sanford, *J. Am. Chem. Soc.*, 2011, **133**, 7577; (e) C.-P. Zhang, Z.-L. Wang, Q.-Y. Chen, C.-T. Zhang, Y.-C. Gu and J.-C. Xiao, *Angew. Chem., Int. Ed.*, 2011, **50**, 1896; (f) H. Morimoto, T. Tsubogo, N. D. Litvinas and J. F. Hartwig, *Angew. Chem., Int. Ed.*, 2011, **50**, 3793; (g) O. A. Tomashenko, E. C. Escudero, M. M. Belmonte and V. V. Grushin, *Angew. Chem., Int. Ed.*, 2011, **50**, 7655; (h) Y. Ye, S. A. Künzi and M. S. Sanford, *Org. Lett.*, 2012, **14**, 4979.
- Cu and Pd-catalysed trifluoromethylation of functionalised arenes: (a) M. Oishi, H. Kondo and H. Amii, *Chem. Commun.*, 2009, 1909; (b) E. J. Cho, T. D. Senecal, T. Kinzel, Y. Zhang, D. A. Watson and S. L. Buchwald, *Science*, 2010, **328**, 1679; (c) R. Shimizu, H. Egami, T. Nagi, J. Chae, Y. Hamashima and M. Sodeoka, *Tetrahedron Lett.*, 2010, **51**, 5947; (d) T. Knauber, F. Arıkan, G.-V. Roeschenthaler and L. J. Goossen, *Chem.-Eur. J.*, 2011, **17**, 2689; (e) J. Xu, D.-F. Luo, B. Xiao, Z.-J. Liu, T.-J. Gong, Y. Fu and L. Liu, *Chem. Commun.*, 2011, **47**, 4300; (f) T. Liu and Q. Shen, *Org. Lett.*, 2011, **13**, 2342; (g) Y. Ye and M. S. Sanford, *J. Am. Chem. Soc.*, 2012, **134**, 9034; (h) T. Schareina, X.-F. Wu, A. Zapf, A. Cotte, M. Gotta and M. Beller, *Top. Catal.*, 2012, **55**, 426.
- Recent C–H Trifluoromethylation of arenes: (a) M. S. Wiehn, E. V. Vinogradova and A. Togni, *J. Fluorine Chem.*, 2009, **131**, 951; (b) X. Wang, L. Truesdale and J.-Q. Yu, *J. Am. Chem. Soc.*, 2010, **132**, 3648; (c) Y. Ji, T. Brueckl, R. D. Baxter, Y. Fujiwara, I. B. Seiple, S. Su, D. G. Blackmond and P. S. Baran, *Proc. Natl. Acad. Sci. U. S. A.*, 2011, **108**, 14411; (d) D. A. Nagib and D. W. C. MacMillan, *Nature*, 2011, **480**, 224; (e) R. N. Loy and M. S. Sanford, *Org. Lett.*, 2011, **13**, 2548; (f) X. Mu, S. Chen, X. Zhen and G. Liu, *Chem.-Eur. J.*, 2011, **17**, 6039; (g) Y. Ye, S. H. Lee and M. S. Sanford, *Org. Lett.*, 2011, **13**, 5464; (h) L. Chu and F.-L. Qing, *J. Am. Chem. Soc.*, 2012, **134**, 1298; (i) E. Mejia and A. Togni, *ACS Catal.*, 2012, **2**, 521; (j) Y. Fujiwara, J. A. Dixon, F. O'Hara, E. D. Funder, D. D. Dixon, R. A. Rodriguez, R. D. Baxter, B. Herlé, N. Sach, M. R. Collins, Y. Ishihara and P. S. Baran, *Nature*, 2012, **492**, 95; (k) A. Hafner and S. Bräse, *Angew. Chem., Int. Ed.*, 2012, **51**, 3713; (l) X.-G. Zhang, H.-X. Dai, M. Wasa and J.-Q. Yu, *J. Am. Chem. Soc.*, 2012, **134**, 11948; (m) X. Wu, L. Chu and F.-L. Qing, *Tetrahedron Lett.*, 2013, **54**, 249.
- (a) W. E. Tyrra, *J. Fluorine Chem.*, 2001, **112**, 149; (b) M. M. Kremlev, A. I. Mushta, W. E. Tyrra, D. Naumann, H. T. M. Fischer and Y. L. Yagupolskii, *J. Fluorine Chem.*, 2007, **128**, 1385; (c) Z. Weng, R. Lee, W. Jia, Y. Yuan, W. Wang, X. Feng and K.-W. Huang, *Organometallics*, 2011, **30**, 3229.
- Silver in organic chemistry*, ed. M. Harmata, Wiley, NY, 2010.
- For silver-catalysed fluorination, see: P. Tang, T. Furuya and T. Ritter, *J. Am. Chem. Soc.*, 2010, **132**, 12150.
- (a) S. Seo, M. Slater and M. F. Greaney, *Org. Lett.*, 2012, **14**, 2650; (b) S. Seo, J. B. Taylor and M. F. Greaney, *Chem. Commun.*, 2012, **48**, 8270.
- G. K. S. Prakash and A. K. Yudin, *Chem. Rev.*, 1997, **97**, 757.
- T. Akiyama, K. Kato, M. Kajitani, Y. Sakaguchi, J. Nakamura, H. Hayashi and A. Sugimori, *Bull. Chem. Soc. Jpn.*, 1988, **61**, 3531.
- C. Luthy, H. Zondler, T. Rapold, G. Seifert, B. Urwyler, T. Heinis, H. C. Steinrücken and J. Allen, *Pest Manage. Sci.*, 2001, **57**, 205.
- Stauffer Chemical Company, *Br. Pat.*, GB1066606, 1967.
- Examples of radical trifluoromethylation: (a) A. Gregorcic and M. Zupan, *J. Org. Chem.*, 1979, **44**, 4120; (b) C. Wakelman and M. Tordeux, *Chem. Commun.*, 1987, 1701; (c) H. Sawada, M. Nakayama, M. Yoshida, T. Yoshida and N. Kamigata, *J. Fluorine Chem.*, 1990, **46**, 423; (d) B. R. Langlois, E. Laurent and N. Roidot, *Tetrahedron Lett.*, 1991, **32**, 7525; (e) K. L. Kirk, M. Nishida, S. Fujii and H. Kimoto, *J. Fluorine Chem.*, 1992, **59**, 197; (f) C. Lai and T. E. Mallouk, *Chem. Commun.*, 1993, 1359; (g) N. Kamigata, T. Ohtsuka, T. Fukushima, M. Yoshida and T. Shimizu, *J. Chem. Soc., Perkin Trans. 1*, 1994, 1339; (h) J.-B. Tommasino, A. Brondex, M. Médebielle, M. Thomalla, B. R. Langlois and T. Billard, *Synlett*, 2002, 1697; (i) T. Kino, Y. Nagase, Y. Ohtsuka, K. Yamamoto, D. Uraguchi, K. Tokuhisa and T. Yamakawa, *J. Fluorine Chem.*, 2010, **131**, 98.
- ArI(OAc)₂ could potentially generate Ag(II) as the oxidant for TMSCF₃ *in situ* under the reaction conditions. For *in situ* silver(II) generation from catalytic Ag(I) and a stoichiometric oxidant in decarboxylation chemistry, see: J. M. Anderson and J. K. Kochi, *J. Am. Chem. Soc.*, 1970, **92**, 1651.
- T. Dohi, M. Ito, N. Yamaoka, K. Morimoto, H. Fujioka and Y. Kita, *Tetrahedron*, 2009, **65**, 10797.

