# ChemComm

#### COMMUNICATION

## **RSC**Publishing

View Article Online View Journal | View Issue

 Dpen Access Article. Published on 23 October 2013. Downloaded on 8/13/2025 2:44:19 PM.

 This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Cite this: Chem. Commun., 2013, 49, 11176

Received 3rd September 2013, Accepted 7th October 2013

DOI: 10.1039/c3cc46720a

www.rsc.org/chemcomm

## Epoxidation of bromoallenes connects red algae metabolites by an intersecting bromoallene oxide – Favorskii manifold<sup>†</sup>

D. Christopher Braddock,\* James Clarke and Henry S. Rzepa

DMDO epoxidation of bromoallenes gives directly  $\alpha$ , $\beta$ -unsaturated carboxylic acids under the reaction conditions. Calculated ( $\omega$ B97XD/ 6-311G(d,p)/SCRF = acetone) potential energy surfaces and <sup>2</sup>H- and <sup>13</sup>C-labeling experiments are consistent with bromoallene oxide intermediates which spontaneously rearrange *via* a bromocyclopropanone in an intersecting bromoallene oxide – Favorskii manifold.

The remarkably wide structural diversity and complexity of halogenated  $C_{15}$  acetogenin metabolites isolated from marine red algae of *Laurencia* species<sup>1</sup> continue to stimulate innovative efforts in their target synthesis,<sup>2</sup> in the discovery of new synthetic transformations<sup>3</sup> and in advancing biosynthetic hypotheses.<sup>4</sup> A recent re-isolation<sup>5</sup> of obtusallene IV (1)<sup>6</sup> from *Laurencia marilzae* provided also 12-epoxyobtusallene IV (2) and unnamed  $\alpha$ , $\beta$ unsaturated carboxylate ester (3) with an identical macrocycle to epoxybromoallene 2 (Fig. 1). It seems reasonable to connect *E*-alkene 1 and *trans*-epoxide 2 biogenetically *via* enzymatic epoxidation,<sup>7</sup> and on the basis of their co-isolation, we propose



Fig. 1 Metabolites 1–3 from *Laurencia marilzae* and proposed biogenesis *via* epoxidation events.

Department of Chemistry, Imperial College London, London, SW7 2AZ, UK. E-mail: c.braddock@imperial.ac.uk; Fax: +44 (0)2075945805; Tel: +44 (0)2075945772

<sup>†</sup> Electronic supplementary information (ESI) available: Notes §, ¶, ¥, ††, \*\*, ‡‡, §§, ¶¶, ¥¥, †††, \*\*\*, ‡‡; general experimental; experimental details and characterising data for compounds leading to bromoallenes 4 (including ESI Scheme S1 for the synthesis of bromoallenes 4), (1-<sup>2</sup>H)-4 and (1-<sup>13</sup>C-4) and epoxidation thereof leading to *E*- and *Z*-5, (*E*-2-<sup>2</sup>H)- and (*Z*-2-<sup>2</sup>H)-5, and (*E*-2-<sup>13</sup>C)-and (*Z*-2-<sup>13</sup>C)-5; Copies of <sup>1</sup>H and <sup>13</sup>C spectra for all compounds showing <sup>2</sup>H and <sup>13</sup>C totopic shifts and coupling constants where appropriate; ESI references. See DOI: 10.1039/c3cc46720a

that bromoallene 2 and  $\alpha$ , $\beta$ -unsaturated carboxylate 3 may also be connected biogenetically by epoxidation.

While the epoxidation of allenes<sup>8,9</sup> and vinyl bromides<sup>10</sup> has been studied, the epoxidation of bromoallenes has not been reported.<sup>11</sup> Herein, we report the hitherto unknown direct conversion of bromoallenes to  $\alpha$ , $\beta$ -unsaturated carboxylic acids *via* an initial epoxidation event and the presumed intermediacy of a bromoallene oxide. We also show by computational modeling and <sup>2</sup>H- and <sup>13</sup>C-labeling studies that the latter's spontaneous reorganization to an  $\alpha$ , $\beta$ -unsaturated carboxylic acid under the reaction conditions is consistent with a bromocyclopropanone intermediate in an intersecting allene oxide – Favorskii manifold.

Bromoallene 4<sup>12</sup> was selected as a suitable substrate for investigating epoxidation and was synthesized by a standard sequence from heptanal (ESI<sup>†</sup>).<sup>13</sup> Much to our delight, epoxidation of bromoallene 4 using dimethyl dioxirane (DMDO), generated either *in situ*<sup>14</sup> or as a solution (ESI<sup>†</sup>)<sup>15</sup> (Scheme 1), gave a mixture of *Z* and *E*- $\alpha$ , $\beta$ -unsaturated carboxylic acids 5 directly in low but reproducible yields (note §, ESI<sup>†</sup>). The low yields can be attributed to decomposition of DMDO<sup>16*a*</sup> under the reaction conditions to methyl radicals,<sup>16*b*</sup> and subsequent radical attack on either of the products *or* starting materials (note ¶, ESI<sup>†</sup>).

Mechanistically, we invoke the following pathway for the formation of  $\alpha$ , $\beta$ -unsaturated carboxylic acids from DMDO mediated epoxidation of bromoallenes (Fig. 2). Initial epoxidation of the bromoallene would give bromoallene oxides of the type **A** and/or **B** (note  $\ddagger$ , ESI $\ddagger$ ). Spontaneous epoxide opening<sup>8c</sup> via bromo oxyallyl cations **C** and **D** (note  $\dagger\dagger$ , ESI $\dagger$ ) respectively converge on the same bromocyclopropanone **E**. This intermediate now intersects with the Favorskii rearrangement manifold of  $\alpha$ , $\alpha$ - and  $\alpha$ , $\alpha'$ -dibromoketones where the resulting bromocyclopropanones **E** are known to collapse after attack by water giving hydrate **F** to  $\alpha$ , $\beta$ -unsaturated carboxylic acids **5** (note **\*\***, ESI $\ddagger$ ).<sup>17,18</sup> Evidently, there is sufficient water in the dioxirane solution to function as a nucleophile here (note  $\ddagger\ddagger$ , ESI $\ddagger$ ).







Fig. 2 Mechanistic rationale for conversion of bromoallenes into  $\alpha$ , $\beta$ -unsaturated carboxylic acids, with the carbon atoms of the functional groups numbered 1–3 showing an interchange of carbon atoms 1 and 2 (see also interactive Fig. 2 in HTML version of this article).

Interestingly, regardless of the initial site of epoxidation, this mechanism predicts that carbon atoms 1 and 2 in bromoallene 4 interchange positions in the  $\alpha$ , $\beta$ -unsaturated carboxylic acid products 5.

This mechanism can be subjected to scrutiny via density functional level ( $\omega$ B97XD/6-311G(d,p)/SCRF = acetone)<sup>19</sup> exploration of the potential energy surface (R = H, Me, presented as an interactive version of Fig. 2 (ref. 20) via a digital data repository<sup>21</sup>). Oxygen transfer from dimethyldioxirane to form both A and B (TS1) have thermally accessible free energy activation barriers  $\Delta G_{208}^{\dagger}$  (R = H, 26.8 for **A**, 27.3 for **B**; R = Me, 26.8 for **A**, 24.6 kcal mol<sup>-1</sup> for **B**), followed by a second, lower energy dyotropic rearrangement (TS2) to give E. An intrinsic reaction coordinate (IRC) reveals that TS2 (R = H,Me) represents the concerted transformation of A or B to E, with C/D acting as "hidden intermediates" in the process.<sup>22</sup> Such hidden intermediates can be potentially transformed to real ones by tuning the substituents, and in this instance changing R from H or Me to OMe is predicted to accomplish this by stabilization of C/D (see interactive Fig. 2). TS2 itself (R = Me) has some early character of C/D; the C-Br bond is calculated to initially contract in length due to a significant stabilising resonance contribution of Br lone pairs, from 1.924/1.896 Å (A and B respectively) via 1.840/1.885 (TS2), 1.856/1.868 (C/D acting as hidden intermediates) to 1.921/1.922 Å (E).<sup>23</sup> Calculations having demonstrated the thermal accessibility of the epoxidation-bromocyclopropanone sequence, <sup>2</sup>H- and <sup>13</sup>C-labeling experiments were necessary to verify the overall reorganization (4 to A/B to E to F to 5, Fig. 2) of the carbon framework.<sup>24</sup>

Deuterated bromoallene  $(1-{}^{2}H)-4$  was prepared by addition of ethynylmagnesium bromide to heptanal, *in situ* deprotonation of the propargylic alkoxide with *n*-butyllithium and quenching with



Scheme 2 Synthesis of deuterated bromoallene (1-<sup>2</sup>H)-4





MeOH-d<sub>4</sub> to give labeled propargylic alcohol (1-<sup>2</sup>H)-**6** (Scheme 2). Subsequent alcohol trisylation<sup>25</sup> gave (1-<sup>2</sup>H)-**7**, and  $S_N 2'$  displacement of the trisylate with bromide under the action of LiCuBr<sub>2</sub> (ref. 26) provided bromoallene (1-<sup>2</sup>H)-**4** with 70% deuterium incorporation at the 1-position.<sup>†</sup>

<sup>13</sup>C-labeled bromoallene (1-<sup>13</sup>C)-4 was similarly targeted, commencing with silyl enol ether **8** formation<sup>27</sup> from octanal (Scheme 3). Oxidation using *m*CPBA gave interrupted Rubottom<sup>28</sup> adduct **9**, which could be acetylated to give acetate **10**. Desilylation using buffered TBAF<sup>29</sup> revealed protected α-hydroxyaldehyde **11**, which we planned to use in a Wittig reaction with a suitably <sup>13</sup>C-labeled phosphorous ylid. To the best of our knowledge, there is only a single report<sup>30</sup> using methyltriphenylphosphonium iodide to generate the Stork–Wittig reagent<sup>31</sup> using an *in situ* deprotonation– iodination–deprotonation procedure which we adapted using <sup>13</sup>C-labeled salt **12** – available from relatively inexpensive 99% atom <sup>13</sup>C-labeled methyl iodide – to give vinyl iodides *Z*-(1-<sup>13</sup>C)-**13**, *E*-(1-<sup>13</sup>C)-**13** and diiodide (1-<sup>13</sup>C)-**14**.<sup>32</sup> Acetate deprotection as a mixture gave the corresponding alcohols *Z*-(1-<sup>13</sup>C)-**15**, *E*-(1-<sup>13</sup>C)-**15** and (1-<sup>13</sup>C)-**16** all with 99% <sup>13</sup>C at the alkene terminus.<sup>†</sup>

Dehydrohalogenation of *Z*- and *E*-iodides  $(1^{-13}C)$ -**15** in the presence of inseparable diiodide  $(1^{-13}C)$ -**16** with LDA gave propargylic alcohol  $(1^{-13}C)$ -**7** in good overall yield, with the unprecedented observation that LDA converts vinyl 1,1-diiodides into terminal alkynes also (note §§, ESI<sup>†</sup>). Interestingly, 4% of the alkyne product was found to be the 2-<sup>13</sup>C isotopomer (ESI<sup>†</sup>), implicating a 1,1-elimination reaction pathway for diiodide **16** 



Fig. 3 <sup>1</sup>H NMR spectra of (a) (E-2-<sup>13</sup>C)-5 and (b) (Z-2-<sup>13</sup>C)-5 displaying the expected <sup>1</sup>J<sub>CH</sub> values for the  $\alpha$ -vinyl protons.

and competitive alkyl group migration from a vinylidene intermediate (note  $\P\P$ , ESI<sup>†</sup>). Alcohol (1-<sup>13</sup>C)-7 was then converted to the desired bromoallene (1-<sup>13</sup>C)-7 (as 4% of its 2-<sup>13</sup>C-isotopomer, ESI<sup>†</sup>) as previously described (*cf.*, Scheme 2).



With (1-<sup>2</sup>H)-4 and (1-<sup>13</sup>C)-4 in hand, epoxidation with DMDO was conducted. For deuterated (1-<sup>2</sup>H)-4, after the reaction was conducted in the usual manner (*cf.*, Scheme 1), *E*-(2-<sup>2</sup>H)-5 and *Z*-(2-<sup>2</sup>H)-5 were isolated each showing 65% deuteration at the  $\alpha$ -position only (note ‡, \Vec{T}, ESI†). Evidently, this result is consistent with the proposed mechanism (*cf.*, Fig. 2) (note  $\dagger \dagger \dagger \dagger$ , ESI†). More compellingly, epoxidation of bromoallene (1-<sup>13</sup>C)-4 gave (*E*-2-<sup>13</sup>C)-5<sup>33</sup> and (*Z*-2-<sup>13</sup>C)-5 (28% isolated yield) where carbon atoms 1 and 2 from the bromoallene have entirely interchanged positions, giving also 4% of each of the (*E*-1-<sup>13</sup>C)-5 and (*Z*-1-<sup>13</sup>C)-5 isotopomers (ESI†). The expected <sup>1</sup>J<sub>CH</sub> coupling constants experienced by the  $\alpha$ -vinyl protons of the major isotopomers are clearly apparent in their <sup>1</sup>H NMR spectra (Fig. 3).

In conclusion we have established that the hitherto unknown direct conversion of bromoallenes to  $\alpha$ , $\beta$ -unsaturated carboxylic acids using DMDO is consistent with an initial epoxidation event (note \*\*\*, ESI†) followed by a spontaneous reorganization *via* a bromocyclopropanone, a mechanism supported by calculations, in an intersecting bromoallene oxide – Favorskii manifold. These experiments support the proposed biogenesis of  $\alpha$ , $\beta$ -unsaturated carboxylate 3 from bromoallene 2 by epoxidation (note ‡‡‡, ESI†).

We thank the EPSRC for DTG funding (to J. C.).

#### Notes and references

- (a) J. W. Blunt, B. R. Copp, R. A. Keyzers, M. H. G. Munroa and M. R. Prinsep, *Nat. Prod. Rep.*, 2013, **30**, 237–323 and earlier reviews in this series; (b) B.-G. Wang, J. B. Gloer, N.-Y. Ji and J.-C. Zhao, *Chem. Rev.*, 2013, **113**, 3632–3685.
- 2 For a comprehensive review of the synthesis of medium ring ethers from *Laurencia* sp., see: (a) K. Fujiwara, *Top. Heterocycl. Chem.*, 2006, 5, 97–148; (b) For recent leading examples see: B. S. Dyson, J. W. Burton, T.-i. Sohn, B. Kim, H. Bae and D. Kim, *J. Am. Chem. Soc.*, 2012, 134, 11781–11790(c) M. J. Kim, T.-i. Sohn, D. Kim and R. S. Paton, *J. Am. Chem. Soc.*, 2012, 134, 20178–20188 and references cited therein.
- 3 For recent representative examples see: (*a*) S. Keshipeddy, I. Martínez, B. F. Castillo II, M. D. Morton and A. R. Howell, *J. Org. Chem.*, 2012, 77, 7883–7890; (*b*) S. A. Snyder, A. P. Brucks, D. S. Treitler and I. Moga, *J. Am. Chem. Soc.*, 2012, **134**, 17714–17721; (*c*) S. A. Snyder, D. S. Treitler, A. P. Brucks and W. Sattler, *J. Am. Chem. Soc.*, 2011, **133**, 15898–15901; (*d*) N. Ortega, V. S. Martín and T. Martín, *J. Org. Chem.*, 2010, 75, 6660–6672 and references cited therein.
- 4 For a review see: (a) A. Murai, in Comprehensive Natural Products Chemistry, ed. D. H. R. Barton, O. Meth-Cohn and K. Nakinishi, Elsevier, Oxford, 1999, vol. 1, pp. 303–324. For representative examples see: (b) K. J. Bonney and D. C. Braddock, J. Org. Chem., 2012, 77, 9574–9584; (c) A. Gutiérrez-Cepeda, J. J. Fernández, M. Norte and M. L. Souto, Org. Lett., 2011, 13, 2690–2693; (d) D. C. Braddock, D. S. Millan, Y. Perez-Fuertes, R. H. Pouwer, R. N. Sheppard, S. Solanki and A. J. P. White, J. Org. Chem., 2009, 74, 1835–1841; (e) D. C. Braddock, Org. Lett., 2006, 8, 6055–6058 and references cited therein.
- 5 A. Gutiérrez-Cepeda, J. J. Fernández, L. V. Gil, M. López-Rodríguez, M. Norte and M. L. Souto, J. Nat. Prod., 2011, 74, 441–448.
- 6 (a) G. Guella, G. Chiasera, I. Mancini, A. Öztunç and F. Pietra, *Chem.-Eur. J.*, 1997, 3, 1223–1231; (b) M. L. Ciavatta, M. Gavagnin, R. Puliti, G. Cimino, E. Martínez, J. Ortea and C. A. Mattia, *Tetrahedron*, 1997, 53, 17343–17350.

- 7 For reviews on enzymatic epoxidation of alkenes see: (a) M. Sono,
  M. P. Roach, E. D. Coulter and J. H. Dawson, *Chem. Rev.*, 1996, 96, 2841–2887; (b) P. R. Ortiz de Montellano and J. J. De Voss, *Nat. Prod. Rep.*, 2002, 19, 477–493.
- 8 For reviews see: (a) G. L'abbé, Angew. Chem., Int. Ed. Engl., 1980, 19, 276–289; (b) W. Smadja, Chem. Rev., 1983, 83, 263–320; (c) T. H. Chan and B. S. Ong, Tetrahedron, 1980, 36, 2269–2289.
- 9 For the first isolated allene oxide and its thermal rearrangement to a cyclopropanone see: R. L. Camp and F. D. Greene, *J. Am. Chem. Soc.*, 1968, **90**, 7349.
- 10 For an early report on their synthesis and reactivity of 2-bromooxiranes see: (*a*) A. Hassner and P. Catsoulacos, *J. Org. Chem.*, 1967, 32, 549–553; For a representative naturally occurring 2-bromooxirane see: (*b*) K. Watanabe, M. Sekine and K. Iguchi, *J. Nat. Prod.*, 2003, 66, 1434–1440.
- 11 There is a single report of a bromoallene oxide functionality: ethyl 2-bromo-3-(diphenylmethylene) oxirane-2-carboxylate was reported in a study of ketenes and aliphatic diazo compounds: H. Staudinger and T. Reber, *Helv. Chim. Acta*, 1921, **4**, 3–23.
- 12 P. C. Ravikumar, L. Yao and F. F. Fleming, J. Org. Chem., 2009, 74, 7294–7299.
- 13 See for example: D. C. Braddock, R. Bhuva, Y. Pérez-Fuertes, R. Pouwer, C. A. Roberts, A. Ruggiero, E. S. E. Stokes and A. J. P. White, *Chem. Commun.*, 2008, 1419–1421 and references cited therein.
- 14 J. K. Crandall, D. J. Batal, F. Lin, T. Reix, G. S. Nadol and R. A. Ng, *Tetrahedron*, 1992, **48**, 1427–1448 and references therein.
- 15 (a) R. W. Murray and M. Singh, Org. Synth., 1997, 74, 91. For the use of cyclic ketones as dioxirane precursors see: (b) R. W. Murray, M. Singh and R. Jeyaraman, J. Am. Chem. Soc., 1992, 114, 1346–1351.
- 16 For leading examples see: (a) M. Singh and R. W. Murray, J. Org. Chem., 1992, 57, 4263–4270; (b) N. N. Kabal'nova, D. V. Kazakov, N. M. Shishlov and V. V. Shereshovets, Russ. Chem. Bull., 1996, 45, 1481–1483.
- 17 For a review of the Favorskii reaction see: J. Mann, in *Comprehensive Organic Synthesis*, ed. B. M. Trost, I. Fleming and G. Pattenden, Pergamon Press, Oxford, 1991, vol. 3, ch. 3.7, pp. 839–859.
- 18 In a previous isolation from red algae *Bonnemaisonia*, both 1,1,3-tribromo-2-ketones and their proposed Favorskii products *E* and *Z*-3-bromo-2-alkenoic acids were co-isolates: O. J. McConnell and W. Fenical, *Phytochemistry*, 1980, **19**, 233–247.
- 19 J.-D. Chai and M. Head-Gordon, Phys. Chem. Chem. Phys., 2008, 6615-6620.
- 20 D. C. Braddock, J. Clarke and H. S. Rzepa, *Figshare*, 2013, DOI: 10.6084/ m9.figshare.785756 and the further digital repository links therein.
- 21 J. Downing, P. Murray-Rust, A. P. Tonge, P. Morgan, H. S. Rzepa, F. Cotterill, N. Day and M. J. Harvey, *J. Chem. Inf. Model.*, 2008, 48, 1571–1581. See also http://www.force11.org/AmsterdamManifesto for the Amsterdam Manifesto on data citation principles.
- 22 D. Cremer and E. Kraka, *Acc. Chem. Res.*, 2010, **43**, 591–601; H. S. Rzepa and C. Wentrup, *J. Org. Chem.*, 2013, **78**, 7565–7574.
- 23 Similar non-linear behavior of a bond is found in the related dyotropic rearrangement of dibromoethanes; D. C. Braddock, D. Roy, D. Lenoir, E. Moore, H. S. Rzepa, J. I.-C. Wu and P. von R. Schleyer, *Chem. Commun.*, 2012, **48**, 8943–8945.
- 24 For pioneering labeling work to elucidate the mechanism of the Favorskii rearrangement and to implicate a symmetrical intermediate, *viz.*, a cyclopropanone see: R. B. Loftfield, *J. Am. Chem. Soc.*, 1951, 73, 4707–4714.
- 25 T. A. Grese, K. D. Hutchinson and L. E. Overman, J. Org. Chem., 1993, 58, 2468–2477.
- 26 C. J. Elsevier, P. Vermeer, A. Gedanken and W. Runge, J. Org. Chem., 1985, 50, 364–367.
- 27 Y. Tanigichi, J. Inanaga and M. Yamaguchi, Bull. Chem. Soc. Jpn., 1981, 54, 3229–3230.
- 28 A. Hassner, R. H. Reuss and H. W. Pinnick, J. Org. Chem., 1975, 40, 3427–3429.
- 29 For a representative example see: J. S. Debenham, R. Rodebaugh and B. Fraser-Reid, *J. Org. Chem.*, 1997, **62**, 4591–4600.
- 30 W. Zhu, M. Jiménez, W.-H. Jung, D. P. Camarco, R. Balachandran, A. Vogt, B. W. Day and D. P. Curran, *J. Am. Chem. Soc.*, 2010, 132, 9175–9187.
- 31 G. Stork and K. Zhao, Tetrahedron Lett., 1989, 2173-2174.
- 32 For the unwanted formation of a vinyl 1,1-diiodide in a Stork–Wittig reaction using [Ph<sub>3</sub>PCH<sub>2</sub>I]I see: P. Li, J. Li, F. Arikan, W. Ahlbrecht, M. Dieckmann and D. Menche, *J. Org. Chem.*, 2010, 75, 2429–2444.
- 33 D. C. Braddock, J. Clarke and H. S. Rzepa, *Figshare*, 2013, DOI: 10.6084/m9.figshare.785753 viz. ref. 20 and 21 for explanation.