Organic & Biomolecular Chemistry

COMMUNICATION

View Article Online View Journal | View Issue

 Open Access Article. Published on 28 February 2013. Downloaded on 9/4/2025 12:56:22 PM.

 V
 This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Cite this: Org. Biomol. Chem., 2013, 11, 2249

Received 10th December 2012, Accepted 18th February 2013

DOI: 10.1039/c3ob27396b

www.rsc.org/obc

Synthesis of imidazo and benzimidazo[2,1-a]isoquinolines by rhodium-catalyzed intramolecular double C–H bond activation†

Vutukuri Prakash Reddy, Takanori Iwasaki and Nobuaki Kambe*

The rhodium-catalyzed intramolecular direct arylation of imidazole and benzimidazole derivatives *via* double C–H bond activation is described. This approach provides new access to a wide range of imidazo and benzimidazo[2,1-a]isoquinoline derivatives in moderate to high yields. This reaction provides an alternative method to the known Pd-catalyzed intramolecular oxidative cross-coupling reactions.

Introduction

Nitrogen-containing heterocycles and their derivatives are often found in natural products and in pharmaceuticals and agrochemicals.^{1,2} Since imidazo and benzimidazo[2,1-a]isoquinoline derivatives show interesting biological activities such as anticancer, anti-HIV-1, antimicrobial and antiviral properties,^{3,4} a variety of synthetic protocols have been developed for the preparation of imidazo and benzimidazo [2,1-a] isoquinoline derivatives based on condensation reactions⁵ and the photocyclization of 1-styrylimidazoles.⁶ In addition, coupling reactions catalyzed by transition metals, such as palladium and copper, have recently been reported and successfully applied to the preparation of such heterocyclic compounds through C-N and C-C cross-coupling.7 However, these protocols require the use of organohalides^{7a-d} or boranes^{7e} or both^{7f} as starting materials. These prefunctionalized starting materials are not readily available from commercial sources and are often difficult to synthesize. Therefore, the development of more general and convenient processes using readily accessible and inexpensive substrates is an important theme. As a promising strategy for the synthesis of the imidazo and

benzimidazo[2,1-*a*]isoquinoline derivatives, the intramolecular oxidative cross-coupling *via* cleavage of two C–H bonds would be a more straightforward and efficient route to these compounds.

It has long been known that Pd catalyzes the oxidative homo-coupling of arenes^{8a,b} and cross-coupling between arenes and olefins.^{8c,d} This transformation has been successfully applied to cyclization by intramolecular oxidative crosscoupling reactions.^{8e-m} This successful Pd-catalyzed functionalization of C–H bonds^{8n,o} prompted the study of Rh catalyzed systems leading to the recent remarkable development of useful transformations *via* C–H bond cleavage⁹ such as oxidative Heck-type reactions,¹⁰ oxidative aryl–aryl coupling,¹¹ as well as addition to carbon–carbon^{10c,d,k,q,12} or carbon–heteroatom¹³ unsaturated bonds.

During the course of our study on transition metal-catalyzed cross-coupling reactions,¹⁴ we developed a straightforward procedure for the synthesis of imidazo and benzimidazo-[2,1-a]isoquinoline derivatives. The method involves a rhodium-catalyzed cross-coupling reaction *via* double C–H bond cleavage as the first example of the rhodium-catalyzed intramolecular oxidative cross-coupling of heteroarenes with arenes, although the corresponding aryl-aryl intramolecular coupling reactions are known (Scheme 1).¹⁵



Scheme 1 Rhodium-catalyzed intramolecular oxidative cross-coupling reaction.

Department of Applied Chemistry, Graduate School of Engineering, Osaka University, Suita, Osaka 565-0871, Japan. E-mail: kambe@chem.eng.osaka-u.ac.jp; Fax: +81-6-6879-7390; Tel: +81-6-6879-7390

 $[\]dagger$ Electronic supplementary information (ESI) available: Full experimental details, characterization data, ¹H and ¹³C NMR spectra for new products. See DOI: 10.1039/c3ob27396b

Table 1 Optimization of the rhodium-catalyzed intramolecular C–H arylation^a



^{*a*} Reaction conditions: **1a** (0.25 mmol), [Cp*RhCl₂]₂ (3.0 mol%), Cu-(OAc)₂ (0.6 mmol), additive (0.5 mmol), solvent (1.0 mL), 24 h, 140 °C. ^{*b*} Isolated yields. ^{*c*} Rh (10 mol%). ^{*d*} [Cp*RhCl₂]₂ (1.5 mol%). ^{*e*} At 70 °C. Cp*: pentamethylcyclopentadienyl; cod: 1,5-cyclooctadiene; nbd: bicyclo[2.2.1]hepta-2,5-diene.

Results and discussion

We carried out the reaction under different conditions using (*Z*)-1-styryl-1*H*-benzimidazole (**1a**) as a model substrate and the results are summarized in Table 1. Among the various solvents tested, DMA and mesitylene afforded the best results (Table 1, entries 1, 2, 5, 8), whereas DMF and toluene gave lower yields (entries 3, 4, 9, 10). PivOH was found to be an effective additive (entry 5). Among the Rh complexes examined, $[Cp*RhCl_2]_2$ proved to be preeminent for this reaction (entries 5, 12–16). When the amount of $[Cp*RhCl_2]_2$ was reduced to half from 3.0 mol%, the yield decreased (entries 5 and 17). No reaction took place in the absence of the metal catalyst (entry 18). The reaction was slow at 70 °C and the coupling product was produced in low yield (entry 19).

Table 2 shows the effect of various oxidants on this reaction and Cu(OAc)₂ was found to be an excellent oxidant (entry 6). The use of AgOAc gave products in low yield (entry 1) and other oxidants such as Ag₂CO₃, AgF, CuCl₂, CuCO₃, $(NH_4)_2S_2O_8$, TBHP (*tert*-butyl hydroperoxide) and K₂S₂O₈ were ineffective (entries 2–5 and 10–12). In the absence of an oxidant, no reaction occurred (entry 13). The best yield was obtained when 3.0 mol% amount of [Cp*RhCl₂]₂ was employed in combination with Cu(OAc)₂ (2.4 equiv.) as the oxidant and PivOH (0.5 mmol) as the additive in mesitylene (1.0 mL) at 140 °C for 24 h (entry 6). This reaction proceeded **Organic & Biomolecular Chemistry**



^{*a*} Reaction conditions: **1a** (0.25 mmol), $[Cp*RhCl_2]_2$ (3.0 mol%), oxidants (0.6 mmol), PivOH (0.5 mmol), mesitylene (1.0 mL), 24 h, 140 °C. ^{*b*} Isolated yields. ^{*c*} $[Cp*RhCl_2]_2$ (5.0 mol%), Cu(OAc)₂·H₂O (0.3 mmol) and AgSbF₆ (20 mol%) in mesitylene (1.0 mL) without using PivOH. ^{*d*} Cu(OAc)₂ (0.3 mmol). ^{*e*} Cu(OAc)₂ (0.45 mmol). ^{*f*} 2.0 equiv.

similarly in the absence of PivOH to give a comparative yield of product **2a** (entry 7).

To explore the scope of this rhodium-catalyzed intramolecular oxidative cross-coupling, the reactions of 1a-n were examined under optimized conditions. As shown in Table 3, the corresponding imidazo and benzimidazo[2,1-a]isoquinoline derivatives 2a-l were obtained in moderate to high yields. Alkyl and methoxy substituents exerted little effect on the yield, but substituents at the ortho position of the benzene ring slightly reduced the yields due to the steric effect (2c and 2j). When 1h carrying a 1,2-diphenylvinyl moiety was employed, the reaction proceeded efficiently to give the corresponding cyclized product 2h in 62% yield, indicating that the phenyl group on the vinylic tether carbon did not affect the reaction.¹⁶ However, 1m, having a thiophene ring, and 1n, having an (E)-styryl group, did not afford coupling products. The procedure using AgSbF₆ and Cu(OAc)₂·H₂O in mesitylene gave slightly better results than the procedure using PivOH as the additive as shown in Table 3.

A plausible catalytic cycle for the reaction is illustrated in Scheme 2. The reaction starts from Cp*Rh(OPiv)₂ or Cp*Rh $(OAc)_2$ which is derived from the ligand exchange between rhodium dimer and the pivalic acid or Cu(OAc)₂. Thus formed $Cp*Rh(OCOR)_2$ is coordinated with a nitrogen lone pair of imidazole to give intermediate II in Path A. Rh then undergoes insertion into the C-H bond at the C2 position assisted by the carboxylate ligand leading to the formation of intermediate III with the loss of carboxylic acid.17 The subsequent C-H bond cleavage by Rh(III) generates a seven-membered rhodacyclic intermediate IV which undergoes reductive elimination to give product 2 and a Rh(I) species, and the latter is oxidized by Cu(II) to complete the catalytic cycle. A possible alternative pathway is shown by Path B, where the catalytic cycle is triggered by the coordination of the nitrogen lone pair of imidazole to Cu(OCOR)₂ affording intermediate V. The insertion of

 Table 3
 Rhodium-catalyzed intramolecular oxidative cross-coupling of various imidazoles^a



^a The same reaction conditions as in Table 2, entry 6. Results obtained under the conditions indicated in Table 2, entry 7 are in parentheses.



Scheme 2 Plausible reaction pathways of rhodium-catalyzed oxidative cross-coupling.

Cu into a C–H bond followed by transmetalation with rhodium then gives intermediate III via VI. 18

Conclusions

We report herein on the rhodium-catalyzed synthesis of imidazo and benzimidazo[2,1-*a*]isoquinolines *via* the intramolecular oxidative cross-coupling reaction through double C–H bond cleavage. This protocol can be applied to the synthesis of various heterocyclic compounds. The scope of the reaction and further applications as well as mechanistic studies of the rhodium-catalyzed C–H activation reactions are currently under investigation.

Acknowledgements

This work was supported by a GCOE and Grant-in-Aid for Scientific Research (S) from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

Notes and references

- 1 *Comprehensive Heterocyclic Chemistry*, ed. A. R. Katritzky and C. W. Rees, Elsevier, New York, 1984, vol. 4, 1–38.
- 2 Z. Jin, Nat. Prod. Rep., 2005, 22, 196.
- 3 (a) R. L. Weinkauf, A. Y. Chen, C. Yu, L. Liu, L. Barrows and E. J. Lavoie, *Bioorg. Med. Chem.*, 1994, 2, 781;
 (b) W. J. Houlihan, P. G. Munder, D. A. Handley, S. H. Cheon and V. A. Parrino, *J. Med. Chem.*, 1995, 38, 234;
 (c) P. Molina, E. Aller, A. Lorenzo, C. Foces and A. L. L. Saiz, *Tetrahedron*, 1996, 52, 13671; (d) Z. X. Lu, N. H. Quazi, L. W. Deady and G. M. Polya, *Biol. Chem. Hoppe-Seyler*, 1996, 377, 373; (e) V. K. Pandey and A. Shukla, *Indian J. Chem., Sect. B: Org. Chem. Incl. Med. Chem.*, 1999, 38, 1381; (f) L. W. Deady, T. Rodemann, G. J. Finlay, B. C. Baguley and W. A. Denny, *Anti-Cancer Drug Des.*, 2000, 15, 339; (g) S. M. Rida, S. A. M. El-Hawash, H. T. Y. Fahmy, A. A. Hazzaa and M. M. M. El-Meligy, *Arch. Pharmacal Res.*, 2006, 29, 826.
- 4 I. Hayakawa, Y. Sugano, T. Agatsuma, H. Furukawa, S. Kurakata and S. Naruto, *PCT Int. Appl.*, WO2002034748, 2002; *Chem. Abstr.*, 2002, **136**, 355236.
- 5 (a) A. R. Katritzky, G. Qiu, Q.-H. Long, H.-Y. He and P. J. Steel, *J. Org. Chem.*, 2000, 65, 9201; (b) G. Dyker, W. Stirner and G. Henkel, *Eur. J. Org. Chem.*, 2000, 1433; (c) L. W. Deady and S. M. Devine, *J. Heterocycl. Chem.*, 2004,

41, 549; (*d*) R.-S. Hou, H.-M. Wang, H.-Y. Huang and L.-C. Chen, *J. Chin. Chem. Soc.*, 2004, **51**, 1417; (*e*) A. D. C. Parenty, Y.-F. Song, C.-J. Richmond and L. Cronin, *Org. Lett.*, 2007, **9**, 2253; (*f*) N. Okamoto, K. Sakurai, M. Ishikura, K. Takeda and R. Yanada, *Tetrahedron Lett.*, 2009, **50**, 4167; (*g*) N. Chernyak and V. Gevorgyan, *Angew. Chem., Int. Ed.*, 2010, **49**, 2743.

- 6 G. Cooper and W. J. Irwin, J. Chem. Soc., Perkin Trans. 1, 1976, 75.
- 7 (a) Q. Cai, Z. Li, J. Wei, L. Fu, C. Ha, D. Pei and K. Ding, Org. Lett., 2010, 12, 1500; (b) S. Nandi, S. Samanta, S. Jana and J. K. Ray, Tetrahedron Lett., 2010, 51, 5294; (c) B. Liu, X.-R. Qin, K.-Z. Li, X.-Y. Li, Q. Guo, J.-B. Lan and J.-S. You, Chem.-Eur. J., 2010, 16, 11836; (d) I. Cerna, R. Pohl, B. Klepetarova and M. Hocek, J. Org. Chem., 2010, 75, 2302; (e) J. Lu and H. Fu, J. Org. Chem., 2011, 76, 4600; (f) Q. Liao, L.-Y. Zhang, S.-T. Li and C.-J. Xi, Org. Lett., 2011, 13, 228.
- 8 For selected reports on palladium-catalyzed C-H activation reactions see: (a) A. Shiotani and H. Itatani, Angew. Chem., Int. Ed. Engl., 1974, 13, 471; (b) B. Akermark, L. Eberson, E. Jonsson and E. Petterson, J. Org. Chem., 1975, 40, 1365; (c) Y. Fujiwara, I. Moritani, S. Danno, R. Asano and S. Teranishi, J. Am. Chem. Soc., 1969, 91, 7166; (d) H. Iida, Y. Yuasa and C. Kibayashi, J. Org. Chem., 1980, 45, 2938; (e) T. Watanabe, S. Ueda, S. Inuki, S. Oishi, N. Fujii and H. Ohno, Chem. Commun., 2007, 4516; (f) B. Liegault, D. Lee, M. P. Huestis, D. R. Stuart and K. Fagnou, J. Org. Chem., 2008, 73, 5022; (g) S. Wurtz, S. Rakshit, J. J. Neumann, T. Droge and F. Glorius, Angew. Chem., Int. Ed., 2008, 47, 7230; (h) H.-J. Knolker, Chem. Lett., 2009, 38, 8; (i) L. Ackermann, R. Jeyachandran, H. K. Potukuchi, P. Novak and L. Buttner, Org. Lett., 2010, 12, 2056; (j) D. G. Pintori and M. F. Greaney, J. Am. Chem. Soc., 2011, 133, 1209; (k) P. Gandeepan, C.-H. Hung and C.-H. Cheng, Chem. Commun., 2012, 48, 9379; (l) G. Meng, H.-Y. Niu, G.-R. Qu, J. S. Fossey, J.-P. Li and H.-M. Guo, Chem. Commun., 2012, 48, 9601; (m) M. Sun, H. Wu, J. Zheng and W. Bao, Adv. Synth. Catal., 2012, 354, 835. For recent reviews, see: (n) S.-L. You and J.-B. Xia, Top. Curr. Chem., 2010, 292, 165; (o) N. Kuhl, M. N. Hopkinson, J. Wencel-Delord and F. Glorius, Angew. Chem., Int. Ed., 2012, 51, 10236.
- 9 For recent reviews on rhodium(III)-catalyzed C-H activation reactions, see: (a) D. A. Colby, R. G. Bergman and J. A. Ellman, *Chem. Rev.*, 2010, **110**, 624; (b) J. Bouffard and K. Itami, *Top. Curr. Chem.*, 2010, **292**, 231; (c) T. Satoh and M. Miura, *Chem.-Eur. J.*, 2010, **16**, 11212; (d) T. Satoh and M. Miura, *Synthesis*, 2010, 3395; (e) D. A. Colby, A.-S. Tsai, R. G. Bergman and J. A. Ellman, *Acc. Chem. Res.*, 2012, **45**, 814.
- 10 (a) T. Matsumoto and H. Toshida, *Chem. Lett.*, 2000, 1064;
 (b) T. Matsumoto, R. A. Periana, D. Taube and H. Yoshida, *J. Catal.*, 2002, 206, 272; (c) K. Ueura, T. Satoh and
 M. Miura, *Org. Lett.*, 2007, 9, 1407; (d) J. Chen, G. Song,
 C.-L. Pan and X. Li, *Org. Lett.*, 2010, 12, 5426; (e) F. Wang,

G. Song and X. Li, Org. Lett., 2010, 12, 5430; (f) A. S. Tsai, M. Brasse, R. G. Bergman and J. A. Ellman, Org. Lett., 2011, 13, 540; (g) F. W. Patureau, T. Besset and F. Glorius, Angew. Chem., Int. Ed., 2011, 50, 1064; (h) S. Rakshit, C. Grohmann, T. Besset and F. Glorius, J. Am. Chem. Soc., 2011, 133, 2350; (i) H. Li, Y. Li, X.-S. Zhang, K. Chen, X. Wang and Z.-J. Shi, J. Am. Chem. Soc., 2011, 133, 15244; (j) S. Mochida, K. Hirano, T. Satoh and M. Miura, J. Org. Chem., 2011, 76, 3024; (k) X. Li and M. Zhao, J. Org. Chem., 2011, 76, 8530; (l) S. H. Park, J. Y. Kim and S. Chang, Org. Lett., 2011, 13, 2372; (m) F. Wang, G. Song, Z. Du and X. Li, J. Org. Chem., 2011, 76, 2926; (n) J. Willwacher, S. Rakshit and F. Glorius, Org. Biomol. Chem., 2011, 9, 4736; (o) S. Mochida, K. Hirano, T. Satoh and M. Miura, J. Org. *Chem.*, 2011, **76**, 3024; (*p*) L. Zheng and J. Wang, Chem.-Eur. J., 2012, 18, 9699; (q) P. Zhao, R. Niu, F. Wang and X. Li, Org. Lett., 2012, 14, 4166; (r) Z. Shi, N. Schrçder and F. Glorius, Angew. Chem., Int. Ed., 2012, 51, 8092.

- 11 (a) K. Morimoto, M. Itoh, K. Hirano, T. Satoh, Y. Shibata, K. Tanaka and M. Miura, Angew. Chem., Int. Ed., 2012, 51, 5359; (b) J. Wencel-Delord, C. Nimphius, F. W. Patureau and F. Glorius, Angew. Chem., Int. Ed., 2012, 51, 2247; (c) P. Wang, H. Rao, R. Hua and C.-J. Li, Org. Lett., 2012, 14, 902; (d) N. Kuhl, M. N. Hopkinson and F. Glorius, Angew. Chem., Int. Ed., 2012, 51, 8230; (e) J. Dong, Z. Long, F. Song, N. Wu, Q. Guo, J. Lan and J. You, Angew. Chem., Int. Ed., 2013, 52, 580; (f) V. P. Reddy, R. Qiu, T. Iwasaki and N. Kambe, Org. Lett., DOI: 10.1021/ol400230y.
- 12 (a) K. Ueura, T. Satoh and M. Miura, J. Org. Chem., 2007, 72, 5362; (b) N. Umeda, H. Tsurugi, T. Satoh and M. Miura, Angew. Chem., Int. Ed., 2008, 47, 4019; (c) T. Fukutani, N. Umeda, K. Hirano, T. Satoh and M. Miura, Chem. Commun., 2009, 5141; (d) M. Shimizu, K. Hirano, T. Satoh and M. Miura, J. Org. Chem., 2009, 74, 3478; (e) T. K. Hyster and T. Rovis, J. Am. Chem. Soc., 2010, 132, 10565; (f) P. C. Too, Y.-F. Wang and S. Chiba, Org. Lett., 2010, 12, 5688; (g) K. Morimoto, K. Hirano, T. Satoh and M. Miura, Org. Lett., 2010, 12, 2068; (h) N. Guimond, S. I. Gorelsky and K. Fagnou, J. Am. Chem. Soc., 2011, 133, 6449; (i) T. Fukutani, K. Hirano, T. Satoh and M. Miura, J. Org. Chem., 2011, 76, 2867; (j) N. Umeda, K. Hirano, T. Satoh, N. Shibata, H. Sato and M. Miura, J. Org. Chem., 2011, 76, 13; (k) F. W. Patureau, T. Besset, N. Kuhl and F. Glorius, J. Am. Chem. Soc., 2011, 133, 2154; (l) K. Morimoto, K. Hirano, T. Satoh and M. Miura, J. Org. Chem., 2011, 76, 9548; (m) H. Wang and F. Glorius, Angew. Chem., Int. Ed., 2012, 51, 7318.
- 13 (a) K. D. Hesp, R. G. Bergman and J. A. Ellman, J. Am. Chem. Soc., 2011, 133, 11430; (b) Y. Li, B.-J. Li, W.-H. Wang, W.-P. Huang, X.-S. Zhang, K. Chen and Z.-J. Shi, Angew. Chem., Int. Ed., 2011, 50, 2115; (c) L. Yang, C. A. Correia and C.-J. Li, Adv. Synth. Catal., 2011, 353, 1269.
- 14 (*a*) J. Terao, A. Oda and N. Kambe, *Org. Lett.*, 2004, 6, 3341;
 (*b*) Y. Minami, H. Kuniyasu and N. Kambe, *Org. Lett.*, 2008,
 10, 2469; (*c*) M. Toyofuku, S.-I. Fujiwara, T. Shin-ike,

H. Kuniyasu and N. Kambe, *J. Am. Chem. Soc.*, 2005, **127**, 9706; (*d*) M. Toyofuku, E. Murase, S.-I. Fujiwara, T. Shin-ike, H. Kuniyasu and N. Kambe, *Org. Lett.*, 2008, **10**, 3957; (*e*) Y. Minami, H. Kuniyasu, A. Sanagawa and N. Kambe, *Org. Lett.*, 2010, **12**, 3744; (*f*) N. Kambe, T. Iwasaki and J. Terao, *Chem. Soc. Rev.*, 2011, **40**, 4937.

15 Recently, palladium-catalyzed intermolecular oxidative cross-coupling of imidazole and benzimidazole derivatives

has been reported: ref. 8*m*. For Rh-catalyzed intramolecular aryl–aryl coupling, see: ref. 11*a*,*c*.

- 16 In ref. 8m, it was claimed that introduction of a Me substituent on an olefinic tether carbon resulted in lower yield (21%) in the case of the Pd/Cu catalytic system.
- 17 D. Lapointe and K. Fagnou, Chem. Lett., 2010, 39, 1118.
- 18 J.-R. Huang, L. Dong, B. Han, C. Peng and Y.-C. Chen, *Chem.-Eur. J.*, 2012, 18, 8896.