A selective fluorescent chemosensor for phosphoserine†

Chad M. Cooley, Kenneth S. Hettie, Jessica L. Klockow, Shana Garrison and Timothy E. Glass*

Cite this: *Org. Biomol. Chem.*, 2013, **11**, 7387Received 16th August 2013,
Accepted 18th September 2013

DOI: 10.1039/c3ob41677a

www.rsc.org/obc

A fluorescent chemosensor for the detection of phosphoserine is reported. The ditopic sensor features a phosphate-coordinating zinc(II)-dipicolylamine (Zn^{2+} -DPA) unit tethered to an amine-binding coumarin aldehyde fluorophore. With phosphoserine, the sensor demonstrates a 30-fold fluorescence enhancement under buffered aqueous conditions.

Introduction

Phospholipids constitute the dominant class of lipids composing the lipid bilayer of cellular membranes and confer structural support, organization of membrane proteins, and intercellular signaling mechanisms.^{1–6} Phosphatidylcholine (PC), sphingomyelin (SM), phosphatidylethanolamine (PE), and phosphatidylserine (PS) represent over 97% of the total phospholipid composition and are asymmetrically distributed between the inner and outer leaflets of a cellular membrane.⁷ In quiescent cells, PC and SM constitute ~87% of the total phospholipid content of the outer leaflet, with PS being exclusively restricted to and comprising ~30% of the total phospholipid content of the inner leaflet (Fig. 1). Cell death randomizes the asymmetric distribution of phospholipids between the inner and outer leaflets, thereby externalizing PS to the outer cell surface.^{8–10} In fact, exposure of PS to the outer leaflet of the cellular membrane is the hallmark molecular biomarker for all types of cell death and serves as a signal for phagocytosis by providing $\sim 10^6$ – 10^9 binding sites per cell.¹¹

Several methods to tag apoptotic cells and phospho-amino acid residues have appeared in the literature.^{12–17} In addition to biological probes (e.g., annexin assays), the vast majority of synthetic approaches have focused on detecting anionic phospholipid head groups, which includes phosphoserine. In particular, zinc(II)-dipicolylamine (Zn^{2+} -DPA) complexes have proven to be admirable receptors for negatively-charged lipid head groups and have been used to tag apoptotic cells as well as bacterial cells.^{11,12,14,15,18,19} These primarily consist of

Fig. 1 (a) Common cellular membrane phospholipid head groups. PC = phosphatidylcholine, SM = sphingomyelin, PS = phosphatidylserine, PE = phosphatidylethanolamine, PG = phosphatidylglycerol, PGP = phosphatidylglucopyranose. (b) Asymmetric phospholipid composition (mol%) of the outer and inner leaflets of healthy human cellular membranes.⁷

two pre-coordinated Zn^{2+} -DPA units tethered to a synthetic fluorophore that associate with the membrane surface of apoptotic cells through strong multipoint interactions with the anionic phospholipid head groups. These molecular probes are phospholipid tags and do not operate in a turn-on fashion. By contrast, a turn-on sensor for PS could give quantitative information about the amount of lipid present with low background as well as enable new types of experiments such as the ability to monitor the progression of cell death in a cell population.

Recently, we have been exploring the use of water-soluble shape-selective molecular tubes for hydrophobic sensing of lipids.²⁰ In order to build selective sensors for PS, we require a

Department of Chemistry, University of Missouri, 601 South College Avenue, Columbia, MO 65211, USA. E-mail: GlassT@missouri.edu; Fax: +1 573-882-2754; Tel: +1 573-882-3813

†Electronic supplementary information (ESI) available: UV/Vis and fluorescence spectroscopic results and ^1H and ^{13}C NMR spectra. See DOI: 10.1039/c3ob41677a

We reasoned that only phospholipid head groups with primary amines should bind to the coumarin aldehyde and form an iminium ion which causes an ~ 40 nm bathochromic

Binding titrations. Sensors **1** and **2** were titrated with several phospholipid head groups: phosphoserine, phosphoethanolamine, phosphocholine, α -glycerolphosphate, and D-glucopyranose 1-phosphate, as well as the primary amines glutamate and glycine under buffered aqueous conditions (50 mM HEPES, 100 mM NaCl, pH 7.4). Glutamate and glycine served as controls for amine binding as they lack phosphate groups. The binding and spectroscopic data are summarized in Table 1.

Table 1 Binding constants (K_a) and spectroscopic properties of sensors **1** and **2** with various analytes^a

Analyte	1		1-Zn²⁺ ^b		2		2-Zn²⁺ ^b	
	K_a (M ⁻¹)	I_{sat}/I_0 ^c	K_a (M ⁻¹)	I_{sat}/I_0 ^c	K_a (M ⁻¹)	I_{sat}/I_0 ^c	K_a (M ⁻¹)	I_{sat}/I_0 ^c
Phosphoserine	3.1	7.2	5.4	6.2	120	5.8	310	30
Phosphoethanolamine	5.7	4.8	5.7	4.9	93	9.1	180	39
Glutamate	4.7	7.0	7.0	4.3	22	13	190	17
Glycine	6.1	6.1	6.2	4.0	5.3	16	6.6	29

^a Analytes of various concentrations were added to sensor (10 μM) in buffer (50 mM HEPES, 100 mM NaCl, pH 7.4). Sensor **1** was titrated in 97 : 3 buffer–DMSO and sensor **2** was titrated in 100% buffer. ^b Zinc acetate (10 μM) was pre-coordinated with sensor prior to addition of analyte. ^c I_{sat} was taken from the theoretical maximum of the binding isotherm; λ_{ex} = 488 nm; $\lambda_{\text{em}, 1}$ = 521 nm, $\lambda_{\text{em}, 2}$ = 513 nm. The phospholipid head groups phosphocholine, α -glycerolphosphate, and D-glucose-1-phosphate did not bind or alter the fluorescence properties of either sensor.

The absorbance of **1-Zn²⁺** shifted to 488 nm upon the addition of phosphoserine, phosphoethanolamine, glutamate, and glycine (see ESI†).

Phosphocholine, α -glycerolphosphate, and D-glucose-1-phosphate did not bind to the coumarin aldehyde of **1-Zn²⁺** and therefore did not alter the spectroscopic properties. All primary amine analytes bound to the coumarin aldehyde with low affinity and relatively low fluorescence enhancements (<7-fold) with emission maxima at 521 nm when excited at 488 nm. The binding results were similar to the control experiments conducted without Zn^{2+} indicating that the Zn^{2+} –DPA had little effect on enhancing analyte binding to the coumarin aldehyde (see ESI†). Indeed, these binding constants are similar to those achieved with simple, unsubstituted coumarin aldehydes.^{21–23}

Sensor **2** displayed similar absorption properties compared to sensor **1**. The absorbance of **2-Zn²⁺** also shifted to the red only upon addition of phosphoserine, phosphoethanolamine, glutamate, and glycine. As seen with sensor **1**, the phospholipid head groups that did not contain primary amines did not bind. The binding for phosphoserine was remarkably higher than for the other analytes tested with $K_a = 310 \text{ M}^{-1}$ (Fig. 3). Titrations with phosphoethanolamine and glutamate also demonstrated reasonable affinity, though not as high as with phosphoserine. The fluorescence increased over 30-fold with the addition of both phosphoserine and phosphoethanolamine, with emission at 513 nm when excited at 488 nm.

For sensor **2**, the spectroscopic properties with Zn^{2+} were much more favorable than the control experiments without Zn^{2+} , though the binding constants of **2** with phosphoserine, phosphoethanolamine, and glutamate were elevated compared to the typical low binding constants seen with monotopic coumarin aldehyde sensors.^{21–23} Presumably, the protonated form of the DPA can interact favorably with the charged phosphate to enhance binding. The fluorescence response of sensor **2** was lower without Zn^{2+} due to photoinduced electron transfer quenching from the uncoordinated DPA.

It is clear from the spectroscopic studies of both sensors that **2-Zn²⁺** is the best sensor for phosphoserine both in terms of binding constant and maximum fluorescence response (Fig. 4). Because of the flexible linker, **2-Zn²⁺** demonstrated more than two orders of magnitude better binding than the

Fig. 3 (a) UV/Vis and (b) fluorescence spectra of **2-Zn²⁺** (10 μM) with aliquots of 100 mM phosphoserine in buffer (50 mM HEPES, 100 mM NaCl, pH 7.4). λ_{ex} = 488 nm. λ_{em} = 513 nm. Inset is the fit to a binding isotherm.

more rigid sensor **1**. Clearly, the rigid structure for **1-Zn²⁺** prohibits a two-point binding interaction. Even sensor **2** lacking Zn^{2+} was a better receptor for phosphoserine than sensor **1**.

Compared to the phospholipid tagging probes, **2-Zn²⁺** is unique in that it can coordinate to the analyte phosphate group and primary amine to achieve selective binding to phosphoserine over other phospholipid head groups. Thus, the head groups phosphocholine, α -glycerolphosphate, and D-glucose-1-phosphate give no response, which is extremely important since the majority of membrane phospholipids are choline-based. In addition, **2-Zn²⁺** is also sensitive to

Fig. 4 (a) Binding constants and (b) fluorescence enhancements (I_{sat}/I_0) of sensors 1 and 2 with phosphoserine.

the charge on the head group. Phosphoethanolamine and glutamate bind less effectively, as each bears one less negative charge than phosphoserine. It should be noted that while glutamate and glycine gave enhanced binding and fluorescence output, they are not expressed at high concentrations on a cellular membrane.

Conclusions

In conclusion, 2-Zn²⁺ demonstrates a means to selectively sense phosphoserine over other major phospholipid head groups by forming a two-point binding interaction with the phosphate and primary amine functional groups of the analyte under buffered aqueous conditions. Future work in this area will focus on joining this head group binding unit to our lipid receptor to produce sensitive and selective sensors for bio-active lipids.

Experimental section

Spectroscopic analysis

Instrumentation. Absorption spectra were recorded on a Cary 1E spectrophotometer at 25 °C. Fluorescence spectra were recorded on a Shimadzu RF-5301 PC spectrofluorometer at 25 °C. NMR spectra were recorded on Bruker DRX 500.

Solution preparation. Solutions for UV/Vis and fluorescence spectroscopy were prepared from a 1 mg mL⁻¹ sensor stock solution in MeOH that was diluted to volume with buffer (50 mM HEPES, 100 mM NaCl, pH 7.4) until a 10 μM sensor

solution was produced following removal of the solvent. For sensor 1, DMSO was added to equal 3% of the total volume to improve solubility. Analytes were measured and diluted to volume with the buffered sensor solution to keep the sensor concentration constant throughout the titration.

Quantum yields. Separate stock solutions of sensors 1 and 2 (10 μM) in buffer (50 mM HEPES, 100 mM NaCl, pH 7.4) were prepared. The solution of sensor 1 contained 3% (v/v) DMSO as a cosolvent. For quantum yields (Φ_f), a 1.0 μM solution of fluorescein (absolute $\Phi_f = 0.85$) in 0.1 N NaOH (pH 13) was used as a fluorescence standard. Sensors 1 and 2 were excited at 473 nm. The slit width was 2 nm for both excitation and emission. Quantum yields were obtained in triplicate and calculated as per the following equation:

$$\Phi_S = \Phi_R \times \frac{I_S}{I_R} \times \frac{A_R}{A_S} \times \frac{n_S^2}{n_R^2} \quad (1)$$

where Φ_S = quantum yield of the sample, Φ_R = quantum yield of the reference, I_S = integrated fluorescence intensity of the sample, I_R = integrated fluorescence intensity of the reference, A_R = absorbance of the reference, A_S = absorbance of the sample, n_S = refractive index of the sample solvent, n_R = refractive index of the reference solvent.

Binding constant calculations. The concentration of analyte was plotted against the fluorescence intensity and fit to a one-site binding isotherm using the following equation:

$$I/I_0 - 1 = \frac{B_{max}[\text{analyte}]}{K_d + [\text{analyte}]} \quad (2)$$

where I = fluorescence intensity, I_0 = initial fluorescence intensity, B_{max} = theoretical maximum fluorescence, and K_d = dissociation constant.

Synthetic procedures

General methods. All reactions were carried out in dried glassware under a blanket of N₂. Before use, tetrahydrofuran and diethyl ether were distilled from sodium benzophenone ketyl while dichloromethane and triethylamine were distilled from CaH₂. NMR spectra were recorded on Bruker DRX 500. IR spectra were recorded on a Nexus 670 FT-IR E.S.P. spectrometer.

Materials. All chemicals were obtained from Sigma Aldrich, Fisher Scientific, Combi-Blocks, TCI America, or Alfa Aesar and were used without further purification. Flash chromatography was performed with 32–63 μm silica gel.

Compound 4. Compound 3 (250 mg, 1.072 mmol), Na₂CO₃ (340 mg, 3.215 mmol), and TsCl (204.4 mg, 1.072 mmol) were dissolved in 10 mL degassed DMF-H₂O (95 : 5). The solution stirred at 60 °C for 1 h. Then 4-(hydroxymethyl)phenylboronic acid (179.2 mg, 1.179 mmol) and PdCl₂ (29 mg, 0.161 mmol) were added and N₂ was bubbled through for 5 min. The solution stirred at 60 °C for 16 h. The inorganic solids were filtered off and rinsed with acetone. The solvent was removed *in vacuo* and the remaining crude material was purified by chromatography (95 : 5 CH₂Cl₂-EtOAc) to yield compound 4 (156.5 mg,

45%) as a yellow solid (mp 62 °C): ^1H NMR (500 MHz, CDCl_3) δ 7.50 (d, 2H, $J = 7.5$ Hz), 7.43 (d, 2H, $J = 7.5$ Hz), 7.23 (d, 1H, $J = 9.0$ Hz), 6.57 (d, 1H, $J = 2.0$ Hz), 6.51 (dd, 1H, $J = 9.0$, 2.0 Hz), 5.99 (s, 2H), 4.80 (s, 2H), 3.42 (q, 4H, $J = 7.0$), 1.21 (t, 6H, 7.0 Hz); ^{13}C NMR (125 MHz, CDCl_3) δ 162.1, 156.8, 155.8, 150.6, 142.1, 135.5, 128.6, 127.8, 127.1, 108.5, 108.2, 107.9, 97.9, 64.8, 44.8, 12.4; IR (neat, cm^{-1}) 3412, 2970, 1695, 1605, 1589, 1417, 1352, 1115; HRMS calculated for $\text{C}_{20}\text{H}_{21}\text{NO}_3\text{Na}$ ($\text{M} + \text{Na}^+$): 346.1414. Found: 346.1414.

Compound 5. Compound 4 (100 mg, 0.309 mmol) and POCl_3 (3 mL) were combined in a round bottom flask and stirred at room temperature for 21 h. The mixture was poured over ice water (10 mL) and extracted with CH_2Cl_2 (10 mL $\times$ 4). The organic solvent was removed *in vacuo* and the resulting material was purified by chromatography (90:10 CH_2Cl_2 -EtOAc) to yield compound 5 (81 mg, 77%) as a yellow oil: ^1H NMR (500 MHz, CDCl_3) δ 7.51 (d, 2H, $J = 8.0$ Hz), 7.43 (d, 2H, $J = 8.0$ Hz), 7.22 (d, 1H, $J = 9.0$ Hz), 6.56 (d, 1H, $J = 2.5$ Hz), 6.51 (dd, 1H, $J = 9.0$, 2.5 Hz), 5.98 (s, 1H), 3.41 (q, 4H, $J = 7.0$ Hz), 1.20 (t, 6H, $J = 7.0$ Hz); ^{13}C NMR (125 MHz, CDCl_3) δ 161.9, 156.7, 155.4, 150.6, 138.5, 136.3, 128.8, 128.7, 127.7, 108.5, 108.3, 107.6, 97.8, 45.6, 44.7, 12.4; IR (neat, cm^{-1}) 2970, 1708, 1614, 1593, 1524, 1417, 1356, 1115; HRMS calculated for $\text{C}_{20}\text{H}_{20}\text{ClNO}_2\text{Na}$ ($\text{M} + \text{Na}^+$): 364.1075. Found: 364.1077.

Compound 6. The Vilsmeier reagent was made by combining 2 mL POCl_3 and 3.6 mL DMF in a flame-dried round bottom flask at 0 °C and stirring for 30 min. In a separate flask, 142.7 mg of compound 5 was dissolved in 3 mL DMF. The starting material was cooled to 0 °C and 3 mL of the Vilsmeier reagent was added dropwise. The reaction was brought to room temperature and stirred for 40 h. The mixture was poured over ice water (30 mL), extracted with CH_2Cl_2 (20 mL $\times$ 6), the organic layers combined and dried over Na_2SO_4 , and the solvent removed *in vacuo*. Purification *via* column chromatography (95:5 CH_2Cl_2 -EtOAc) yielded compound 6 as a yellow oil (97 mg, 62%): ^1H NMR (500 MHz, CDCl_3) δ 9.89 (s, 1H), 7.54 (d, 2H, $J = 8.0$ Hz), 7.26 (d, 2H, $J = 8.0$ Hz), 6.94 (d, 1H, $J = 9.5$ Hz), 6.49–6.53 (m, 2H), 4.68 (s, 2H), 3.45 (q, 4H, $J = 7.0$ Hz), 1.23 (t, 6H, $J = 7.0$ Hz); ^{13}C NMR (125 MHz, CDCl_3) δ 188.2, 160.8, 160.1, 157.7, 153.1, 138.3, 133.4, 131.0, 128.6, 128.5, 112.0, 109.8, 108.9, 97.0, 45.7, 45.2, 12.4; IR (neat, cm^{-1}) 1614, 1559, 1494, 1417, 1352; HRMS calculated for $\text{C}_{21}\text{H}_{20}\text{ClNO}_3\text{Na}$ ($\text{M} + \text{Na}^+$): 392.1024. Found: 392.1022.

Sensor 1. Compound 6 (102 mg, 0.276 mmol), di(2-picolyl)-amine (55 μL , 0.290 mmol), K_2CO_3 (114 mg, 0.828 mmol), and 7 mL THF were combined in a round bottom flask at stirred at 55 °C for 21 h. The mixture was filtered and rinsed with acetone. The filtrate solvent was removed *in vacuo* and the remaining residue was purified *via* column chromatography (100% $\text{CH}_2\text{Cl}_2 \rightarrow 100\%$ EtOAc $\rightarrow 80:20$ EtOAc-MeOH) to yield sensor 1 as a yellow oil (54 mg, 47%). ^1H NMR (500 MHz, CDCl_3) δ 9.76 (s, 1H) 8.55 (d, 2H, $J = 5.0$ Hz), 7.69 (td, 2H, $J = 7.5$, 1.5 Hz), 7.61 (d, 2H, $J = 8.0$ Hz), 7.54 (d, 2H, $J = 8.0$ Hz), 7.21 (d, 2H, $J = 8.0$ Hz), 7.17 (m, 2H), 6.93 (d, 1H, $J = 9.5$ Hz), 6.45–6.51 (m, 2H), 3.88 (s, 4H), 3.80 (s, 2H), 3.43 (q, 4H, $J =$

7.0 Hz), 1.21 (t, 6H, $J = 7.0$ Hz); ^{13}C NMR (125 MHz, CDCl_3) δ 188.3, 162.2, 159.5, 159.4, 157.7, 153.0, 148.9, 140.2, 136.5, 131.5, 130.9, 128.7, 128.5, 122.8, 122.0, 112.2, 109.6, 108.9, 96.9, 60.1, 58.1, 45.1, 12.4; IR (neat, cm^{-1}) 1746, 1611, 1495, 1422, 1356, 1133, 731; HRMS calculated for $\text{C}_{33}\text{H}_{32}\text{N}_4\text{O}_3\text{Na}$ ($\text{M} + \text{Na}^+$): 555.2367. Found: 555.2366.

Compound 8. Compound 7 (0.0412 g, 0.1358 mmol), iodine (0.0655 g, 0.2580 mmol), triphenylphosphine (0.0641 g, 0.2444 mmol) and imidazole (0.0185 g, 0.2716 mmol) were dissolved in dichloromethane (7.0 mL) and stirred at room temperature for 2 h. The mixture was filtered and columned twice on silica with 9:1 CH_2Cl_2 -EtOAc to give compound 8 (48.9 mg, 90%) as a yellow solid (decomp 138–141 °C). ^1H NMR (500 MHz, CDCl_3) δ 10.35 (s, 1H), 7.73 (d, 1H, $J = 16.0$ Hz), 6.69 (dd, 1H, $J = 15.5$, 4.5 Hz), 6.47 (d, 1H, $J = 4.5$ Hz), 3.33–3.52 (m, 8H), 2.05–2.15 (m, 2H), 1.26 (t, 6H, $J = 7.0$ Hz); ^{13}C NMR (125 MHz, CDCl_3) δ 190.9, 162.9, 161.5, 157.6, 152.8, 128.6, 111.8, 110.1, 108.4, 97.3, 45.1, 33.5, 28.9, 12.5, 6.8; IR (neat, cm^{-1}) 2913, 1716, 1671, 1614, 1556, 1503, 1454, 1352; HRMS calculated for $\text{C}_{17}\text{H}_{20}\text{INO}_3\text{Na}$ ($\text{M} + \text{Na}^+$): 436.0380. Found: 436.0372.

Sensor 2. In a flame-dried flask, compound 8 (0.0485 g, 0.1174 mmol), potassium carbonate (0.0243 g, 0.1761 mmol), and di(2-picolyl)amine (31.7 μL , 0.1761 mmol) were dissolved in acetonitrile (4.1 mL) and the flask was flushed with N_2 . The mixture was stirred at room temperature for 60 h. Column chromatography on silica with 1:1 CH_2Cl_2 -EtOAc gave sensor 2 (24.1 mg, 56%) as a yellow solid (mp 109 °C). ^1H NMR (500 MHz, CDCl_3) δ 10.34 (s, 1H), 8.55 (d, 2H, $J = 7.0$ Hz), 7.56–7.71 (m, 5H), 7.14–7.19 (m, 2H), 6.52 (dd, 1H, $J = 9.5$, 4.5 Hz), 6.44 (d, 1H, $J = 4.5$ Hz); ^{13}C NMR (125 MHz, CDCl_3) δ 190.7, 163.8, 163.1, 159.6, 157.5, 152.6, 149.0, 136.4, 128.7, 123.0, 121.9, 111.7, 109.8, 108.5, 97.2, 60.2, 54.3, 45.0, 27.8, 25.8, 12.4; IR (neat, cm^{-1}) 2970, 2921, 1712, 1676, 1615, 1559, 1508, 1452, 1355, 1148, 1131; HRMS calculated for $\text{C}_{29}\text{H}_{32}\text{N}_4\text{O}_3\text{Na}$ ($\text{M} + \text{Na}^+$): 507.2367. Found: 507.2357.

Acknowledgements

This work was supported by the National Science Foundation (CHE-1112194).

Notes and references

- 1 M. I. Gurr, J. L. Harwood and K. N. Frayn, *Lipid Biochemistry: An Introduction*, Oxford, UK, 5th edn, 2002, pp. 215–259.
- 2 R. O. Ryan and D. J. van der Horst, *Annu. Rev. Entomol.*, 2000, **45**, 233.
- 3 M. Dykstra, A. Cherukuri, H. W. Sohn, S.-J. Tzeng and S. K. Pierce, *Annu. Rev. Immunol.*, 2003, **21**, 457.
- 4 M. Kielian, P. K. Chatterjee, D. L. Gibbons and Y. E. Lu, *Subcell. Biochem.*, 2000, **34**, 409.

- 5 K. Lauber, E. Bohn, S. M. Kröber, Y.-J. Xiao, S. G. Blumenthal, R. K. Lindemann, P. Marini, C. Wiedig, A. Zobywalski, S. Baksh, Y. Xu, I. B. Autenrieth, K. Schulze-Osthoff, C. Belka, G. Stuhler and S. Wesselborg, *Cell*, 2003, **113**, 717.
- 6 S. A. Islam, S. Y. Thomas, C. Hess, B. D. Medoff, T. K. Means, C. Brander, C. M. Lilly, A. M. Tager and A. D. Luster, *Blood*, 2006, **107**, 444.
- 7 J. A. Virtanen, K. H. Cheng and P. Somerharju, *Proc. Natl. Acad. Sci. U. S. A.*, 1998, **95**, 4964.
- 8 Y. Y. Tyurina, A. A. Shvedova, K. Kawai, V. A. Tyurin, C. Kommineni, P. J. Quinn, N. F. Schor, J. P. Fabisiak and V. E. Kagan, *Toxicology*, 2000, **148**, 93.
- 9 G. Preta and B. Fadeel, *PLoS One*, 2012, **7**, 1.
- 10 T. T. Tung, K. Nagaosa, Y. Fujita, A. Kita, H. Mori, R. Okada, S. Nonaka and Y. Nakanishi, *J. Biochem.*, 2013, **153**, 483.
- 11 B. A. Smith, S. T. Gammon, S. Xiao, W. Wang, S. Chapman, R. McDermott, M. A. Suckow, J. R. Johnson, D. Piwnicka-Worms, G. W. Gokel, B. D. Smith and W. M. Leevy, *Mol. Pharmaceutics*, 2011, **8**, 583.
- 12 B. A. Smith and B. D. Smith, *Bioconjugate Chem.*, 2012, **23**, 1989.
- 13 B. A. Smith, S. Xiao, W. Wolter, J. Wheeler, M. A. Suckow and B. D. Smith, *Apoptosis*, 2011, **16**, 722.
- 14 R. G. Hanshaw and B. D. Smith, *Bioorg. Med. Chem.*, 2005, **13**, 5035.
- 15 S. W. Bae, M. S. Cho, A. R. Jeong, B.-R. Choi, D.-E. Kim, W.-S. Yeo and J.-I. Hong, *Small*, 2010, **6**, 1499.
- 16 P. Atkinson, B. S. Murray and D. Parker, *Org. Biomol. Chem.*, 2006, **4**, 3166.
- 17 T. Anai, E. Nakata, Y. Koshi, A. Ojida and I. Hamachi, *J. Am. Chem. Soc.*, 2007, **129**, 6232.
- 18 A. G. White, B. D. Gray, K. Y. Pak and B. D. Smith, *Bioorg. Med. Chem. Lett.*, 2012, **22**, 2833.
- 19 C. Lakshmi, R. G. Hanshaw and B. D. Smith, *Tetrahedron*, 2004, **60**, 11307.
- 20 C. T. Avetta, B. J. Shorthill, C. Ren and T. E. Glass, *J. Org. Chem.*, 2012, **77**, 851.
- 21 K. S. Hettie, X. Liu, K. D. Gillis and T. E. Glass, *ACS Chem. Neurosci.*, 2013, **4**, 918.
- 22 K. E. Secor and T. E. Glass, *Org. Lett.*, 2004, **6**, 3727.
- 23 E. K. Feuster and T. E. Glass, *J. Am. Chem. Soc.*, 2003, **125**, 16174.
- 24 C. A. Parker and W. T. Rees, *Analyst*, 1960, **85**, 587.