Chemical tweaking of a non-fluorescent GFP chromophore to a highly fluorescent coumarinic fluorophore: application towards photo-uncaging and stem cell imaging†

Cite this: *RSC Adv.*, 2013, **3**, 24021Received 31st July 2013
Accepted 10th October 2013

DOI: 10.1039/c3ra44034f

www.rsc.org/advances

Tanmay Chatterjee, Debjit Roy, Ananya Das, Anup Ghosh, Partha Pratim Bag and Prasun K. Mandal*

Three step chemical tweaking of nonfluorescent parent GFP chromophore has yielded a three hundred times as bright coumarinic fluorophore. A fluorogenic compound has been prepared which after photo-uncaging yields the same highly bright fluorophore that could be used for live stem cell imaging.

The quest to enhance the fluorescence quantum yield of nonfluorescent GFP (green fluorescent protein)^{1,2} chromophores (*p*-HBDI, see Scheme 1) has engaged many researchers from biology and chemistry for more than a decade. Strong deviation of fluorescence quantum yield (ϕ) of *p*-HBDI ($\phi < 0.001$) from that of wild type GFP ($\phi \sim 0.8$) has been noted in solution.^{3,4} Different mechanisms have been put forward as plausible reasons for the low fluorescence quantum yield of GFP chromophore.^{5,6} Several successful attempts^{7,8} to increase ϕ of GFP chromophore by restricting the torsional motion include mimicking GFP barrel encapsulation in human serum albumin, in octaacid cavitand, and using truncated or split GFP technique, and selective complexation with RNA aptamers, complexation with metal salts (such as Zn^{2+}) or BF_2 moiety. However, concerns have been raised regarding their usage due to lability of metal- BF_2 complexes in solution.⁹ Different chemical modifications/substitutions in benzene and imidazole ring of *p*-HBDI have been tried but so far a maximum of only $\sim 10\%$ fluorescence quantum yield could be achieved in GFP chromophore analogues.¹⁰ Thus, the quest for chemically modified GFP chromophore analogue with enhanced fluorescence quantum yield is still very much active.

Photoactivation process provides spatial and temporal control over the release of desired chemical¹¹ and thus photo-activable compounds and fluorescent proteins are quite

important towards bioimaging.¹² There are quite a few number of photoactivable compounds, however, there is only one report so far where a photoactivable compound has been made with GFP chromophore analogue.¹³

In this manuscript we describe how three step chemical tweaking of non-fluorescent parent GFP chromophore (*p*-HBDI) could lead to a highly fluorescent coumarinic fluorophore (Scheme 1). Three modifications that has been incorporated are (i) *p*-hydroxy has been replaced by *o*-hydroxy, (ii) *N,N*-diethyl-amino group has been introduced at the *p*-position, and (iii) sp^3 nitrogen in imidazole ring has been replaced by oxygen. In this direction we have made four compounds: 4-(2-hydroxy-4-*N,N*-diethylamino-benzylidene)-1,2-dimethyl-1*H*-imidazol-5(4*H*)-one (OHIM), 4-(2-methoxy-4-*N,N*-diethylamino-benzylidene)-1,

Scheme 1 (a) Three step chemical tweaking (in red) of *p*-HBDI. (b) Chemical structure and synthesis of all four dyes. (c) Possible pathway for conversion of OHBO to cOHBO.

Department of Chemical Sciences, Indian Institute of Science Education and Research (IISER)-Kolkata, Mohanpur Campus, West-Bengal, 741252, India. E-mail: prasunchem@iiserkol.ac.in

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c3ra44034f

Table 1 Photophysical properties of OHIM and cOHBO

Dye	Solvent	λ_{max}^a (nm)	ϵ (M ⁻¹ cm ⁻¹)	ϕ	τ
OHIM	Toluene	484(448)	29 075	2.2×10^{-3}	0.76, 32.8 ps
	Methanol	505(451)	30 580	1.4×10^{-3}	0.73, 13.9 ps
cOHBO	Toluene	444(377)	19 030	0.90	3.27 ns
	Methanol	485(392)	19 000	0.70	3.88 ns

^a Emission maximum, absorption maximum (nm) in parentheses.

2-dimethyl-1*H*-imidazol-5(4*H*)-one (OMIM) and two of their respective point atom mutation analogues (OHBO and OMBO respectively) where the nitrogen atom at 1-position have been replaced by oxygen atom (see Scheme 1b) (for synthesis and characterisation see ESI†). Absorption and emission spectra of OHIM and OHBO in different solvents are shown in ESI† and the photophysical parameters are depicted in Table 1. (For OMIM and OMBO see ESI†).

Surprisingly, we have noticed that ϕ as well as the fluorescence lifetime (τ) of OHBO are drastically different from that of OHIM (or other two derivatives). To the best of our knowledge this is the first example of chemical modification of GFP chromophore leading to a highly bright fluorescent dye with near unity fluorescence quantum yield, *i.e.* more than 300 fold fluorescence enhancement from the GFP chromophore in solution. It was observed that the fluorescence lifetime of OHIM, OMIM, and OMBO are in femtosecond or picosecond time scale, whereas fluorescence lifetime of OHBO is in nanosecond time scale. Thus an enhancement of 100 to 4000 times of τ is noted for OHBO.

Such an enhancement of fluorescence quantum yield and fluorescence lifetime made us rethink what could be the possible reason and hence we had to look deeper into the spectroscopic characterisation. Based on careful analysis we propose here that although OHBO is produced (previous literature reports suggest that hydrolysis of methoxy group with BBr₃ yields hydroxy group)^{10b} in the reaction but it undergoes an intramolecular rearrangement and hence formation of a coumarinic structure is a possibility (Scheme 1c). So, OHBO plausibly remains not in hydroxyl derivative state but in coumarinic structure (cOHBO). This proposition has been supported by IR, and NMR spectroscopic results. For

rearrangement, presence of –OH group at *ortho* position and oxygen in oxazole ring are necessary conditions. For OHIM, OMIM or OMBO these conditions are not fulfilled and thus no intramolecular rearrangement is possible. Thus, fluorescence properties of OHIM, OMIM, and OMBO are similar to each other but quite different from that of cOHBO. Absorption maximum of cOHBO matches with that of many other coumarinic probes that absorb at around 375–400 nm.¹⁴ The best proof in support of coumarinic structure stems from the single crystal X-ray structure of cOHBO (see Fig. 1). (For lattice parameters *etc.* see ESI†).

It is quite interesting to note that at room temperature intramolecular rearrangement (Scheme 1c) is spontaneous (*i.e.* not externally photoinduced or thermo-induced). Thus, it is evident that three step tweaking especially the last step of point atom mutation in OHIM (*N* being replaced by *O*) followed by rearrangement and thus forming cOHBO is responsible for the much improved fluorescence properties. Most of the bright dyes available are either coumarin, rhodamine based. Hereby we have converted nonfluorescent GFP chromophore to fluorescent coumarinic dye. This methodology can be used for making many other nonfluorescent fluorescent protein chromophore to a bright fluorescent one. Work in this direction is currently underway.

Photoactivable fluorophores and fluorescent proteins^{11,12} are quite important towards understanding the cellular processes using different imaging techniques. Exploiting the idea of spontaneous rearrangement of OHBO leading to cOHBO (see Scheme 1c) and hence highly bright fluorescence; we have introduced an *o*-nitrobenzoyl group (see Scheme 2) in OHBO. Details of synthesis, and characterization of ONBYOHBO is given in ESI.† This photoactivable compound could be cleaved by light (at ~370 nm) leading to formation of highly fluorescent cOHBO.

The fact that indeed cOHBO is formed because of photocleavage of ONBYOHBO (see Fig. 1) is confirmed by fluorescence spectroscopy. Steady state emission spectrum of cOHBO matches quite well with that of the photo-irradiated product (see ESI†). A better proof is shown in Fig. 2. Fluorescence decay curve and the decay parameters of cOHBO (see ESI†) and that of the photoirradiated product match quite well.

This conclusively proves that cOHBO is formed from ONBYOHBO. The decay pattern of cOHBO is clearly different from photoactivable ONBYOHBO, thus, the former could be spectroscopically distinguished easily from the latter. Earlier

Fig. 1 Molecular structure of cOHBO and ONBYOHBO. Thermal ellipsoids were drawn at the 50% probability level.

Scheme 2 Synthesis of ONBYOHBO and its photocleavage to cOHBO.

Fig. 2 Fluorescence decay of ONBYOHBO, irradiated photoproduct, and cOHBO in DMF. $\lambda_{\text{ex}} = 377$ nm.

reports of introduction of *o*-nitrobenzoyl group to GFP chromophore yielded nonfluorescent product after photocleavage.¹³ Our approach has yielded more than 300 times brighter fluorescent product (cOHBO) than that in the reported literature.

Imaging of live cancerous cells like HeLa, U2OS with fluorescent dyes is quite common. However, non-invasive *in vivo* stem cell imaging¹⁵ is more modern and quite important towards regenerative medicinal therapy towards curing incurable diseases like cancer, Alzheimer disease, Parkinson disease *etc.* Fluorescence imaging of stem cells have so far been done using fluorescent proteins, quantum dots *etc.* Application of fluorescent dyes for live stem cell imaging is at its nascent stage. Thus we thought of using cOHBO for this purpose. To our delightful surprise we could image live stem cells using cOHBO (Fig. 3). (Details of imaging can be found in ESI†). We strongly believe that it is an important step towards using fluorescent dyes for stem cell imaging. In recent future many more dyes will be tried to improve the applicability of fluorescent dyes towards stem cell imaging.

To conclude, we have shown that by three point chemical tweaking parent GFP chromophore can be converted to coumarinic fluorophore with more than three hundred times enhanced fluorescence quantum yield and lifetime. This coumarinic probe could be successfully employed in live cell imaging. Employing this idea a photocleavable compound has been made which after photocleavage yields same coumarinic fluorophore. Employing our methodology many more nonfluorescent protein chromophores can be converted to highly fluorescent fluorophores. Moreover, using the photoactivation strategy highly bright fluorophores emitting at different wavelength ranging from blue to red can be prepared.

Fig. 3 Live stem cell imaging with cOHBO. Left one is fluorescence, middle one is DIC and the right one is the merged image.

PKM thanks IISER-Kolkata for instrumental facilities and financial help. Support from the Fast-Track Project (SR/FT/CS-52/2011) of DST-India is gratefully acknowledged. TC, DR, AD, PPB thank CSIR, and AG thanks UGC for their fellowship. PKM acknowledges Professor P. Ramamurthy (NCUFP), Malancha Ta (IISER-K) for Femtosecond decay measurement, and Stem cell imaging respectively.

Notes and references

- (a) R. Y. Tsien, *Annu. Rev. Biochem.*, 1998, **67**, 509–544; (b) *Green fluorescent protein: properties, application, and protocols*, ed. M. Chalfie and S. R. Kein, Wiley Interscience, New Jersey, 2nd edn, 2006.
- (a) M. Zimmer, *Chem. Rev.*, 2002, **102**, 759–781; (b) R. N. Day and M. W. Davidson, *Chem. Soc. Rev.*, 2009, **38**, 2887–2921; (c) M. Zimmer, *Chem. Soc. Rev.*, 2009, **38**, 2823–2832.
- K. Y. Chen, Y. M. Cheng, C. H. Lai, C. C. Hsu, M. L. Ho, G. H. Lee and P. T. Chou, *J. Am. Chem. Soc.*, 2007, **129**, 4534–4535.
- (a) M. Vengris, I. H. M. van Stokkum, X. He, A. F. Bell, P. J. Tonge, R. van Grondelle and D. S. Larsen, *J. Phys. Chem. A*, 2004, **108**, 4587–4598; (b) S. R. Meech, *Chem. Rev.*, 2009, **38**, 2922–2934.
- (a) W. Weber, V. Helms, J. A. McCammon and P. W. Langhoff, *Proc. Natl. Acad. Sci. U. S. A.*, 1999, **96**, 6177–6182; (b) P. Altoe, F. Bernardi, M. Garavelli, G. Orlandi and F. Negri, *J. Am. Chem. Soc.*, 2005, **127**, 3952–3963; (c) A. Usman, O. F. Mohammed, E. T. J. Nibbering, J. Dong, K. M. Solntsev and L. M. Tolbert, *J. Am. Chem. Soc.*, 2005, **127**, 11214–11215; (d) S. S. Stavrov, K. M. Solntsev, L. M. Tolbert and D. J. Huppert, *J. Am. Chem. Soc.*, 2006, **128**, 1540–1546; (e) J. Dong, K. M. Solntsev, O. Poizat and L. M. Tolbert, *J. Am. Chem. Soc.*, 2007, **129**, 10084–10085; (f) J. S. Yang, G. J. Huang, Y. H. Liu and S. M. Peng, *Chem. Commun.*, 2008, 1344–1346; (g) S. Olsen and S. C. Smith, *J. Am. Chem. Soc.*, 2008, **130**, 8677–8689; (h) A. Baldrige, S. R. Samanta, N. Jayaraj, V. Ramamurthy and L. M. Tolbert, *J. Am. Chem. Soc.*, 2010, **132**, 1498–1499; (i) A. Baldrige, S. R. Samanta, N. Jayaraj, V. Ramamurthy and L. M. Tolbert, *J. Am. Chem. Soc.*, 2011, **133**, 712–715.
- (a) N. M. Webber, K. L. Litvinenko and S. R. Meech, *J. Phys. Chem. B*, 2001, **105**, 8036–8039; (b) K. L. Litvinenko, N. M. Weber and S. R. Meech, *J. Phys. Chem. A*, 2003, **107**, 2616–2623; (c) D. Mandal, T. Tahara and S. R. Meech, *J. Phys. Chem. B*, 2004, **108**, 1102–1108; (d) S. Rafiq, B. K. Rajbongshi, N. N. Nair, P. Sen and P. Ramanathan, *J. Phys. Chem. A*, 2011, **115**, 13733–13742.
- (a) L. Wu and K. Burgess, *J. Am. Chem. Soc.*, 2008, **130**, 4089–4096; (b) K. P. Kent, L. M. Oltrogge and S. G. Boxer, *J. Am. Chem. Soc.*, 2009, **131**, 15988–15989; (c) A. Baldrige, S. R. Samanta, N. Jayaraj, V. Ramamurthy and L. M. Tolbert, *J. Am. Chem. Soc.*, 2010, **132**, 1498–1499; (d) A. Baldrige, K. M. Solntsev, C. Song, T. Tanioka, J. Kowalik, K. Hardcastle and L. M. Tolbert, *Chem.*

- Commun.*, 2010, **46**, 5686–5688; (e) K. P. Kent and S. G. Boxer, *J. Am. Chem. Soc.*, 2011, **133**, 4046–4052; (f) K. Do and S. G. Boxer, *J. Am. Chem. Soc.*, 2011, **133**, 18078–18081; (g) A. Baldrige, S. Feng, Y. T. Chang and L. M. Tolbert, *ACS Comb. Sci.*, 2011, **13**, 214–217; (h) J. S. Paige, K. Y. Wu and S. R. Jaffrey, *Science*, 2011, **333**, 642–646.
- 8 M. S. Baranov, K. A. Lukyanov, A. O. Borissova, J. Shamir, D. Kosenkov, L. V. Slipchenko, L. M. Tolbert, I. V. Yampolsky and K. M. Solntsev, *J. Am. Chem. Soc.*, 2012, **134**, 6025–6032.
- 9 J. Kang, G. Zhao, J. Xu and W. Yang, *Chem. Commun.*, 2010, **46**, 2868–2870.
- 10 (a) P. E. Ivashkin, I. V. Yampolsky and K. A. Lukyanov, *Russ. J. Bioorg. Chem.*, 2009, **35**, 726–743; (b) W. T. Chuang, C. C. Hsieh, C. H. Lai, C. H. Lai, C. W. Shih, K. W. Chen, W. Y. Hung, Y. H. Hsu and P. T. Chou, *J. Org. Chem.*, 2011, **76**, 8189–8202.
- 11 (a) D. Maurel, S. Banala, T. Laroche and K. Johnsson, *ACS Chem. Biol.*, 2010, **5**, 507–516; (b) P. Klan, T. Solomek, G. C. Bochet, A. Blanc, R. Givens, M. Rubina, V. Popik, A. Kostikov and J. Wirz, *Chem. Rev.*, 2013, **113**, 119–191; (c) Special issue, *Photochem. Photobiol. Sci.*, 2012, **11**, 433–600.
- 12 (a) G. H. Patterson and J. Lippincott-Schwartz, *Science*, 2002, **297**, 1873–1877; (b) R. Ando, H. Hama, M. Yamamoto-Hino, H. Mizun and A. Miyawaki, *Proc. Natl. Acad. Sci. U. S. A.*, 2002, **99**, 12651–12656; (c) H. Tsutsui, S. Karasawa, H. Shimizu, N. Nukina and A. Miyawaki, *EMBO Rep.*, 2005, **6**, 233–238; (d) S. Habuchi, R. Ando, P. Dedeker, W. Verheijen, H. Mizuno, A. Miyawaki and J. Hofkens, *Proc. Natl. Acad. Sci. U. S. A.*, 2005, **102**, 9511–9516; (e) N. Gagey, M. Emond, P. Neveu, C. Benbrahim, B. Goetz, I. Aujard, J. B. Baudin and L. Jullien, *Org. Lett.*, 2008, **10**, 2341–2344; (f) E. Betzig, G. H. Patterson, R. Sougrat, O. W. Lindwasser, S. Olenych, J. S. Bonifacio, M. W. Davidson, J. Lippincott-Schwartz and H. F. Hess, *Science*, 2006, **313**, 1642–1645.
- 13 D. Groff, F. Wang, S. Jockusch, N. J. Turro and P. J. Schultz, *Angew. Chem., Int. Ed.*, 2010, **49**, 7677–7679.
- 14 U. Brackmann, *Lamdaochrome Laser Dyes Lamda Physik*, Goettingen, Germany, 1986.
- 15 (a) Y. Gao, J. K. Y. Chan and C. Xu, *Am. J. Nucl. Med. Mol. Imaging*, 2013, **3**, 232–246; (b) M. R. Porcel, J. C. Wu and S. S. Gambhir, Molecular imaging of stem cells (July 30, 2009), StemBook, ed. The Stem Cell Research Community, StemBook, DOI: 10.3824/stembook.1.49.1.