

# Anhydrides as $\alpha,\beta$ -unsaturated acyl ammonium precursors: isothiourea-promoted catalytic asymmetric annulation processes†

Emily R. T. Robinson, Charlene Fallan, Carmen Simal, Alexandra M. Z. Slawin and Andrew D. Smith\*

Cite this: *Chem. Sci.*, 2013, 4, 2193

Received 22nd January 2013

Accepted 1st March 2013

DOI: 10.1039/c3sc50199j

www.rsc.org/chemicalscience

The asymmetric annulation of a range of  $\alpha,\beta$ -unsaturated acyl ammonium intermediates, formed from isothiourea HBTM 2.1 and anhydrides with either 1,3-dicarbonyls,  $\beta$ -ketoesters or azaaryl ketones gives either functionalised esters (upon ring opening), dihydropyranones or dihydropyridones in good yields (up to 93%) and high enantioselectivity (up to 97% ee).

## Introduction and background

Asymmetric organocatalysis has developed tremendously as a synthetic strategy within the last decade and a range of methodologies and catalysts have emerged that provide functionalised products with high levels of stereocontrol.<sup>1</sup> Ideally a given organocatalyst architecture should be able to participate in a range of reaction processes and display diverse modes of reactivity, while showing good catalytic efficiency and delivering products with high levels of enantioselectivity. Within this area, isothioureases,<sup>2</sup> initially employed by Birman and Okamoto as efficient *O*-acyl transfer reagents,<sup>3</sup> have been utilised in a range of kinetic resolution,<sup>4</sup> asymmetric desymmetrisation,<sup>5</sup> *C*-acylation and *C*-carboxylation processes,<sup>6</sup> as well as *O*-silylation reactions.<sup>7</sup> Recent advances have showcased the utility of isothioureases to generate ammonium enolates<sup>8</sup> from carboxylic acids and their applications in aldol-<sup>9</sup> and Michael-lactonisation processes (Fig. 1A).<sup>10</sup>

Building upon these precedents, this work demonstrates the previously unexplored ability of isothioureases to generate asymmetry by promoting the addition of a range of nucleophiles to a stereodefined  $\alpha,\beta$ -unsaturated acyl ammonium species (Fig. 1B). While Peters and Ye have invoked  $\alpha,\beta$ -unsaturated acyl ammoniums as precursors to dienolate formation,<sup>11</sup> to the best of our knowledge there are currently no processes that form C–C bonds directly *via* such intermediates. Related work in the literature has shown that NHCs<sup>12</sup> can catalytically generate  $\alpha,\beta$ -unsaturated acyl azolium intermediates through an internal redox process from alkynals,<sup>13</sup> from enals using a stoichiometric

oxidant,<sup>14</sup> directly from  $\alpha,\beta$ -unsaturated acyl fluorides or enol esters,<sup>15</sup> or alternatively from  $\alpha$ -bromo enones.<sup>16</sup> The oxidative approach from enals has been applied to a range of asymmetric C–C bond-forming reactions including aza-Claisen,<sup>17</sup> Coates-Claisen,<sup>18</sup> cyclopropanation<sup>19</sup> and Michael addition processes.<sup>20</sup> Given the recognised difficulties in accessing a wide variety of ynals and enals for such NHC-catalysed approaches and limitations associated with the scope and generality of such processes,<sup>21</sup> we envisaged a direct strategy to generate an  $\alpha,\beta$ -unsaturated acyl ammonium species from readily available  $\alpha,\beta$ -unsaturated carboxylic acids or their anhydrides. Described herein are our results concerning isothiourea-promoted asymmetric addition of 1,3-diketones,  $\beta$ -ketoesters and azaaryl ketones to  $\alpha,\beta$ -unsaturated acyl ammonium intermediates for the preparation of a range of functionalised esters, stereo-defined dihydropyranones and dihydropyridones in highly enantioenriched form (up to 97% ee).




Fig. 1 Proposed access to enantioenriched annulation products *via* an unexplored  $\alpha,\beta$ -unsaturated acyl ammonium species.


EaStCHEM, School of Chemistry, University of St Andrews, North Haugh, St Andrews, KY16 9ST, UK. E-mail: ads10@st-andrews.ac.uk

† Electronic supplementary information (ESI) available: All spectroscopic data and procedures for novel compounds. CCDC 916302. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c3sc50199j



## Proof of concept and reaction optimisation: annulations using diketone nucleophiles

Initial investigations focused upon generation of an  $\alpha,\beta$ -unsaturated acyl ammonium directly from cinnamic acid **1** (Scheme 1A), *via in situ* activation using 4-methoxybenzoic anhydride (PMBA) and isothioureia HBTM 2.1 **5** (20 mol%).<sup>22</sup> Employing diketone **4** as the nucleophile provided dihydropyranone **2** in modest 25% isolated yield, albeit with an encouraging 95% ee. Cinnamic anhydride **3** was next evaluated as an alternative acyl ammonium precursor,<sup>23</sup> giving dihydropyranone **2** in improved 49% yield and 95% ee (Scheme 1B). These initial findings served as a benchmark for further optimisation with the aim to lower catalyst loadings and improve isolated yields.<sup>24</sup>



**Scheme 1** Initial proof of concept studies. <sup>a</sup>Isolated yield of **2**; <sup>b</sup>determined by HPLC analysis.

Further studies showed that *in situ* ring opening of dihydropyranone **2** with MeOH led to consistently higher isolated yields of the functionalised ester product **6** (Table 1). Variation

**Table 1** Optimisation studies

Entry	HBTM 2.1 (mol%)	Base	Yield <sup>a</sup>	ee <sup>b</sup> (%)
1	20	EtN( <i>i</i> Pr) <sub>2</sub>	77	96
2	10	EtN( <i>i</i> Pr) <sub>2</sub>	70	95
3	5	EtN( <i>i</i> Pr) <sub>2</sub>	62 <sup>c</sup>	—
4 <sup>d</sup>	5	EtN( <i>i</i> Pr) <sub>2</sub>	45	99
5	20	DBU	86	73
6	20	PS-BEMP	85	95
7	10	PS-BEMP	82	95
8	5	PS-BEMP	83	96
9 <sup>d</sup>	5	PS-BEMP	50	83
10	2.5	PS-BEMP	69	93
11	1	PS-BEMP	62	93


<sup>a</sup> Isolated yield of **6**. <sup>b</sup> Determined by HPLC analysis. <sup>c</sup> Conversion determined by <sup>1</sup>H NMR spectroscopic analysis of the unpurified reaction mixture. <sup>d</sup> Reaction conducted in THF.

of the base showed that EtN(*i*Pr)<sub>2</sub>, DBU or PS-BEMP (polymer-supported 2-*t*-butylimino-2-diethylamino-1,3-dimethylperhydro-1,3,2-diazaphosphorine) could be used in this process, although DBU gave **6** with reduced enantioselectivity (entry 5).<sup>25</sup> The use of PS-BEMP proved optimal, allowing the catalyst loading of HBTM 2.1 **5** to be reduced to 1 mol% without compromising product enantioselectivity, albeit with reduced product yields (entry 11). Performing the reaction in THF led to a reduced yield (entries 4 and 9). While catalytic asymmetric Michael additions to nitro-olefins and enones are well documented,<sup>26</sup> this strategy formally allows the asymmetric Michael addition of diketones to  $\alpha,\beta$ -unsaturated carboxylic acid derivatives for which there is only limited precedent.<sup>27</sup>

## Reaction scope and generality

The generality of this process was next probed, initially through variation of the  $\alpha,\beta$ -unsaturated anhydride component (Table 2). Using HBTM 2.1 **5** (5 mol%) and a number of symmetrical diketones, this protocol tolerates a range of 2-, 3-, and 4-substituted  $\beta$ -aryl groups containing either electron-withdrawing or electron-donating groups, as well as heteroaryl

**Table 2** Substrate scope




<sup>a</sup> Isolated yield of **6–21**. <sup>b</sup> Determined by HPLC analysis. <sup>c</sup> 10 mol% catalyst. <sup>d</sup> Reaction carried out at –78 °C. <sup>e</sup> Reaction conducted in THF.



substituents, giving the corresponding functionalised esters in good yield (up to 86%) and high enantioselectivity (90–97% ee, **6–14**). Notable reactivity trends within this series indicate that anhydrides containing electron deficient  $\beta$ -aryl units give higher product conversion and isolated yields than their electron rich  $\beta$ -aryl counterparts (product **9**).  $\beta$ -Alkyl substituents within the anhydride are also tolerated (a significant advantage over the NHC-catalysed systems that typically exhibit low enantioselectivity in similar transformations),<sup>20b</sup> although low temperatures are necessary to achieve optimal enantioselectivity, resulting in only moderate reaction efficiency and reduced product yields (**15** and **16**). Variation of the diketone functionality was next investigated (Table 2, **17–21**). A range of substituted aryl and heteroaryl diketones participate in this reaction process, giving functionalised esters **17–21** in moderate to good yield and high enantioselectivity.<sup>28</sup>

Further investigations probed the stereospecificity of this asymmetric annulation protocol (Scheme 2). While (*E,E*)-cinnamic anhydride **3** gave functionalised ester (*S*)-**6** in 83% yield and 96% ee (Scheme 2A), (*Z,Z*)-cinnamic anhydride **22** gave (*R*)-**6** in reduced 41% yield and only 30% ee (Scheme 2B),<sup>29</sup> indicating the necessity of the (*E*)-configuration for maximum enantiocontrol.




**Scheme 2** Stereospecificity of asymmetric annulation process. <sup>a</sup>Isolated yield of **6**; <sup>b</sup>determined by HPLC analysis.

Subsequent studies probed the ability of non-symmetric dicarbonyls to participate in this protocol. Using cinnamic anhydride **3**, 1-phenyl-1,3-dibutanone **23** generated a 70 : 30 mixture of regioisomeric dihydropyranones **24** and **25** in 88% overall yield and 70% and 61% ee, respectively (Scheme 3A),<sup>30</sup> while ethyl benzoylacetate **26** gave dihydropyranone **27** in 60% isolated yield as a single regioisomer and in 94% ee (Scheme 3B).


The generality of this process was next examined (Table 3), with  $\beta$ -aryl and  $\beta$ -heteroaryl substituents within the anhydride tolerated, in all cases giving the corresponding dihydropyranones **27–32** in acceptable yield (46–70%) and high ee (89–96%).<sup>31</sup>

Having probed the viability of this process, our attention turned to 1,3-dicarbonyl systems that are not tolerated in related NHC-catalysed processes. For example, aliphatic cyclic Michael donors such as 1,3-cyclohexanedione **33** have been reported by




**Scheme 3** Regioselectivity of annulation using unsymmetrical nucleophiles. <sup>a</sup>Isolated yield; <sup>b</sup>determined by HPLC analysis.

**Table 3** Use of ethyl benzoylacetate as nucleophile




<sup>a</sup> Isolated yield. <sup>b</sup> Determined by HPLC analysis.

Bode to be ineffective in NHC-catalysis using  $\alpha,\beta$ -unsaturated acyl azoliums.<sup>21</sup> However, in this organocatalysed process, diketone **33** demonstrated favourable reactivity and provided **34** in high yield and good enantioselectivity (Scheme 4).




**Scheme 4** 1,3-Cyclohexanedione as a Michael donor. <sup>a</sup>Isolated yield; <sup>b</sup>determined by HPLC analysis.





**Scheme 5** Asymmetric addition of azaaryl ketone **35**. <sup>a</sup>Isolated yield; <sup>b</sup>determined by HPLC analysis; <sup>c</sup>determined from <sup>1</sup>H NMR of unpurified reaction mixture; <sup>d</sup>following a single recrystallisation.

**Table 4** Anhydride scope with azaaryl ketone **35** as a nucleophile in asymmetric annulation



<sup>a</sup> Isolated yield. <sup>b</sup> Determined by HPLC analysis. <sup>c</sup> Determined from <sup>1</sup>H NMR of unpurified reaction mixture.

## Beyond 1,3-dicarbonyls: azaaryl ketones as nucleophiles

Encouraged by the novel and complementary reactivity of the isothiourea catalysis, the scope of this asymmetric annulation was extended beyond the use of 1,3-dicarbonyl nucleophiles. Gratifyingly, azaaryl ketone **35** proved a competent nucleophile and displayed improved reactivity compared with simple 1,3-dicarbonyls. This increased reactivity allows bench grade solvents in an open flask atmosphere to be employed, PS-BEMP can be replaced with more cost efficient EtN(*i*Pr)<sub>2</sub> and a lower catalyst loading of 1 mol% could be routinely employed. The use of this nucleophile led to intriguing regioselectivity. Cyclisation occurs preferentially through the benzothiazole nitrogen, generating dihydropyridone **36a** as the major product, in addition to dihydropyranone **36b** as the minor product (88 : 12 regioisomeric ratio). These heterocycles were readily separable by chromatography, with the major product **36a** obtained in excellent yield and high enantioselectivity (97% ee after a single recrystallisation, Scheme 5).

Next, the substrate scope with this nucleophile was examined with respect to the anhydride component (Table 4). The increased reactivity of the azaaryl ketone **35** provided a wide range of enantioenriched heterocycles in excellent yields and enantioselectivities. For example, while anhydrides bearing electron-rich and aliphatic substituents were modestly tolerated using diketone **4** (Table 2) in terms of both reactivity and enantioselectivity, the use of azaaryl ketone **35** with the same anhydrides leads to enantioenriched products in high yields under our reaction conditions (Table 4).

## Mechanistic investigations

In related NHC-catalysed processes involving  $\alpha,\beta$ -unsaturated acyl azolium intermediates two potential mechanistic pathways have been proposed; a Michael addition–lactonisation process with diketonyls, ketene acetals and enamines (favoured by Studer and Mayr)<sup>32</sup> or alternatively an initial 1,2-addition followed by a [3,3]-Claisen rearrangement with Kojic acid and enamine derivatives (favoured by Bode)<sup>21</sup> to facilitate the formation of enantioenriched products. Similarly, in our isothiourea-promoted annulation, related catalytic cycles depicted in Fig. 2(a) and (b), could potentially be responsible for the generation of the annulation products with high enantiocontrol. Both cycles involve an initial *N*-acylation of HBTM 2.1 **5** with an anhydride to generate the corresponding  $\alpha,\beta$ -unsaturated acyl ammonium **41**. The *s-cis* conformation of ammonium **41** is presumably favoured, with the carbonyl oxygen hypothesised to adopt a *syn*-conformation with respect to the isothiourea S atom due to a stabilising non-bonding O–S interaction ( $n_o$  to  $\sigma_{C-S}^*$ ).<sup>33</sup> In pathway (a), Michael addition<sup>31</sup> of diketone enolate **42** to the *Re* face of the  $\alpha,\beta$ -unsaturated acyl ammonium **41** gives intermediate **43**,<sup>34</sup> subsequent proton transfer followed by lactonisation generates the desired dihydropyranone **44**, which can be either isolated or subsequently ring opened with MeOH to generate the ester products **45** in high ee. Alternatively, pathway (b) demonstrates that







Fig. 2 Proposed mechanisms of asymmetric dihydropyranone formation via (a) Michael addition–lactonisation or (b) 1,2-addition, [3,3]-rearrangement.


1,2-addition to the  $\alpha,\beta$ -unsaturated acyl ammonium **42** to generate intermediate **46** followed by a [3,3]-rearrangement and subsequent proton transfer/lactonisation process would lead to the same enantioenriched products.

To gain insight into the favoured mechanistic pathway, potential intermediates were synthesised and subjected to the reaction conditions. Interestingly, Lupton has previously shown that 1,2-addition of NHCs to enol esters such as **47** facilitates formation of dihydropyranone **49** in the presence of NHC catalyst **48**, albeit with moderate enantioselectivity (Scheme 6A). In this regard, **50** was prepared from cinnamoyl chloride and dicarbonyl **4**. This potential [3,3]-rearrangement precursor was examined under the reaction conditions, resulting in no conversion into dihydropyranone **2** (Scheme 6B)<sup>15a-c</sup> after 24 h at room temperature. While this result does not rule out either mechanistic pathway, it indicates that **50** is not a likely intermediate in this process. Additionally the absence of any 1,2-addition products such as **50** in the <sup>1</sup>H NMR of all unpurified reaction mixtures provides further evidence to support this view.

A. Lupton (ref 15a):




B. [3,3]-rearrangement?




Scheme 6 Probing the mechanism.

Evidence supporting the intermediacy of an acyl ammonium species was next obtained. Treatment of *trans*-cinnamoyl chloride **51** with HBTM 2.1 **5** in  $\text{CH}_2\text{Cl}_2$  gave the  $\alpha,\beta$ -unsaturated acyl ammonium salt **52** that was isolated in high yield (Scheme 7).<sup>35</sup> X-ray crystallography confirmed the structure of **52** and provided further support for the *syn* geometry of the carbonyl oxygen and the isothioureia S atom, presumably due to the previously hypothesised stabilising non-bonding O–S interaction ( $n_{\text{O}}$  to  $\sigma_{\text{C-S}}^*$ ).<sup>33</sup>



Scheme 7 Isolation and representation of the X-ray crystal structure of  $\alpha,\beta$ -unsaturated acyl ammonium salt **52**.





**Scheme 8**  $\alpha,\beta$ -Unsaturated acyl ammonium salt **52** as organocatalyst.

The isolated acyl ammonium salt **52** was next examined as a precatalyst, employing cinammic anhydride **3** and diketone **4** under our optimised reaction conditions (Scheme 8). The functionalised ester **6** was isolated in similar yield and enantioselectivity (82% yield, 91% ee) compared with the use of organocatalyst HBTM 2.1 **5** (83% yield, 95% ee) in this process, consistent with acyl ammonium salt **52** being an intermediate in this asymmetric annulation.

At this juncture we cannot rule out any plausible mechanistic pathway that involves an  $\alpha,\beta$ -unsaturated acyl ammonium ion, although we currently favour a catalytic cycle involving a Michael addition–lactonisation sequence, described in Fig. 2(a). Further mechanistic investigations and DFT calculations to provide insight into the pathway in operation are underway and will be reported in due course.

## Conclusions

To conclude, HBTM 2.1 **5** promotes the asymmetric annulation of a range of nucleophiles including 1,3-diketones,  $\beta$ -ketoesters and azaryl ketones to (*E,E*)- $\alpha,\beta$ -unsaturated anhydrides, giving either functionalised esters (upon ring opening), dihydropyranones, or dihydropyridones in good yields (up to 86%) and high enantioselectivity (up to 97% ee) via a postulated  $\alpha,\beta$ -unsaturated acyl ammonium intermediate. Current research from this laboratory is directed toward developing alternative uses of isothioureas and other Lewis bases in asymmetric catalysis, and exploiting  $\alpha,\beta$ -unsaturated acyl ammonium intermediates for a range of synthetic procedures.

## Acknowledgements

We thank the Royal Society for a University Research Fellowships (ADS), the EPSRC (ER) and the EU (CF and CS) for funding, as well as European Research Council under the European Union's Seventh Framework Programme (FP7/2007–2013) ERC grant agreement n° 279850. We also thank the EPSRC National Mass Spectrometry Service Centre (Swansea).

## Notes and references

- For selected reviews on organocatalysis see: (a) P. I. Dalko and L. Moisan, *Angew. Chem., Int. Ed.*, 2004, **43**, 5138–5175; (b) D. W. C. MacMillan, *Nature*, 2008, **455**, 304–308; (c) A. Erkkilä, I. Majaner and P. M. Pihko, *Chem. Rev.*, 2007, **107**, 5416–5470; (d) S. Mukherjee, J. W. Yang, S. Hoffmann and B. List, *Chem. Rev.*, 2007, **107**, 5471–5569; (e) M. J. Gaunt and C. C. C. Johansson, *Chem. Rev.*, 2007, **107**, 5596–5605; (f) S. E. Denmark and G. L. Beutner, *Angew. Chem., Int. Ed.*, 2008, **47**, 1560–1638; (g) C. Gronal, M. Jeanty and D. Enders, *Nat. Chem.*, 2010, **2**, 167–178; (h) A. Moyana and R. Rios, *Chem. Rev.*, 2011, **111**, 4703–4832; (i) F. Giacalone, M. Gruttadauria, P. Agrigento and R. Noto, *Chem. Soc. Rev.*, 2012, **41**, 2406–2447.
- For a comprehensive review of the catalytic uses of guanidines, amidines and isothioureas see: J. E. Taylor, S. D. Bull and J. M. J. Williams, *Chem. Soc. Rev.*, 2012, **41**, 2109–2121.
- (a) V. B. Birman, H. Jiang, X. Li, L. Guo and E. W. Uffman, *J. Am. Chem. Soc.*, 2006, **128**, 6536–6537; (b) M. Kobayashi and S. Okamoto, *Tetrahedron Lett.*, 2006, **47**, 4347–4350; (c) V. B. Birman and X. Li, *Org. Lett.*, 2006, **8**, 1351–1354.
- For kinetic resolutions using anhydrides as acylating agents see: (a) V. B. Birman and L. Guo, *Org. Lett.*, 2006, **8**, 4859–4861; (b) V. B. Birman and X. Li, *Org. Lett.*, 2008, **10**, 1115–1118; (c) D. Belmessieri, C. Joannesse, P. A. Woods, C. MacGregor, C. Jones, C. D. Campbell, C. P. Johnston, N. Duguet, C. Concellón, R. A. Bragg and A. D. Smith, *Org. Biomol. Chem.*, 2011, **9**, 559–570; (d) X. Li, H. Jiang, E. W. Uffman, L. Guo, Y. Zhang, X. Yang and V. B. Birman, *J. Org. Chem.*, 2012, **77**, 1722–1737. For kinetic resolutions using carboxylic acids as acylating agents utilising *in situ* formation of a reactive mixed anhydride see: (e) I. Shiina and K. Nakata, *Tetrahedron Lett.*, 2007, **48**, 8314–8317; (f) I. Shiina, K. Nakata, M. Sugimoto, Y. Onda, T. Iizumi and K. Ono, *Heterocycles*, 2009, **77**, 801–810; (g) X. Yang and V. B. Birman, *Adv. Synth. Catal.*, 2009, **351**, 2301–2304; (h) I. Shiina and K. Nakata, *Heterocycles*, 2010, **80**, 169–175; (i) I. Shiina, K. Nakata and Y. Onda, *Eur. J. Org. Chem.*, 2008, 5887–5890; (j) I. Shiina, K. Nakata, K. Ono, M. Sugimoto and A. Sekiguchi, *Chem.–Eur. J.*, 2010, **16**, 167–172; (k) K. Nakata, Y. Onda, K. Ono and I. Shiina, *Tetrahedron Lett.*, 2010, **51**, 5666–5669; (l) I. Shiina, K. Ono and K. Nakata, *Chem. Lett.*, 2011, **40**, 147–149. For the kinetic resolution of  $\beta$ -lactams by asymmetric alcoholysis see: (m) V. D. Bumbu and V. B. Birman, *J. Am. Chem. Soc.*, 2011, **133**, 13902–13905; by asymmetric acylation see: (n) X. Yang, V. D. Bumbu and V. B. Birman, *Org. Lett.*, 2012, **13**, 4755–4757.
- (a) X. Yang, G. Lu and V. B. Birman, *Org. Lett.*, 2010, **12**, 892–895; (b) X. Yang and V. B. Birman, *Angew. Chem., Int. Ed.*, 2011, **50**, 5553–5555.
- For *C*-acylation and *C*-carboxylation reaction processes see: (a) F. R. Dietz and H. Gröger, *Synthesis*, 2009, 4208–4218; (b) C. Joannesse, C. P. Johnston, C. Concellón, C. Simal, D. Philp and A. D. Smith, *Angew. Chem., Int. Ed.*, 2009, **48**,



- 8914–8918; (c) P. A. Woods, L. C. Morrill, T. Lebl, A. M. Z. Slawin, R. A. Bragg and A. D. Smith, *Org. Lett.*, 2010, **12**, 2660–2663; (d) C. Joannesse, L. C. Morrill, C. D. Campbell, A. M. Z. Slawin and A. D. Smith, *Synthesis*, 2011, 1865–1879; (e) B. Viswambharan, T. Okimura, S. Suzuki and S. Okamoto, *J. Org. Chem.*, 2011, **76**, 6678–6685; (f) P. A. Woods, L. C. Morrill, R. A. Bragg and A. D. Smith, *Chem.–Eur. J.*, 2011, **17**, 11060–11067.
- 7 C. I. Sheppard, J. L. Taylor and S. L. Wiskur, *Org. Lett.*, 2011, **13**, 3794–3797.
- 8 (a) G. S. Cortez, R. L. Tennyson and D. Romo, *J. Am. Chem. Soc.*, 2001, **123**, 7945–7946; (b) G. S. Cortez, S. H. Oh and D. Romo, *Synthesis*, 2001, 1731–1736; (c) S. H. Oh, G. S. Cortez and D. Romo, *J. Org. Chem.*, 2005, **70**, 2835–2838; (d) H. Henry-Riyad, C. Lee, V. C. Purohit and D. Romo, *Org. Lett.*, 2006, **8**, 4363–4366; (e) G. Ma, H. Nguyen and D. Romo, *Org. Lett.*, 2007, **9**, 2143–2146; (f) H. Nguyen, G. Ma and D. Romo, *Chem. Commun.*, 2010, **46**, 4803–4805; (g) K. A. Morris, K. M. Arendt, S. H. Oh and D. Romo, *Org. Lett.*, 2010, **12**, 3764–3767; (h) H. Nguyen, G. Ma, T. Gladysheva, T. Fremgen and D. Romo, *J. Org. Chem.*, 2011, **76**, 2–12.
- 9 (a) V. C. Purohit, A. S. Matla and D. Romo, *J. Am. Chem. Soc.*, 2008, **130**, 10478–10479; (b) C. A. Leverett, V. C. Purohit and D. Romo, *Angew. Chem., Int. Ed.*, 2010, **49**, 9479–9483.
- 10 (a) D. Belmessieri, L. C. Morrill, C. Simal, A. M. Z. Slawin and A. D. Smith, *J. Am. Chem. Soc.*, 2011, **133**, 2714–2720; (b) C. Simal, T. Lebl, A. M. Z. Slawin and A. D. Smith, *Angew. Chem., Int. Ed.*, 2012, **51**, 3653–3657; (c) L. C. Morrill, T. Lebl, A. M. Z. Slawin and A. D. Smith, *Chem. Sci.*, 2012, **3**, 2088–2093.
- 11 There is only limited precedent for the potential intermediacy of  $\alpha,\beta$ -unsaturated acyl ammonium ions in asymmetric catalysis. For dienolate formation that may proceed via an  $\alpha,\beta$ -unsaturated acyl ammonium ion see: (a) P. S. Tiseni and R. Peters, *Angew. Chem., Int. Ed.*, 2007, **46**, 5325–5328; (b) P. S. Tiseni and R. Peters, *Chem.–Eur. J.*, 2010, **16**, 2503–2517; (c) L. T. Shen, L. H. Sun and S. Ye, *J. Am. Chem. Soc.*, 2011, **133**, 15894–15897. Recently,  $\alpha,\beta$ -unsaturated acyl ammonium intermediates have been accessed from  $\alpha,\beta$ -unsaturated acid fluorides and employed in 1,3-dipolar cycloaddition reactions, albeit with low enantioselectivity, see: (d) S. Pandiancherri, S. J. Ryan and D. W. Lupton, *Org. Biomol. Chem.*, 2012, **10**, 7903–7911.
- 12 For reviews see: (a) D. Enders, O. Niemeier and A. Henseler, *Chem. Rev.*, 2007, **107**, 5606–5655; (b) N. Marion, S. Diez-González and S. P. Nolan, *Angew. Chem., Int. Ed.*, 2007, **46**, 2988–3000; (c) V. Nair, S. Vellalath and B. P. Babu, *Chem. Soc. Rev.*, 2008, **37**, 2691–2698; (d) J. L. Moore and T. Rovis, in *Topics in Current Chemistry*, ed. B. List, Springer, Berlin, vol. 291, 2009, pp. 77–144; (e) C. D. Campbell, K. B. Ling and A. D. Smith, in *N-Heterocyclic Carbenes in Transition Metal Catalysis and Organocatalysis, Catalysis by Metal Complexes 32*, ed. C. S. J. Cazin, Springer, Berlin, 2010, pp. 263–297; (f) S. J. Ryan, L. Candish and D. W. Lupton, *Chem. Soc. Rev.*, 2013, DOI: 10.1039/c3cs35522e.
- 13 (a) K. Zeitler, *Org. Lett.*, 2006, **8**, 637–640. For a mechanistic investigation of the formation of  $\alpha,\beta$ -unsaturated acyl azoliums from alkynals see: (b) J. Mahatthananchai, P. Zheng and J. W. Bode, *Angew. Chem., Int. Ed.*, 2010, **49**, 9479–9483.
- 14 (a) S. De Sarkar and A. Studer, *Angew. Chem., Int. Ed.*, 2010, **49**, 9266–9269; (b) R. S. Reddy, J. N. Rosa, L. F. Veiros, S. Caddick and P. M. P. Gois, *Org. Biomol. Chem.*, 2011, **9**, 3126–3129; (c) S. D. Sarkar and A. Studer, *Org. Lett.*, 2010, **12**, 1992–1995.
- 15 (a) S. J. Ryan, L. Candish and D. W. Lupton, *J. Am. Chem. Soc.*, 2009, **131**, 14176–14177; (b) L. Candish and D. W. Lupton, *Org. Lett.*, 2010, **12**, 4836–4839; (c) S. J. Ryan, L. Candish and D. W. Lupton, *J. Am. Chem. Soc.*, 2011, **133**, 4694–4697; (d) C. Yao, D. Wang, J. Lu, T. Li, W. Jiao and C. Yu, *Chem.–Eur. J.*, 2012, **18**, 1914–1917; (e) L. Candish and D. W. Lupton, *J. Am. Chem. Soc.*, 2013, **135**, 58–61.
- 16 (a) C. Yao, D. Wang, J. Lu, T. Li, W. Jiao and C. Yu, *Chem.–Eur. J.*, 2012, **18**, 1914–1917; (b) F.-G. Sun, L.-H. Sun and S. Ye, *Adv. Synth. Catal.*, 2012, **353**, 3134–3138; (c) B. Zhang, P. Feng, Y. Cui and N. Jiao, *Chem. Commun.*, 2012, **48**, 7280–7282.
- 17 B. Wanner, J. Mahatthananchai and J. W. Bode, *Org. Lett.*, 2011, **13**, 5378–5381.
- 18 J. Kaeobamrung, J. Mahatthananchai, P. Zheng and J. W. Bode, *J. Am. Chem. Soc.*, 2010, **132**, 8810–8812.
- 19 A. Biswas, S. De Sarkar, L. Tebben and A. Studer, *Chem. Commun.*, 2012, **48**, 5190–5192.
- 20 (a) Z. Q. Zhu and J. C. Xiao, *Adv. Synth. Catal.*, 2010, **352**, 2455–2458; (b) Z.-Q. Rong, M.-Q. Jia and S.-L. You, *Org. Lett.*, 2011, **13**, 4080–4083; (c) Z.-Q. Zhu, X.-L. Zheng, N.-F. Jiang, X. Wang and J.-C. Xiao, *Chem. Commun.*, 2011, **47**, 8670–8672. For a Michael-initiated cascade process see: (d) A. Biswas, S. De Sarkar, R. Frohlich and A. Studer, *Org. Lett.*, 2011, **13**, 4966–4969.
- 21 J. Mahatthananchai, J. Kaeobamrung and J. W. Bode, *ACS Catal.*, 2012, **2**, 494–503.
- 22 Screening of a range of isothioureas for catalytic activity and asymmetry in these processes showed that HBTM 2.1 was optimal.
- 23 All  $\alpha,\beta$ -unsaturated anhydrides in this manuscript were readily prepared from the corresponding  $\alpha,\beta$ -unsaturated acid by EDCI-mediated coupling. See ESI† for further information.
- 24 The absolute configuration within **2** was proven by comparison of its specific rotation  $[\alpha]_{\text{D}}^{22} -7.6$  (*c* 0.5 in  $\text{CHCl}_3$ ) with the literature  $\{[\text{lit.}^{20b} [\alpha]_{\text{D}}^{22} -6.5$  (*c* 1.0 in  $\text{CHCl}_3$ )\}.
- 25 Control studies showed that DBU alone can act as a catalyst for this transformation, as 0.6 equivalents of DBU in the absence of isothiourea catalyst afforded 75% consumption of diketone starting material.
- 26 For selective reviews on asymmetric Michael additions see: (a) M. P. Sibi and S. Manyem, *Tetrahedron*, 2000, 8033–8061; (b) N. Krause and A. Hoffmann-Röder, *Synthesis*, 2001, **2**, 171–196; (c) J. Christoffers and A. Baro, *Angew. Chem., Int. Ed.*, 2003, **42**, 1688–1690; (d) S. B. Tsogoeva, *Eur. J. Org. Chem.*, 2007, 1701–1716.



- 27 The use of  $\alpha,\beta$ -unsaturated acyl phosphonates as ester equivalents has been demonstrated see: (a) H. Jiang, M. W. Paixão, D. Monge and K. A. Jørgensen, *J. Am. Chem. Soc.*, 2010, **132**, 2775–2783. For Michael additions of ethyl benzoylacetates to *N*-alkylmaleimides using chiral guanidines see: (b) Z. Jiang, W. Ye, Y. Yang and C.-H. Tana, *Adv. Synth. Catal.*, 2008, **350**, 2345–2351.
- 28 Aliphatic diketones such as 2,4-pentanedione led to products with low levels of enantiocontrol. With cinnamic anhydride, 2,4-pentanedione in the presence of HBTM 2.15 gave the corresponding ester in 64% yield and 37% ee. See ESI† for details.
- 29 (*Z,Z*)-Cinnamic anhydride **22** was prepared by Lindlar reduction of ethyl 3-phenylpropionate, ester hydrolysis to give (*Z*)-cinnamic acid and subsequent anhydride formation. For full experimental procedures see ESI.†
- 30 *In situ* ring opening by addition of methanol to dihydropyranones derived from unsymmetrical Michael donors led to a statistical mixture of diastereoisomers and so this strategy was not pursued further.
- 31 Addition of ethyl acetoacetate to cinnamic anhydride **3** afforded the corresponding dihydropyranone in 72% yield, 70% ee; addition of ethyl benzoylacetate to crotonic anhydride afforded the corresponding dihydropyranone in 45% yield, 65% ee. See ESI† for further information.
- 32 For a mechanistic investigation regarding the preferred 1,4-addition of diketone enolates and alternative nucleophiles to  $\alpha,\beta$ -unsaturated acyl azoliums see: R. C. Samantha, B. Maji, S. De Sarkar, K. Bergander, R. Fröhlich, C. Mück-Lichtenfeld, H. Mayr and A. Studer, *Angew. Chem., Int. Ed.*, 2012, **51**, 5234–5238. Additionally, a recent computational study highlights the similarity of the two potential mechanistic pathways: E. Lyngvi, J. W. Bode and F. Schoenebeck, *Chem. Sci.*, 2012, **3**, 2346–2350.
- 33 Such non-bonded S–O interactions have been widely recognised; for overviews see: (a) V. I. Minkin and R. M. Minyaev, *Chem. Rev.*, 2001, **101**, 1247–1266; (b) K. A. Brameld, B. Kuhn, D. C. Reuter and M. Stahl, *J. Chem. Inf. Model*, 2008, **48**, 1–24. For  $n_{\text{O}}$  to  $\sigma_{\text{C-S}}^*$  interactions discussed with respect to (acylimino) thiadiazoline derivatives see: (c) Y. Nagao, T. Hirata, S. Goto, S. Sano, A. Kakehi, K. Iizuka and M. Shiro, *J. Am. Chem. Soc.*, 1998, **120**, 3104–3310. In the context of isothiourea-mediated catalysis the importance of these interactions have been recognised see: (d) V. B. Birman, X. Li and Z. Han, *Org. Lett.*, 2007, **9**, 37–40; (e) P. Liu, X. Yang, V. B. Birman and K. N. Houk, *Org. Lett.*, 2012, **14**, 3288–3291. See also ref. 9b.
- 34 *Re* face addition to the  $\alpha,\beta$ -unsaturated acyl ammonium is favoured for  $\beta$ -aryl substituents, with *Si* face addition favoured to  $\beta$ -alkyl substituents due to CIP priority changes.
- 35 The acyl ammonium chloride salt **52** was characterised by single-crystal X-ray diffraction.†

