Cite this: *Analyst*, 2014, 139, 6058

Correction: Spatial organization of *Pseudomonas aeruginosa* biofilms probed by combined matrix-assisted laser desorption ionization mass spectrometry and confocal Raman microscopy

Rachel N. Masyuko,^a Eric J. Lanni,^b Callan M. Driscoll,^c Joshua D. Shrout,^c
Jonathan V. Sweedler^{*b} and Paul W. Bohn^{*ad}

DOI: 10.1039/c4an90079k

www.rsc.org/analyst

Correction for 'Spatial organization of *Pseudomonas aeruginosa* biofilms probed by combined matrix-assisted laser desorption ionization mass spectrometry and confocal Raman microscopy' by Rachel N. Masyuko *et al.*, *Analyst*, 2014, DOI: 10.1039/c4an00435c.

In the original study for the DOI detailed above, Masyuko *et al.* examined the Raman and mass spectrometric microspectra and images accompanying transition from planktonic cells to biofilms in wild-type *Pseudomonas aeruginosa* and in a quorum sensing (QS)-deficient mutant, $\Delta lasI\Delta rhII$. In the course of follow-up studies, we became aware that some of the Raman spectra attributed to wild-type biofilm could have arisen from a contaminant, microparticulate polystyrene (PS). Subsequently, a new series of wild-type *Pseudomonas* biofilms were grown under rigorously PS-free conditions, and new spectra and images were acquired. For the new data shown in the figures below, Fig. 1 replaces the spectrum shown in Fig. 1(b) of the original and Fig. 3 replaces the images shown in Figs. 3(b) and S2 of the original. A detailed consideration of the bands and their assignments in the new biofilm spectrum is given in the Supplemental Information (see below).

While detailed analysis of the biofilm spectrum, Fig. 1(b), and the image acquired, Fig. 3(b), differ from the corresponding data in the original, the major conclusions of the paper are unaffected. In particular:

- (1) the wild-type biofilm spectra are markedly different than the corresponding wild-type planktonic cell spectra, *cf.* Fig. 1(b) *vs.* Fig. 1(a);
- (2) wild-type biofilms produce spectra that are characteristic of rhamnolipids, secreted as part of the biofilm formation process, and by a co-secreted protein/peptide component; and
- (3) these data are consistent with the formation of a thick (relative to the sampling depth of the CRM) biofilm after 72 h in the wild type cells, but not in the QS mutant.

Supplemental Information

Text in the original describing the Raman spectra (Figs. 1(b) and 3(b) in the original) should be replaced with the following.

Upon biofilm formation, dramatic changes occur in the spectrum, Fig. 1(b). Biofilms cultivated on bare Si exhibit a greatly reduced SiO₂ background (915–1015 cm⁻¹), presumably because biofilms at 72 h are much thicker than the <1 μm confocal depth of the CRM. In addition, the strong DNA/RNA-related bands at 747 (thymine), 1126 (cytosine), and 1310 cm⁻¹ (adenine) are all diminished in relative intensity. The strong band at 1585 cm⁻¹ is reduced in strength and a new band at 1606 cm⁻¹ grows in. In addition, narrow bands with peaks centered at 1005 cm⁻¹ and 1034 cm⁻¹ appear as well as bands centered at 830 cm⁻¹, 1166 cm⁻¹, 1186 cm⁻¹, 1417 cm⁻¹, 1445 cm⁻¹ and 1558 cm⁻¹. Most striking is the intense peak at 1005 cm⁻¹ attributed to symmetric ring breathing vibrations in phenylalanine and tryptophan, indicative of proteins.¹ Other bands characteristic of proteins are at 830 cm⁻¹ ring breathing vibrations in tyrosine (with a possible contribution from rhamnolipid), 1186 cm⁻¹ arising from C–H in plane bending vibrations in tyrosine and phenylalanine,² 1214 cm⁻¹ and 1246 cm⁻¹ both assigned to amide III vibrations, 1558 cm⁻¹ linked to the C–C pyrrole ring stretching vibrations in tryptophan and the two bands at 621 cm⁻¹ and 1605 cm⁻¹ arising from in plane ring deformation and C=C stretching vibrations in phenylalanine, respectively.¹

^aDepartment of Chemistry and Biochemistry, University of Notre Dame, Notre Dame, IN 46556, USA. E-mail: pbohn@nd.edu^bDepartment of Chemistry, University of Illinois at Urbana-Champaign, 600 S. Mathews Ave., Urbana, IL 61801, USA. E-mail: jsweedle@illinois.edu^cDepartment of Civil and Environmental Engineering and Earth Sciences, University of Notre Dame, Notre Dame, IN 46556, USA^dDepartment of Chemical and Biomolecular Engineering, University of Notre Dame, Notre Dame, IN 46556, USA

Fig. 1 Raman microspectra of wild type *Pseudomonas aeruginosa*. (a) Planktonic cells, as deposited on Au-coated Si substrate. (b) Biofilm after 72 h formation on uncoated Si. Raman spectra are baseline corrected using a fourth order polynomial function. Vertical dashed lines added to aid comparison between spectra. Spectrum 1(a) is identical to that in the original.

Fig. 3 Composite Raman images of different *Pseudomonas aeruginosa* samples. (a) Plan view image of planktonic cells constructed from scattering in the region $2800\text{--}3050\text{ cm}^{-1}$. Scale bar $0.7\text{ }\mu\text{m}$. (b) Plan view image of a biofilm constructed from a combination of bands in the regions $2800\text{--}3050\text{ cm}^{-1}$ (red) and $980\text{--}1020\text{ cm}^{-1}$ (blue), representing all organic components and protein scattering, respectively. Scale bar $10\text{ }\mu\text{m}$. (c) Depth profile of the biofilm in the $x\text{--}z$ plane constructed from $2800\text{--}3050\text{ cm}^{-1}$ (red), $1560\text{--}1620\text{ cm}^{-1}$ (blue) and $725\text{--}775\text{ cm}^{-1}$ (green). Scale bar $7\text{ }\mu\text{m}$. Panels (a) and (c) are identical to those in the original.

The important band at 1034 cm^{-1} , as well as bands at 1129 cm^{-1} and 1163 cm^{-1} , lie in the carbohydrate region of the spectrum, and are observed in the wild-type biofilm spectrum, Fig. 1(b), but not in the spectrum from the planktonic cells, Fig. 1(a). These are classified as C–O stretching with contributions from in plane C–H deformations in phenylalanine (1034 cm^{-1}), C–C and C–O symmetric (1128 cm^{-1}) and asymmetric (1163 cm^{-1}) ring breathing vibrations.^{3–5} In the context of *Pseudomonas*-derived biofilms, these bands are consistent with the presence of rhamnolipids,⁶ a specific class of glycolipids known to be secreted by *Pseudomonas* species concurrently with biofilm formation,^{7,8} an assignment confirmed *via* MS. Comparison of representative microspectra from a *Pseudomonas* wild-type biofilm, Fig. 1(b) with a mixed rhamnolipid standard shows that the standard has contributions from common Raman bands at 1030 cm^{-1} , 1068 cm^{-1} , and 1155 cm^{-1} (wide band that includes the contributions at 1163 cm^{-1}) characteristic of the sugar moieties, as well as a strong band at 1445 cm^{-1} as described above consistent with the presence of

Table S1 Raman band assignments for *P. aeruginosa* biofilm spectrum

Frequency (cm ⁻¹)	Assignment
621	In plane ring breathing deformation in phenylalanine ¹
674	Ring breathing modes in DNA bases ⁹
704	
746	
830	Tyrosine ring breathing ^{3,10}
849	Tyrosine ring breathing ^{3,10}
879	C–C stretching, C–O–C glycosidic link ^{4,5,10}
908	C–O–C stretching ¹¹
952	
1005	Ring breathing phenylalanine and tryptophan ¹
1034	C–H in plane phenylalanine, C–C and C–O stretching ^{3–5}
1064	C–C and C–O stretching ^{3–5}
1099	C–O–C glycosidic link symmetric ring breathing ^{3–5}
1128	C–C str, C–O–C glycosidic link symmetric ring breathing ^{3–5}
1163	C–C str, C–O–C glycosidic link asymmetric ring breathing ^{3–5}
1186	Tyrosine, phenylalanine
1215	Amide III ^{3–5,10}
1246	Amide III ^{3–5,10}
1310	δ (CH) in carbohydrates and proteins ^{4,5,10}
1341	δ (CH) in carbohydrates and proteins ^{4,5,10}
1392	COO ⁻ symmetric stretch
1417	COO ⁻ symmetric stretch
1445	CH ₂ deformation ^{5,10}
1558	Tryptophan ring ¹²
1587	Ring stretch doublet – benzene derivative
1605	C=C phenylalanine ¹

ramnolipids in the biofilm matrix. Confirming the assignment of these bands to ramnolipids, the MS data not only show the presence of ramnolipids, but also allow their assignment to individual congeners.

Band assignments for the biofilm spectrum are summarized in Table S1.

References

- 1 G. J. Puppels, H. S. P. Garritsen, G. M. J. Segersnolten, F. F. M. Demul and J. Greve, *Biophys. J.*, 1991, **60**, 1046–1056.
- 2 N. Uzunbajakava, A. Lenferink, Y. Kraan, B. Willekens, G. Vrensen, J. Greve and C. Otto, *Biopolymers*, 2003, **72**, 1–9.
- 3 I. Notingher, S. Verrier, H. Romanska, A. E. Bishop, J. M. Polak and L. L. Hench, *Spectrosc. Int. J.*, 2002, **16**, 43–51.
- 4 M. Harz, P. Rosch, K. D. Peschke, O. Ronneberger, H. Burkhardt and J. Popp, *Analyst*, 2005, **130**, 1543–1550.
- 5 U. Neugebauer, U. Schmid, K. Baumann, W. Ziebuhr, S. Kozitskaya, V. Deckert, M. Schmitt and J. Popp, *ChemPhysChem*, 2007, **8**, 124–137.
- 6 M. Jadhav, S. Kalme, D. Tamboli and S. Govindwar, *J. Basic Microbiol.*, 2011, **51**, 385–396.
- 7 B. R. Boles, M. Thoendel and P. K. Singh, *Mol. Microbiol.*, 2005, **57**, 1210–1223.
- 8 M. E. Davey, N. C. Caiazza and G. A. O'Toole, *J. Bacteriol.*, 2003, **185**, 1027–1036.
- 9 J. W. Chan, D. S. Taylor, T. Zwerdling, S. M. Lane, K. Ihara and T. Huser, *Biophys. J.*, 2006, **90**, 648–656.
- 10 K. Maquelin, C. Kirschner, L. P. Choo-Smith, N. van den Braak, H. P. Endtz, D. Naumann and G. J. Puppels, *J. Microbiol. Methods*, 2002, **51**, 255–271.
- 11 J. De Gelder, K. De Gussem, P. Vandenabeele, M. Vancanneyt, P. De Vos and L. Moens, *Anal. Chim. Acta*, 2007, **603**, 167–175.
- 12 J. T. Pelton and L. R. McLean, *Anal. Biochem.*, 2000, **277**, 167–176.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.