



Cite this: *Chem. Commun.*, 2014, 50, 13334

Received 23rd June 2014,  
Accepted 11th September 2014

DOI: 10.1039/c4cc04767b


www.rsc.org/chemcomm

## On the interaction of copper(II) with disulfiram†

David J. Lewis,<sup>‡ab</sup> Parikshit Deshmukh,<sup>a</sup> Aleksander A. Tedstone,<sup>a</sup> Floriana Tuna<sup>c</sup>  
and Paul O'Brien<sup>§\*ab</sup>

In combination with copper(II) ions, disulfiram (DSF) has been reported to be a potentially potent anticancer agent based on *in vitro* results. The interaction of DSF with copper(II) chloride in solution has been studied using a range of spectroscopic techniques. There is strong evidence for the rapid formation of the bis(*N,N*-diethyl dithiocarbamato)copper(II) complex *in situ*. Kinetic experiments were used to determine rate laws for the reaction that give insight into the mechanism of the process which may help to explain the observed *in vitro* cytotoxicity.

Disulfiram (tetraethylthiuram disulfide, DSF), the two-electron oxidised dimer of diethyl dithiocarbamate (Scheme 1), is a useful pharmaceutical agent in the treatment of chronic alcoholism.<sup>1</sup> Marketed under the name *Antabuse*, the drug acts primarily by irreversible inhibition of the enzyme acetaldehyde dehydrogenase, leading to unpleasant physical feelings accompanied by nausea or vomiting when ethanol is consumed. Recently, there has been renewed interest in DSF, primarily for its use in adjuvant therapies (e.g. clinical trials gov. identifier NCT00312819). DSF is also rapidly becoming considered as a potentially convincing anti-cancer drug. Reports have emerged recently of the induction of apoptosis *in vitro* in cancer cell lines generally thought to be chemotherapy-resistant, such as *glioblastoma multiforme*, by DSF either alone or in synergy with an anticancer agent (gemcitabine).<sup>2,3</sup> However, the cytotoxic activity observed in both cases was reliant on



Scheme 1

the presence of copper(II) salts and the active species responsible for the remarkable anticancer activity was not identified. Cen *et al.* reported a similar copper dependency in the induction of apoptosis in human melanoma cells by DSF/Cu<sup>2+</sup>.<sup>4</sup> Conticello *et al.* reported the effective use of DSF or DSF-copper(II) against human haematological malignancies *in vitro*.<sup>5</sup> It is known that DSF can also be used with other transition metals to effect anti-cancer activity,<sup>6,7</sup> though copper(II) is interesting due to its natural presence *in vivo*, requiring, in theory, only treatment of a patient with DSF alone to establish a prototype chemotherapy.

Dithiocarbamates are a class of sulfur-containing bidentate chelating ligands. The ability of the diethyldithiocarbamate ligands to complex a range of transition metal ions including Cu<sup>2+</sup> is well-known.<sup>8</sup> We have previously been interested in using metal-dithiocarbamate and metal-diselenocarbamate complexes as single-source precursors<sup>9</sup> for various semiconductors for photo-voltaic applications<sup>10–16</sup> as well as for nanocrystalline semi-conducting *quantum dots*.<sup>17,18</sup> Bifunctional dithiocarbamates have been used to functionalise the surface of noble metal nanoparticles with transition metal or lanthanide(III) complexes.<sup>19,20</sup>

The electronic properties of bis(*N,N*-diethyldithiocarbamato)-copper(II) (CuL<sub>2</sub>, Scheme 1) have been investigated by electron paramagnetic resonance (EPR) spectroscopy, giving insights into its self-association.<sup>21</sup> Similarly, the photochemistry of CuL<sub>2</sub> has been explored, with the complex displaying an intense ligand-to-metal charge transfer band (LMCT), with maximum absorbance at around 430 nm, a signature of its formation.<sup>22,23</sup>

<sup>a</sup> School of Chemistry, University of Manchester, Oxford Road, M13 9PL, UK.  
E-mail: Paul.O'Brien@manchester.ac.uk; Fax: +44 (0)161 275 4616;  
Tel: +44 (0)161 275 4653

<sup>b</sup> School of Materials, University of Manchester, Oxford Road, M13 9PL, UK


<sup>c</sup> EPSRC EPR National Research Facility and Service, Photon Science Institute, University of Manchester, Oxford Road, M13 9PL, UK

† Electronic supplementary information (ESI) available: General experimental and instrumental details, synthetic procedures, electrospray mass spectrum, and initial rate plots. See DOI: 10.1039/c4cc04767b

‡ Parts of this communication were written by DJL whilst a visiting Scholar at the University of Illinois at Urbana-Champaign, The United States of America. DJL and POB would like to thank Professors Shen J. Dillon, Pascal Bellon and Robert Averback for their kind invitation.

§ We wish to thank warmly Dr. K. A. F. O'Brien for turning our attention to the copper(II)-DSF question.





It has been claimed that DSF forms CuL<sub>2</sub> species in solution upon addition of copper(II) salts. The chelating ligand, diethyldithiocarbamate, is the product of the two-electron reduction of DSF. Chen *et al.* noted a “dramatic colour change” on addition of copper(II) salts to DSF, probably due to the aforementioned LMCT transition in CuL<sub>2</sub> formed *in situ*.<sup>24</sup> Farmer and co-workers reported that the mechanism of formation of CuL<sub>2</sub> from a mixture of Cu(II) and DSF proceeded through the spontaneous decomposition of a small fraction of DSF in the presence of water to produce 30 electrons (Scheme 2, A) which then proceed to reduce 15 molecules of DSF to form the diethyldithiocarbamate, L, which then complexes copper(II) *in situ* to form CuL<sub>2</sub> in a theoretical 93% yield.<sup>4</sup>

The main evidence presented for such a mechanism was an assay based on UV-Vis absorbance spectroscopy. The work was performed in biphasic systems (soluble Cu<sup>2+</sup> and solid DSF) formed on the addition of DSF to aqueous CuCl<sub>2</sub>, requiring extraction of reaction products into chloroform prior to analysis, as well as long reaction times (*e.g.* 24 h). Here we report the direct observation of CuL<sub>2</sub> species and study the EPR properties compared to isolated CuL<sub>2</sub>. The reaction kinetics of the interaction of DSF with Cu<sup>2+</sup> ions are studied in both unbuffered and buffered (HEPES, pH 7.4) mixed-solvent solutions.

A 1:1 molar ratio of DSF and CuCl<sub>2</sub> was analysed by positive-mode electrospray ionisation mass spectrometry in order to observe directly if the dithiocarbamate-copper(II) complex, CuL<sub>2</sub>, is formed. We were able to identify the product in solution: a peak manifold was observed at *m/z* 359, consistent with CuL<sub>2</sub> (ESI<sup>+</sup>). The isotope pattern of the peak observed at *m/z* 359 confirmed the incorporation of a single copper centre as-compared to the theoretical pattern expected for protonated CuL<sub>2</sub>. Accurate mass determination of the [M+H]<sup>+</sup> peak by high-resolution mass spectrometry (HRMS, ES<sup>+</sup>) gave a value that corresponded exactly to the theoretical mass of protonated CuL<sub>2</sub> (calc. for C<sub>10</sub>H<sub>21</sub>N<sub>2</sub>S<sub>4</sub>Cu, [M+H]<sup>+</sup>: 359.9878. Found: 359.9878), thus confirming that the species analysed by this method *in situ* is indeed protonated CuL<sub>2</sub>.




Fig. 1 S-band (3.8 GHz) EPR spectra at 80 K of 95:5 THF:water glasses of (a) isolated CuL<sub>2</sub> and (c) a mixture of DSF and CuCl<sub>2</sub> in 1:1 molar ratio. The red line (b) represents the calculated EPR spectrum of CuL<sub>2</sub>, using the parameters  $g_{\parallel} = 2.071$ ;  $g_{\perp} = 2.002$ ;  $A_{\parallel} = 165$  G and  $A_{\perp} = 43$  G.

EPR spectroscopy was used to probe structural similarities between a solution of laboratory-synthesised CuL<sub>2</sub> (synthesised by the method of Jeliaskova *et al.*,<sup>22</sup> characterisation data in ESI<sup>+</sup>) and a mixture of DSF and CuCl<sub>2</sub> in 1:1 molar ratio in a 95:5 THF:water glass at 80 K (Fig. 1) in order to compare the structure of the species formed *in situ* when stoichiometric amounts of DSF and CuCl<sub>2</sub> are mixed together in a solution. We also simulated the EPR spectrum of CuL<sub>2</sub> to compare both solutions to theory. The EPR spectra of both mixtures and the simulated spectrum of CuL<sub>2</sub> revealed striking similarities, thus suggesting that the species formed when DSF is mixed with a stoichiometric amount of Cu<sup>2+</sup> is CuL<sub>2</sub>. The EPR parameters obtained from experiment are in good agreement with the published EPR data of CuL<sub>2</sub>.<sup>21</sup>

UV-Vis absorption spectroscopy in a mixed solvent system (aqueous THF, 0.7% v/v) was used to probe complex formation under steady-state and kinetic conditions. The LMCT transition of CuL<sub>2</sub>, which has an absorbance maximum at *ca.* 425 nm can be used as a spectroscopic handle to confirm the formation of CuL<sub>2</sub> species *in situ* when CuCl<sub>2</sub> and DSF are mixed in stoichiometric amounts. Titration of a solution of DSF in THF into a solution of copper(II) chloride in water demonstrated the steady growth of the LMCT absorption band, with peak maxima at 425 nm, characteristic of CuL<sub>2</sub> (ESI<sup>+</sup>). It is important that the solution remained homogeneous throughout the titration, ensuring that all the species formed were analysed. Kinetic studies of the using a 1:1 ratio of DSF:CuCl<sub>2</sub> exhibited a marked increase in the LMCT absorption at 425 nm to a plateau in around 150 min, indicating that complexation was fully complete only after this time (Fig. 2). Isosbestic points in absorbance spectra give important information regarding the relationship between two species in solution. The presence of three well-defined isosbestic points at 250 nm, 290 nm and 325 nm demonstrate that the copper(II)-bound and unbound ligand (*i.e.* Cu(II)-DSF and DSF) species are, most likely, related linearly by stoichiometry as expected.

Continuous variation plots using the LMCT electronic transition at 425 nm were used to further probe the stoichiometry of






Fig. 2 UV-Vis absorption time-course plot for a 1:1 ratio of  $\text{Cu}^{2+}$  and DSF in THF/water. Solid-headed arrows show the either the hyperchromic (upward arrows) or hypochromic (downward arrows) nature of the electronic transitions during the study. The three isosbestic points at 250, 290 and 325 nm are marked with hollow-headed arrows. Inset: the growth of the LMCT electronic transition over time, with an interval of 5 min between points.

the complex formed between  $\text{CuCl}_2$  and DSF *in situ*. A continuous variation plot (the method of Job<sup>25</sup>) of the mole fraction of DSF,  $x$ , (such that  $x = [\text{DSF}] / ([\text{Cu}^{2+}] + [\text{DSF}])$ ) plotted vs. absorption at 425 nm ( $A_{425}$ ) in HEPES buffer revealed a value of  $x$  to be  $0.61 \pm 0.07$ , interpolated *via* the most appropriate linear fits to both halves of the plot (Fig. 3 and ESI† for calculation of intersect and calculation of random error in linear regression). This means therefore that the overall stoichiometry of the system is in slight excess of DSF; a value of  $x = 0.50$  corresponds to  $\text{CuL}_2$ , thus the slight shift to  $x > 0.5$  indicates a slight stoichiometric excess of DSF in the system. This overall stoichiometry is fully consistent with the stoichiometric equations which were subsequently derived from kinetic experiments in buffered solution (*vide infra* and Scheme 2, C).

UV-Vis absorption spectroscopy allowed the determination of a rate law for the reaction in unbuffered solution using the method of initial rates.<sup>26</sup> In this method, initial rates for the reaction being studied ( $v_0$ ) are measured for reactants isolated by concentration and plots of  $\log v_0$  vs.  $\log[R]$  (where  $[R]$  is the concentration of the reactant being studied) give straight lines with slope equal to the order of the reaction with respect to reactant  $R$ , and y-intercepts equivalent to  $\log k$ , where  $k$  is the observed rate constant with respect to the isolated reactant. The rate of formation of  $\text{CuL}_2$ -type species can thus be quantified by UV-Vis spectroscopy using the absorbance of the LMCT transition, and the orders of reaction determined for both DSF and Cu. Initial rates of reaction ( $v_0$ ;  $t \leq 5$  min) were measured for a range of concentrations of DSF and  $\text{Cu}^{2+}$  in a mixed solvent system (aqueous THF, 0.4–14% *v/v* see ESI† for full details). Plots of  $\log v_0$  vs.  $\log[\text{DSF}]$  or  $\log v_0$  vs.  $\log[\text{Cu}^{2+}]$  for isolated reactants ( $N = 5$  at every point with full analysis of systematic and random error for each point; ESI†) gave linear relationships with slopes approximately equal to the order of reaction for each individual reactant; the rate is first-order with respect to DSF, but zero-order with respect to copper *i.e.*  $v = k_1[\text{DSF}][\text{Cu}^{2+}]^0$  or simply  $v = k_1[\text{DSF}]$ , with the observed rate constant for the DSF isolated reaction equal to  $1.653 \times 10^{-2} \text{ s}^{-1}$ . The rate law derived here gives great credence to the mechanism proposed by Farmer and co-workers (*vide supra*);<sup>4</sup> disproportionation




Fig. 3 Continuous variation plot of mole fraction of DSF,  $x$ , (such that  $x = [\text{DSF}] / ([\text{Cu}^{2+}] + [\text{DSF}])$ ) vs. absorption at 425 nm ( $A_{425}$ ).

of DSF in water (*i.e.* *pseudo* first-order kinetics) to release the electrons required for reduction, followed by complexation of copper(II) and acidic runaway would indeed be *pseudo* first-order with respect to DSF, and probably rate-limiting, assuming that the complexation step is relatively rapid in comparison.

In HEPES-buffered solution an interesting change in the rate law to the form  $v_{\text{buff}} = k_{\text{r,buff}}[\text{DSF}][\text{Cu}^{2+}]$  is observed (ESI†) *i.e.* the reaction becomes first-order with respect to copper(II) whilst remaining first-order with respect to DSF. The observed rate constants determined are  $k = 1.854 \times 10^{-4} \text{ mol}^{-1} \text{ dm}^3 \text{ s}^{-1}$  with respect to isolated DSF and  $k = 4.851 \times 10^{-5} \text{ mol}^{-1} \text{ dm}^3 \text{ s}^{-1}$  with respect to isolated copper. We propose that the mechanism in buffered solutions occurs *via* reduction of  $\text{Cu}^{2+}$  to  $\text{Cu}^+$  by the initial oxidation of DSF to the bitt-4<sup>2+</sup> intermediate<sup>4</sup> which is followed by a rapid two-electron reduction of another DSF molecule by two  $\text{Cu}^+$  ions (Scheme 2, B) to form  $\text{CuL}_2$  and regenerate a single  $\text{Cu}^{2+}$  ion. The rapid disproportionation of bitt-4<sup>2+</sup> intermediate to release 30 electrons plus by-products then facilitates the formation of further amounts of  $\text{CuL}_2$  *via* the two-electron reduction of DSF (Scheme 2, C). Pathways to complexation *via* the bitt-4<sup>2+</sup> intermediate have been suggested by Farmer and co-workers.<sup>4</sup> The overall stoichiometry of the reaction with respect to  $\text{Cu}^{2+}$ :DSF ratio is *ca.* 0.9:1.0, as observed in the Job plot (*vide supra*). It is likely that this may be the mechanism for the reaction of DSF with copper(II) ions under biological conditions. It is also likely that the release of copper(I) ions during the formation of  $\text{CuL}_2$  and the catastrophic decomposition of bitt-4<sup>2+</sup> could lead to massive oxidative stress on cells *in vitro* assays and result in the apoptosis observed.

In conclusion, the interaction of DSF with copper(II) ions in solution has been comprehensively studied. Mass spectrometry and EPR suggest the species formed in solution is  $\text{CuL}_2$  where  $\text{L} = (\text{S}_2\text{CNET}_2)$ . UV-Vis absorption spectroscopy titrations at steady-state demonstrate the growth of the characteristic LMCT absorbance for this species. Kinetic studies using UV-Vis absorption spectroscopy to monitor the LMCT absorption band demonstrate that the rate law is first-order with respect to DSF in unbuffered solution, which suggests a rate-limiting step



reliant only on the concentration of DSF within the mixture. The rate law changes to a first order dependence on both DSF and copper(II) ions in buffered solutions, with a mechanism for the formation of CuL<sub>2</sub> involving an oxidised ligand intermediary, bitt-4<sup>2+</sup>, and copper(I) ions. In this sense, DSF may behave non-innocently. We have, therefore, confirmed the first step of the mechanism proposed previously to this study,<sup>4</sup> which can involve a pH runaway, as well as proposing a mechanism for the altered kinetics observed in buffered solutions.

These observations potentially have serious implications regarding the potential use of DSF as a therapeutic agent *in vivo*. It is likely that in *in vitro* assays, on addition of the copper(II) ions to the media, the cells are exposed to a rapid decomposition of DSF to bitt-4<sup>2+</sup> and copper(I) ions with a catastrophic release of reactive oxygen species such as H<sub>2</sub>O<sub>2</sub> arising from Fenton chemistry, the latter which are known to cause apoptosis.<sup>27</sup> The role of reactive oxygen species in apoptosis of cancer cells treated with DSF and DSF-copper(II) cocktails has not escaped the attention of researchers though a potential explanation for this has, until now, been elusive.<sup>5</sup> The oxidation reactions suggested here, as we have seen, are likely to be relatively rapid and thus may be highly cytotoxic (produces a greater dose of H<sub>2</sub>O<sub>2</sub> over time).<sup>28</sup> Therefore, the induction of apoptosis in tumour cells by a copper(II) DSF cocktail *in vivo* is difficult to envisage as it is probably not caused by a discrete copper-DSF complex but rather is due to a reaction.

Obviously, if the product of this reaction were a therapeutic molecule there would be a candidate drug that could be delivered *in vivo* but this does not seem to be the case. The situation is rather similar to the recent case of cisplatin solvated in DMSO, where reactivity of the solvent medium may have produced misleading results from *in vitro* assays.<sup>29</sup> Care must therefore be taken when it is suggested that the anticancer activity of DSF in combination with copper(II) observed *in vitro* can be translated *in vivo*.

The authors would like to acknowledge the EPSRC National EPR Research Facility (NS/A000014/1).

## Notes and references

- 1 R. K. Fuller, L. Branchey, D. R. Brightwell, R. M. Derman, C. D. Emrick, F. L. Iber, K. E. James, R. B. Lacoursiere, K. K. Lee, I. Lowenstam, I. Maany, D. Neiderhiser, J. J. Nocks and S. Shaw, *J. Am. Med. Assoc.*, 1986, **256**, 1449–1455.
- 2 X. Guo, B. Xu, S. Pandey, E. Goessl, J. Brown, A. L. Armesilla, J. L. Darling and W. Wang, *Cancer Lett.*, 2010, **290**, 104–113.
- 3 P. Liu, S. Brown, T. Goktug, P. Channathodiyil, V. Kannappan, J. P. Hugnot, P. O. Guichet, X. Bian, A. L. Armesilla, J. L. Darling and W. Wang, *Br. J. Cancer*, 2012, **107**, 1488–1497.
- 4 D. Z. Cen, D. Brayton, B. Shahandeh, F. L. Meyskens and P. J. Farmer, *J. Med. Chem.*, 2004, **47**, 6914–6920.
- 5 C. Conticello, D. Martinetti, L. Adamo, S. Buccheri, R. Giuffrida, N. Parrinello, L. Lombardo, G. Anastasi, G. Amato, M. Cavalli, A. Chiarenza, R. De Maria, R. Giustolisi, M. Gulisano and F. Di Raimondo, *Int. J. Cancer*, 2012, **131**, 2197–2203.
- 6 E. M. Nagy, L. Ronconi, C. Nardon and D. Fregona, *Mini-Rev. Med. Chem.*, 2012, **12**, 1216–1229.
- 7 D. Buac, S. Schmitt, G. Ventro, F. R. Kona and Q. P. Dou, *Mini-Rev. Med. Chem.*, 2012, **12**, 1193–1201.
- 8 A. Hulanicki, *Talanta*, 1967, **14**, 1371–1392.
- 9 M. A. Malik, M. Afzaal and P. O'Brien, *Chem. Rev.*, 2010, **110**, 4417–4446.
- 10 K. Ramasamy, M. A. Malik and P. O'Brien, *Chem. Sci.*, 2011, **2**, 1170–1172.
- 11 K. Ramasamy, V. L. Kuznetsov, K. Gopal, M. A. Malik, J. Raftery, P. P. Edwards and P. O'Brien, *Chem. Mater.*, 2013, **25**, 266–276.
- 12 M. Afzaal, K. Ellwood, N. L. Pickett, P. O'Brien, J. Raftery and J. Waters, *J. Mater. Chem.*, 2004, **14**, 1310–1315.
- 13 M. B. Hursthouse, M. A. Malik, M. Motevalli and P. O'Brien, *Organometallics*, 1991, **10**, 730–732.
- 14 M. B. Hursthouse, M. A. Malik, M. Motevalli and P. O'Brien, *Polyhedron*, 1992, **11**, 45–48.
- 15 M. Lazell, P. O'Brien, D. J. Otway and J.-H. Park, *J. Chem. Soc., Dalton Trans.*, 2000, 4479–4486.
- 16 P. O'Brien, J. R. Walsh, I. M. Watson, L. Hart and S. R. P. Silva, *J. Cryst. Growth*, 1996, **167**, 133–142.
- 17 B. Ludolph, M. A. Malik, P. O'Brien and N. Revaprasadu, *Chem. Commun.*, 1998, 1849–1850.
- 18 M. A. Malik, N. Revaprasadu and P. O'Brien, *Chem. Mater.*, 2001, **13**, 913–920.
- 19 E. R. Knight, A. R. Cowley, G. Hogarth and J. D. E. T. Wilton-Ely, *Dalton Trans.*, 2009, 607–609.
- 20 S. Sung, H. Holmes, L. Wainwright, A. Toscani, G. J. Stasiuk, A. J. P. White, J. D. Bell and J. D. E. T. Wilton-Ely, *Inorg. Chem.*, 2014, **53**, 1989–2005.
- 21 N. D. Yordanov and D. Shopov, *J. Chem. Soc., Dalton Trans.*, 1976, 883–886.
- 22 B. G. Jeliakova and G. C. Sarova, *J. Photochem. Photobiol., A*, 1996, **97**, 5–9.
- 23 V. F. Plyusnin, A. V. Kolomeets, V. P. Grivin, S. V. Larionov and H. Lemmetyinen, *J. Phys. Chem. A*, 2011, **115**, 1763–1773.
- 24 D. Chen, Q. C. Cui, H. Yang and Q. P. Dou, *Cancer Res.*, 2006, **66**, 10425–10433.
- 25 J. S. Renny, L. L. Tomasevich, E. H. Tallmadge and D. B. Collum, *Angew. Chem., Int. Ed.*, 2013, **52**, 11998–12013.
- 26 P. W. Atkins and J. de Paula, *The Elements of Physical Chemistry*, W. H. Freeman, 5th edn, 2009.
- 27 K. D. Held, F. C. Sylvester, K. L. Hopcia and J. E. Biaglow, *Radiat. Res.*, 1996, **145**, 542–553.
- 28 K. D. Held and J. E. Biaglow, *Radiat. Res.*, 1994, **139**, 15–23.
- 29 M. D. Hall, K. A. Telma, K.-E. Chang, T. D. Lee, J. P. Madigan, J. R. Lloyd, I. S. Goldlust, J. D. Hoeschele and M. M. Gottesman, *Cancer Res.*, 2014, **74**, 3913–3922.

