Cite this: *Chem. Commun.*, 2014, 50, 10979

Received 14th July 2014,
Accepted 28th July 2014

DOI: 10.1039/c4cc05421k

www.rsc.org/chemcomm

Stepwise assembly of an adamantoid Ru_4Ag_6 cage by control of metal coordination geometry at specific sites†

Alexander J. Methereil and Michael D. Ward*

The geometrically pure 'complex ligand' $\text{fac}[\text{Ru}(\text{L}^{\text{Ph}})_3]^{2+}$, in which three pendant bidentate binding sites are located on one face of the complex, reacts with $\text{Ag}(\text{I})$ ions to form the adamantoid decanuclear cage $[\{\text{Ru}(\text{L}^{\text{Ph}})_3\}_4\text{Ag}_6](\text{PF}_6)_{14}$ which contains a 6-coordinate $\text{Ru}(\text{II})$ ion at each vertex of a large tetrahedron and a 4-coordinate $\text{Ag}(\text{I})$ ion along each edge.

The self-assembly and host-guest chemistry of metal-ligand coordination cages continue to fascinate. Such high-symmetry cages represent appealing synthetic targets for supramolecular chemists to test their skills at controlling self-assembly so as to generate elaborate, multi-component assemblies from simple starting materials.¹ In addition the ability of cages to bind guest molecules in their central cavity leads to potential applications² ranging from catalysis³ to drug delivery.⁴

The vast majority of cages are formed from just two types of component: one type of metal ion and one type of ligand. A few examples of mixed-ligand cages are known in which the self-assembly process specifically occurs with selection of two different types of ligand, resulting in a heteroleptic complex being favoured over the homoleptic alternatives.⁵ Likewise a few examples of mixed-metal cages exist in which two different types of metal ion occupy different vertex positions in a cage structure.^{6–8} This can occur when the two types of metal ion have different geometric preferences and the self-assembly requires both: for example, octahedral tris-chelate metal ions at the vertices of a cube and square planar ions with four monodentate ligands at the face centres.⁷ Alternatively, we showed recently how different types of metal ion can be positioned at specific sites in a polyhedral array if kinetically inert metal complex subcomponents

are prepared first and then combined with a second labile metal ion to complete the assembly in a stepwise manner.⁸

Our extensive family of polyhedral cage complexes generally contain an octahedral tris-chelate metal ion at each vertex, and a bis-bidentate bridging ligand (containing two pyrazolyl-pyridine chelating termini) along each edge.^{1c} In these complexes the geometric isomerism (*fac* vs. *mer*) of the metal centres turns out to play a crucial role in the nature of the assembly that forms. In some complexes, such as a family of M_4L_6 tetrahedra, all four metal centres have a *fac* tris-chelate geometry;⁹ in contrast, in a series of $\text{M}_{12}\text{L}_{18}$ truncated tetrahedra, all metal centres have a *mer* tris-chelate geometry.¹⁰ In several other types of cage assembly however there is a 3 : 1 mixture of *mer* : *fac* tris-chelate vertices.¹¹ Therefore, the ability to control the self-assembly of such cages – particularly mixed-metal versions – relies on the ability to prepare kinetically stable, geometrically pure *fac* or *mer* tris-chelate sub-components as starting points to propagate a specific assembly.

We report here the use of this principle – *viz.* control of geometric isomerism at specific sites in a cage as a way of directing assembly – in the formation of an unusual $[\text{Ru}_4\text{Ag}_6(\text{L}^{\text{Ph}})_{12}]^{14+}$ mixed-metal cage which combines octahedral *fac* tris-chelate $\text{Ru}(\text{II})$ vertices and pseudo-tetrahedral $\text{Ag}(\text{I})$ bis-chelate edges in an adamantane-type cage structure having tetrahedral symmetry. The novelty lies both in the structure of the cage [a combination of three-connected and two-connected metal vertices based on $\text{Ru}(\text{II})$ and $\text{Ag}(\text{I})$ respectively], and in the use of the pre-formed, kinetically stable $[\text{Ru}(\text{L}^{\text{Ph}})_3]^{2+}$ units as purely the *fac* isomer to direct the course of the self-assembly.

As *fac*- $[\text{ML}_3]^{2+}$ units from this family occur at specific sites in many of our cages,^{1c} we wished to start with *fac*- $[\text{Ru}(\text{L}^{\text{Ph}})_3]^{2+}$ in which the three pendant binding sites, where cage propagation occurs by coordination to additional metal ions, have a *fac* arrangement. Simple reaction of $\text{Ru}(\text{dmsO})_4\text{Cl}_2$ with > 3 equivalents of L^{Ph} afforded $[\text{Ru}(\text{L}^{\text{Ph}})_3]^{2+}$ as a 1 : 3 statistical mixture of *fac* : *mer* isomers as shown by the ^1H NMR spectrum in which every type of proton (*e.g.* coordinated pyridyl H^6) occurred in four different environments in a 1 : 1 : 1 : 1 ratio. However column chromatography or HPLC under a range of conditions did not give a good separation of the geometric isomers.

Department of Chemistry, University of Sheffield, Sheffield S3 7HF, UK.

E-mail: m.d.ward@sheffield.ac.uk

† Electronic supplementary information (ESI) available: Crystallographic data in CIF format; details of crystallographic data collections and refinements; selected bond distances for the structures; synthesis and characterisation of new complexes, including ^1H NMR, COSY and DOSY spectra; additional figures of the cage $[\{\text{Ru}(\text{L}^{\text{Ph}})_3\}_4\text{Ag}_6](\text{PF}_6)_{14}$. CCDC 1013864 and 1013865. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c4cc05421k

Scheme 1 Preparation of fac -[Ru(L^{Ph})₃](PF₆)₂.

We therefore went back a step in the synthesis, to fac -[Ru(PyPzH)₃]²⁺ [Scheme 1; PyPzH = 3-(2-pyridyl)-pyrazole] which can be readily separated from the *mer* isomer using a method reported earlier.¹² In fac -[Ru(PyPzH)₃]²⁺ the three pyrazole rings, with their acidic NH protons, lie of course on the same face of the complex. Alkylation of these^{12a} with the bromomethyl compound **A** completed the formation of the L^{Ph} ligands coordinated to the metal centre at one end, to give fac -[Ru(L^{Ph})₃]²⁺ which was isolated as the hexafluorophosphate salt (see ESI[†]). The ¹H NMR spectrum showed 20 ¹H environments confirming the threefold symmetry with all three ligands equivalent. Notably the CH₂ protons close to the Ru(II) chiral centre are diastereotopic, giving a coupled pair of doublets at 5.5 and 4.8 ppm, whereas the CH₂ protons more remote from the Ru(II) centre give a singlet at 5.3 ppm (Fig. S3, ESI[†]). The crystal structure of the complex cation of fac -[Ru(L^{Ph})₃](PF₆)₂·acetone is shown in Fig. 1; (ESI[†]) it is clear how the three pendant pyrazolyl-pyridine arms are directed to the same face of the complex. The phenyl group of each pendant arm forms a π -stacking interaction with a coordinated pyrazolyl-pyridine group from another ligand, as we have observed in related complexes.

Ag(I) generally forms four-coordinate bis-chelate complexes with pyrazolyl-pyridine ligands of this type.¹³ On the basis that three pendant bidentate sites are available for coordination in fac -[Ru(L^{Ph})₃](PF₆)₂, we combined (ESI[†]) fac -[Ru(L^{Ph})₃](PF₆)₂ with 1.5 equivalents of AgPF₆ to maximise the likelihood of a structure forming that conforms to the principle of maximum site occupancy, with all metal ions coordinatively saturated and

all ligands fully coordinated.¹⁴ If each pendant ligand fragment coordinates to a different Ag(I) ion, as is likely on steric grounds given the distance between the pendant pyrazolyl-pyridine units, we expect a mixed-metal cage in which each fac -[Ru(L^{Ph})₃]²⁺ unit caps a triangular array of Ag(I) ions.

Slow crystallisation of the reaction mixture afforded X-ray quality crystals of what proved to be a decanuclear Ru₄Ag₆ cage [Ru(L^{Ph})₃]₄Ag₆(PF₆)₁₄ (Fig. 2–4) (ESI[†]). The cage has an adamantane-like structure, with a Ru(II) tris-chelate unit at each of the four three-connected vertices which are arranged in an approximate tetrahedron. An Ag(I) bis-chelate unit occupies each of the six two-connected vertices. Thus the structure can be described as a tetrahedral array of Ru(II) ions with an Ag(I) ion lying in the centre of each Ru···Ru edge (Fig. 2), with every adjacent Ru(II)–Ag(I) pair connected by a bis-bidentate bridging ligand L^{Ph}.

The molecule lies astride a crystallographic C₂ axis such that half of it is unique. This axis passes through Ag(2) and Ag(3) such that these lie on special positions with 50% occupancy in the asymmetric unit, whereas Ag(1) and Ag(4) are in general positions. There is a (non-crystallographic) C₃ axis through each Ru(II) tris-chelate vertex, with all four being homochiral; thus the complex belongs to the pure rotation symmetry point group *T* which is a common consequence of removing mirror planes from high-symmetry polyhedra.

Fig. 1 Structure of the complex cation of fac -[Ru(L^{Ph})₃](PF₆)₂·acetone with the three ligands coloured differently for clarity.

Fig. 2 Two views of the structure of [Ru(L^{Ph})₃]₄Ag₆(PF₆)₁₄: (a) the adamantane-like arrangement of metal ions, with the four anions that lie within the cavity also shown; (b) the metal superstructure with three of the bridging ligands included (coloured differently for clarity).

Fig. 3 A view of the complete complex cation of $[(\text{Ru}(\text{L}^{\text{Ph}})_3)_4\text{Ag}_6](\text{PF}_6)_{14}$. The two ligands coordinated to each $\text{Ag}(\text{I})$ have the same colour. Labels A and B denote the electron-deficient (pyrazolyl-pyridine) and electron-rich (phenyl) units involved in the pairwise π -stacking interactions.

Fig. 4 Space-filling views of the complex cation of $[(\text{Ru}(\text{L}^{\text{Ph}})_3)_4\text{Ag}_6](\text{PF}_6)_{14}$. (a) A view down one of the threefold axes, through a $\text{Ru}(\text{II})$ tris-chelate centre; (b) a view from the opposite side of the complex looking at one of the Ru_3Ag_3 faces, with one of the encapsulated $[\text{PF}_6]^-$ anions (F atoms in green) visible through the portal.

The six $\text{Ag}(\text{I})$ ions lie on the three C_2 axes associated with T symmetry of which one [the $\text{Ag}(\text{2}) \cdots \text{Ag}(\text{3})$ axis, as mentioned above] occurs in the crystal structure; necessarily, all six $\text{Ag}(\text{I})$ ions have the same chirality associated with their two non-symmetrical chelating ligands. The nearest-neighbour $\text{Ru} \cdots \text{Ag}$ separations (*i.e.* along an edge spanned by a bridging ligand) lie in the range 8.86–9.32 Å, averaging 9.06 Å.

The flexibility of the ligands associated with the CH_2 ‘hinges’ allows them to adopt a conformation which maximises inter-ligand π -stacking – a key driver for assembly of such cages.^{16,15} This can be seen in the view shown in Fig. 3, in which the octahedral disposition of the six $\text{Ag}(\text{I})$ ions is emphasised with these being placed top/bottom, left/right and front/back with each pair of $\text{Ag}(\text{I})$ ions lying on a C_2 axis. In this view, $\text{Ag}(\text{4})/\text{Ag}(\text{4A})$ form the ‘vertical’ C_2 axis. The two ligands attached to each $\text{Ag}(\text{I})$ ion have the same colour (*i.e.* the twelve ligands are coloured in six sets of two). The ligands

are disposed such that a central phenyl ring of a bridging ligand (denoted ‘B’ in Fig. 3) lies parallel to, and overlapping with, a pyrazolyl-pyridine unit of another ligand coordinated to the adjacent $\text{Ru}(\text{II})$ ion (denoted ‘A’ in Fig. 3), forming a charge-assisted π -stack between electron-rich (phenyl) and electron-deficient (coordinated pyrazolyl-pyridine) ligands. In the view in Fig. 3 we can readily see four such A/B stacked pairs; there are necessarily, therefore, twelve such interactions overall – involving every phenyl group – as the orientations with $\text{Ag}(\text{1})/\text{Ag}(\text{1A})$ and $\text{Ag}(\text{2})/\text{Ag}(\text{2A})$ as the ‘vertical’ axis are equivalent.

An alternative space-filling view, looking down one of the C_3 axes associated with a $\text{Ru}(\text{II})$ tris-chelate centre, is in Fig. 4a. The cage complex has an approximate cavity size of 178 Å³ (calculated assuming that the windows are blocked; Fig. S2, ESI[†]). The cavity is occupied by a tetrahedral array of four $[\text{PF}_6]^-$ anions (Fig. 2a), each one blocking the window in one of the Ru_3Ag_3 faces of the cage, as shown in Fig. 4b in which three of the F atoms of the $[\text{PF}_6]^-$ anion in that window can be clearly seen. The P $\cdots$ P separations between the four encapsulated anions are in the range of 5.44–5.61 Å, resulting in peripheral F $\cdots$ F contacts between anions of ≈ 3 Å, which is the sum of the van der Waals’ radii of two F atoms. Each anion is involved in a range of $\text{CH} \cdots \text{F}$ interactions with ligand H atoms. Fig. S1(a) (ESI[†]) shows one of the anions embedded in the window in one of the Ru_3Ag_3 faces, with dotted lines indicating some of the short non-bonded $\text{C} \cdots \text{F}$ contacts (≤ 3.15 Å) which are indicative of weak hydrogen-bonding interactions between anion and ligand. This view also nicely shows how the array of six ligands around each Ru_3Ag_3 face forms a cyclic helicate with every ligand in the cycle having the same sense of ‘under and over’ around the ring. Fig. S1(b) (ESI[†]) shows how the four anions fill the cavity.

The structural integrity of the complex in solution was confirmed by ES mass spectrometry, which showed peaks corresponding to the species $[(\text{Ru}_4\text{Ag}_6(\text{L}^{\text{Ph}})_{12})(\text{PF}_6)_{14-n}]^{n+}$ ($n = 3, 4, 6$), and also by ^1H NMR spectroscopy (ESI[†]). The ^1H NMR spectrum at room temperature is very broad, indicative of molecular motions [possibly associated with the highly flexible $\text{Ag}(\text{I})$ centres] at a rate comparable to the ^1H NMR timescale. However at 75 °C the spectrum sharpened satisfactorily and showed the expected 20 independent ^1H signals associated with one environment for L^{Ph} with no internal symmetry (Fig. S6, ESI[†]); this spectrum is considerably different from that of *fac*- $[\text{Ru}(\text{L}^{\text{Ph}})_3](\text{PF}_6)_2$. Significantly, the chirality associated with the $\{\text{Ag}(\text{NN})_2\}^+$ centres ensures that *both* independent sets of CH_2 protons are now diastereotopic, giving two pairs of coupled doublets in the 4.5–5.5 ppm region (Fig. S5, ESI[†]). That this species is a large assembly is confirmed by its DOSY spectrum which clearly shows that all of its ^1H signals belong to a single species which has a much lower diffusion rate [$\log D$ ($\text{m}^2 \text{s}^{-1}$) = -9.2] than *fac*- $[\text{Ru}(\text{L}^{\text{Ph}})_3](\text{PF}_6)_2$ [$\log D$ ($\text{m}^2 \text{s}^{-1}$) = -8.4] (Fig. S7, ESI[†]).

Assembly of this cage with its adamantane-like structure thus relies on two different types of geometric control at specific metal sites. Firstly it requires the appropriate combination of metal vertices that are three-connected [each tris-chelate, $\text{Ru}(\text{II})$ ion is connected to three $\text{Ag}(\text{I})$ ions] and two connected [each bis-chelate $\text{Ag}(\text{I})$ ion is connected to two $\text{Ru}(\text{II})$ ions]. This is achieved by using metal ions with different stereoelectronic preference, *i.e.* a combination of

6-coordinate Ru(II) and 4-coordinate Ag(I) ions at alternating sites. Secondly, the structure relies on exclusive use of pre-formed, kinetically inert *fac* isomers of the $[\text{Ru}(\text{L}^{\text{ph}})_3]^{2+}$ unit. We note that there are a few other examples of mixed-metal $\text{M}_6\text{M}'_4(\mu\text{-L})_{10}$ complexes with an adamantane-like core structure.^{16,17} Many of these arise from one-pot reactions, but some – which use cyanide bridges along the M–M' edges – are based on a kinetically stable, pre-formed hexacyanometallate unit as one precursor in a manner related to ours.¹⁷ In principle the method we have reported here should have high generality as it could be extended to other kinetically inert octahedral d^6 metal complexes of the type *fac*- $[\text{M}(\text{PyPzH})_3]^{n+}$ (M = Os, $n = 2$; M = Rh, Ir, $n = 3$ etc.).

We thank EPSRC for financial support, Mr Will Cullen for assistance with the NMR measurements, and the EPSRC National Crystallography Service for the crystallographic data collections.

Notes and references

- (a) D. Fiedler, D. H. Leung, R. G. Bergman and K. N. Raymond, *Acc. Chem. Res.*, 2005, **38**, 349; (b) M. Fujita, M. Tominaga, A. Hori and B. Therrien, *Acc. Chem. Res.*, 2005, **38**, 369; (c) M. D. Ward, *Chem. Commun.*, 2009, 4487; (d) J. J. Perry, J. A. Perman and M. J. Zaworotko, *Chem. Soc. Rev.*, 2009, **38**, 1400; (e) H. Amouri, C. Desmarests and J. Moussa, *Chem. Rev.*, 2012, **112**, 2015; (f) A. F. Williams, *Coord. Chem. Rev.*, 2011, **255**, 2104; (g) Z. Laughrey and B. Gibb, *Chem. Soc. Rev.*, 2011, **40**, 363; (h) P. Jin, S. J. Dalgarno and J. L. Atwood, *Coord. Chem. Rev.*, 2012, **254**, 1760; (i) R. J. Chakrabarty, P. S. Mukherjee and P. J. Stang, *Chem. Rev.*, 2011, **111**, 6810; (j) Y. Inokuma, M. Kawano and M. Fujita, *Nat. Chem.*, 2011, **3**, 349; (k) M. D. Pluth, R. G. Bergman and K. N. Raymond, *Acc. Chem. Res.*, 2009, **42**, 1650; (l) M. M. J. Smulders, I. A. Riddell, C. Browne and J. R. Nitschke, *Chem. Soc. Rev.*, 2013, **42**, 1728; (m) T. Nakamura, H. Ube and M. Shionoya, *Chem. Lett.*, 2014, **42**, 328.
- M. D. Ward and P. R. Raithby, *Chem. Soc. Rev.*, 2013, **42**, 1619.
- (a) C. J. Brown, R. G. Bergman and K. N. Raymond, *J. Am. Chem. Soc.*, 2009, **131**, 17530; (b) C. J. Hastings, M. D. Pluth, R. G. Bergman and K. N. Raymond, *J. Am. Chem. Soc.*, 2010, **132**, 6938; (c) J. L. Bolliger, A. M. Belenguer and J. R. Nitschke, *Angew. Chem., Int. Ed.*, 2013, **52**, 7958.
- (a) J. W. Yi, N. P. E. Barry, M. A. Furrer, O. Zava, P. J. Dyson, B. Therrien and B. H. Kim, *Bioconjugate Chem.*, 2012, **23**, 461; (b) B. Therrien, *Chem. – Eur. J.*, 2013, **19**, 8378; (c) J. E. M. Lewis, E. L. Gavey, S. A. Cameron and J. D. Crowley, *Chem. Sci.*, 2012, **3**, 778.
- (a) N. K. Al-Rasbi, I. Tidmarsh, S. P. Argent, H. Adams, L. P. Harding and M. D. Ward, *J. Am. Chem. Soc.*, 2008, **130**, 11641; (b) S. Hiraoka, Y. Kubota and M. Fujita, *Chem. Commun.*, 2000, 1509; (c) S. Hiraoka, M. Shiro and M. Shionoya, *J. Am. Chem. Soc.*, 2004, **126**, 1214; (d) P. N. W. Baxter, J.-M. Lehn, G. Baum and D. Fenske, *Chem. – Eur. J.*, 1999, **5**, 102; (e) B. Olenyuk, J. A. Whiteford, A. Fechtenkötter and P. J. Stang, *Nature*, 1999, **398**, 796; (f) M. Wang, Y.-R. Zheng, K. Ghosh and P. J. Stang, *J. Am. Chem. Soc.*, 2010, **132**, 6282; (g) J.-R. Li and H.-C. Zhou, *Angew. Chem., Int. Ed.*, 2009, **48**, 8465.
- (a) W. J. Ramsay, T. K. Ronson, J. K. Clegg and J. R. Nitschke, *Angew. Chem., Int. Ed.*, 2013, **52**, 13439; (b) X. Sun, D. W. Johnson, D. L. Caulder, K. N. Raymond and E. H. Wong, *J. Am. Chem. Soc.*, 2001, **123**, 2752; (c) F. E. Hahn, M. Offermann, C. Schulzelsfort, T. Pape and R. Frohlich, *Angew. Chem., Int. Ed.*, 2008, **47**, 6794; (d) S. Hiraoka, Y. Sakata and M. Shionoya, *J. Am. Chem. Soc.*, 2008, **130**, 10058; (e) H.-B. Wu and Q.-M. Wang, *Angew. Chem., Int. Ed.*, 2009, **48**, 7343; (f) A. J. Metherell and M. D. Ward, *RSC Adv.*, 2013, **3**, 14281.
- (a) M. L. Saha and M. Schmittle, *J. Am. Chem. Soc.*, 2013, **135**, 17743; (b) M. M. J. Smulders, A. Jimenez and J. R. Nitschke, *Angew. Chem., Int. Ed.*, 2012, **51**, 6681; (c) K. Li, L.-Y. Zhang, C. Yan, M. Pan, L. Zhang and C.-Y. Su, *J. Am. Chem. Soc.*, 2014, **136**, 4456; (d) F. Reichel, J. K. Clegg, K. Gloe, K. Gloe, J. J. Weigand, J. K. Reynolds, C.-G. Li, J. R. Aldrich-Wright, C. J. Kepert, L. F. Lindoy, H.-C. Yao and F. Li, *Inorg. Chem.*, 2014, **53**, 688; (e) M. Otte, P. F. Kuijpers, O. Troeppner, I. Ivanović-Burmazović, J. N. H. Reek and B. de Bruin, *Chem. – Eur. J.*, 2013, **19**, 10170; (f) A. Galstyan, P. J. Sanz Miguel, K. Weise and B. Lippert, *Dalton Trans.*, 2013, **42**, 16151; (g) P. de Wolf, S. L. Heath and J. A. Thomas, *Chem. Commun.*, 2002, 2540.
- A. J. Metherell and M. D. Ward, *Chem. Commun.*, 2014, **50**, 6330.
- (a) J. S. Fleming, K. L. V. Mann, C.-A. Carraz, E. Psillakis, J. C. Jeffery, J. A. McCleverty and M. D. Ward, *Angew. Chem., Int. Ed.*, 1998, **37**, 1279; (b) R. L. Paul, Z. R. Bell, J. C. Jeffery, J. A. McCleverty and M. D. Ward, *Proc. Natl. Acad. Sci. U. S. A.*, 2002, **99**, 4883; (c) I. S. Tidmarsh, B. F. Taylor, M. J. Hardie, L. Russo, W. Clegg and M. D. Ward, *New J. Chem.*, 2009, **33**, 366.
- (a) Z. R. Bell, J. C. Jeffery, J. A. McCleverty and M. D. Ward, *Angew. Chem., Int. Ed.*, 2002, **41**, 2515; (b) S. P. Argent, H. Adams, T. Riis-Johannessen, J. C. Jeffery, L. P. Harding, O. Mamula and M. D. Ward, *Inorg. Chem.*, 2006, **45**, 3905.
- (a) I. S. Tidmarsh, T. B. Faust, H. Adams, L. P. Harding, L. Russo, W. Clegg and M. D. Ward, *J. Am. Chem. Soc.*, 2008, **130**, 15167; (b) A. Stephenson, S. P. Argent, T. Riis-Johannessen, I. S. Tidmarsh and M. D. Ward, *J. Am. Chem. Soc.*, 2011, **133**, 858; (c) B. R. Hall, L. E. Manck, I. S. Tidmarsh, A. Stephenson, B. F. Taylor, E. J. Blaikie, D. A. Vander Griend and M. D. Ward, *Dalton Trans.*, 2011, **40**, 12132.
- (a) A. J. Metherell, W. Cullen, A. Stephenson, C. A. Hunter and M. D. Ward, *Dalton Trans.*, 2014, **43**, 71; (b) M. H. W. Lam, S. T. C. Cheung, K.-M. Fung and W.-T. Wong, *Inorg. Chem.*, 1997, **36**, 4618.
- (a) A. Stephenson and M. D. Ward, *Chem. Commun.*, 2012, **48**, 3605; (b) A. Stephenson and M. D. Ward, *RSC Adv.*, 2012, **2**, 10844.
- J.-M. Lehn and A. V. Eliseev, *Science*, 2001, **291**, 2331.
- A. Stephenson, D. Sykes and M. D. Ward, *Dalton Trans.*, 2013, **42**, 6756.
- (a) L. Wang, Y. Li, Y. Peng, Z. Liang, J. Yu and R. Xu, *Dalton Trans.*, 2012, **41**, 6242; (b) X. Kuang, X. Wu, R. Yu, J. P. Donahue, J. Huang and C.-Z. Lu, *Nat. Chem.*, 2010, **2**, 461.
- (a) T. Shiga, F. Iijima, T. Tetsuka, G. N. Newton and H. Oshio, *Macromol. Symp.*, 2012, **317–318**, 286; (b) T. Shiga, G. N. Newton, J. S. Mathieson, T. Tetsuka, M. Nihei, L. Cronin and H. Oshio, *Dalton Trans.*, 2010, **39**, 4730; (c) T. Shiga, T. Tetsuka, K. Sakai, Y. Sekine, M. Nihei, G. N. Newton and H. Oshio, *Inorg. Chem.*, 2014, **53**, 5899.