


ChemComm

Chemical Communications

www.rsc.org/chemcomm



ISSN 1359-7345



COMMUNICATION

Urs F. Fritze and Max von Delius

Dynamic disulfide metathesis induced by ultrasound

175 YEARS



Cite this: *Chem. Commun.*, 2016, 52, 6363

Received 7th March 2016,
Accepted 29th March 2016

DOI: 10.1039/c6cc02034h

www.rsc.org/chemcomm

Dynamic disulfide metathesis induced by ultrasound†

Urs F. Fritze and Max von Delius*

The reversible metathesis of disulfide bonds is generally induced by a combination of a reducing agent and base or by irradiation with ultraviolet light. Here we report that ultrasound irradiation is suitable for generating clean equilibrium mixtures of disulfides within one hour or one day, depending on the sonication source. Preliminary mechanistic investigations suggest that the solvent plays an active role in producing initiator radicals.

The dynamic chemistry of disulfide bonds plays a vital part in cysteine-containing proteins,¹ and since 2000² it has emerged as one of the most powerful tools in the field of dynamic covalent/combinatorial chemistry (DCC).³

In nature, as well as in most man-made systems, disulfide exchange proceeds by the nucleophilic attack of a free thiolate at the disulfide bond, furnishing a new pair of disulfide and thiolate (Fig. 1a).⁴ When starting from air-stable disulfides, this type of reaction is somewhat limited in scope, because it requires long equilibration times and a combination of a reducing agent and base, which is not always compatible with the application one has in mind for a dynamic library. Progress towards alleviating these limitations was recently reported by several groups. Ramström and coworkers developed a Brønsted base-free, phosphine-catalyzed method for disulfide metathesis,^{5a} while Belenguer, Friščić and Sanders described a mechanochemical approach that requires a base, but no reducing agent.^{5b,c} The group of Pittelkow found that diselenides readily exchange at neutral pH, which in turn can be used to catalyze disulfide exchange.^{5d}

An alternative approach for facilitating disulfide metathesis is based on the homolytic cleavage of the disulfide bond and a subsequent radical chain reaction (Fig. 1b). For disulfides, this process can be induced by irradiation with harsh ultraviolet




Fig. 1 Disulfide metathesis: (a and b) conventional methods; (c) ultrasound-induced.

(UV) light,⁶ but for diselenides Xu and coworkers have recently demonstrated that visible light is suitable for generating sufficient initiator radicals for an efficient exchange reaction.^{7a} Interestingly, the authors were able to use this process for a spatially controlled “healing” of damaged areas in diselenide-based polymers.^{7b} These studies illustrate that unprecedented methods for generating dynamic libraries ultimately translate into new applications of dynamic systems.

Here we report that the metathesis of two or more disulfides can be induced by simple sonication of a chloroform solution of the starting materials (Fig. 1c). While the use of a sonotrode leads to remarkably short equilibration times, it is also possible to carry out the reaction in an ultrasonic bath, a piece of equipment that is available in most chemical laboratories.

Researchers in the field of polymer mechanochemistry⁸ routinely make use of intense ultrasound for transmitting mechanical force onto functional groups (mechanophores). This process is not applicable to small molecules, however, because it is based on cavitation shear forces that only affect macromolecules ($M_w > 30$ kDa). Against this background, we were surprised when we discovered that small-molecule disulfides readily exchanged when subjected to ultrasound.

As shown in Fig. 2, irradiation of benzylic disulfides **AA** and **BB** for one hour with ultrasound (sonotrode) furnished mixed

Friedrich-Alexander University Erlangen-Nürnberg (FAU), Department of Chemistry and Pharmacy & Interdisciplinary Center of Molecular Materials (ICMM), Henkestrasse 42, 91054 Erlangen, Germany. E-mail: max.vondelius@fau.de

† Electronic supplementary information (ESI) available: General experimental procedures and spectroscopic data. See DOI: 10.1039/c6cc02034h






Fig. 2 (a) Ultrasound-induced disulfide metathesis. Reaction conditions: **AA** (10 mM) and **BB** (10 mM), sonotrode irradiation (25% amplitude, pulse sequence 0.5 s on, 1.0 s off), CHCl_3 , 0 °C. (b) Analytical HPLC (70% → 100% gradient acetonitrile/water, 0.3 mL min⁻¹, Ascentis amide column) of starting materials **AA** and **BB**. (c) Analytical HPLC after 30 minutes of sonication. (d) Analytical HPLC after 60 minutes of sonication.

disulfide **AB** in a statistical mixture with **AA** and **BB**. We confirmed that the reaction outcome corresponds indeed to an equilibrium distribution by also using the established methods for the same reaction (Table 1, entries 6 and 7).

Intrigued by this initial finding, we proceeded to investigate a range of reaction parameters for the reaction of **AA** with **BB**. A screening of various organic solvents (Table 1, entry 3 and the ESI†) revealed that chloroform is uniquely suited for this

reaction: even similar chlorinated solvents, such as dichloromethane or 1,1,2,2-tetrachloroethane, failed to produce significant quantities of product **AB**. Only bromoform (CHBr_3) could be identified as a second excellent, albeit more toxic, solvent for the reaction (entry 2), providing a first indication that a radical mechanism might be at hand.† The use of an entirely different solvent (tetrahydrofuran), an important aspect regarding possible applications of this method, turned out to be possible when a stoichiometric amount of additive iodoform (CHI_3) was added to the reaction mixture (entry 4).

Because sonotrodes are not widely available in chemical laboratories, we tested whether the reaction would also take place in an ultrasonic bath. We were pleased to find that a clean equilibrium mixture could be obtained after irradiating a sealed flask in a temperature-controlled ultrasonic bath for 24 hours (entry 5). To make sure that this result was not due to the make-up of one particular piece of equipment, we tested three different ultrasonic baths from two manufacturers and obtained identical reaction outcomes in all three cases (see the ESI†).

In addition to the conventional protocols for disulfide metathesis (Table 1, entries 6 and 7), we conducted three control experiments (entries 8–10), designed to provide qualitative evidence for either a thermal (“hot spots”) or a radical-initiated reaction mechanism. Heating the reaction mixture at 110 °C in a sealed tube or at 170 °C under microwave irradiation led to only negligible formation of product **AB**, whereas the use of radical initiator DBPO at 120 °C proved to be effective, pointing towards a radical mechanism.

Studies on the scope of the described ultrasound-induced disulfide exchange are summarized in Table 2. Both benzylic and aliphatic compounds are suitable substrates (entries 1, 3 and 4), whereas aromatic disulfides react only sluggishly (entry 2). This reactivity trend is complementary to the classic base-induced method, in which aromatic thiols/disulfides are most commonly used.³ Our preliminary studies confirm that ether and alcohol functions are tolerated (substrates **BB** and **EE**), whereas a Boc-protected cystine substrate (substrate **FF**, entry 5) led to the formation of several side products, indicating that this protocol is not suitable for peptide substrates. Finally, a ternary mixture of substrates **AA**, **BB** and **EE** was found to lead to clean exchange (entry 6).

To gain a deeper mechanistic understanding, we monitored the reaction between **AA** and **BB** under different reaction conditions (Fig. 3). The kinetic profiles shown in Fig. 3a provide two insights: (i) the formation of product **AB** follows a sigmoidal curve, indicating the presence of an induction period. (ii) The reaction rate is the highest for the most dilute reaction.§ Both these observations can be explained by a radical chain reaction whose initiator radicals are provided *via* sonolysis of the solvent. This rationale is strongly supported by a study of Riesz that described the sonochemical generation and subsequent trapping of $\bullet\text{CHCl}_2$ radicals derived from chloroform.⁹

Further evidence for this mechanistic scenario is provided by the data shown in Fig. 3b. As one would expect for a radical chain reaction that has proceeded beyond the initial induction period, switching the ultrasound source off after 20 minutes

Table 1 Reaction conditions for inducing the metathesis of disulfides **AA** and **BB**

AA + BB		conditions		AB	
Entry	Method	Conc. [mM]	Solvent	Time ^a	AB ^b [%]
1	Sonotrode ^c	10	CHCl ₃	1 h	48
2	Sonotrode ^c	10	CHBr ₃	45 s	50
3	Sonotrode ^c	10	Other ^d	1 h	<5 ^e
4	CHI ₃ additive (100%)	10	THF	1 h ^f	49
5	Ultrasonic bath ^g	10	CHCl ₃	24 h	53
6	10% DBU/DTT ^h	10	CHCl ₃	6 d	49
7	UV light ⁱ	10	CHCl ₃	3 h	49
8	110 °C	10	CHCl ₃	5 h	— ^e
9	Microwave, 170 °C ^j	10	CHCl ₃	1 h	<5
10	5% DBPO, 120 °C ^k	10	CHCl ₃	5 h	47

^a Minimum equilibration times. ^b Reaction progress was monitored by analytical HPLC (area percent). ^c 13 mm titanium tip, pulsing at 20 kHz with 25% of maximal amplitude, 0.5 s pulse and 1.0 s break, 0 °C (ice bath). ^d Investigated solvents: acetone, tetrahydrofuran (THF), acetonitrile, dichloromethane, 1,1,2,2-tetrachloroethane, chloroform/water (emulsion). ^e **AB** not detected. ^f 1 h sonotrode irradiation, 44 h equilibration at RT. ^g Reaction conducted in a sealed glass tube, non-stop sonication, 27 °C. ^h DBU: 1,8-diazabicyclo[5.4.0]undec-7-ene; DTT: 1,4-dithio-D-threitol. ⁱ 400 W Hg lamp. ^j 300 W, 170 °C, 11 bar. ^k DBPO: dibenzoyl peroxide.



Table 2 Scope of disulfide starting materials

$R^1S-S-R^1 + R^2S-S-R^2 \xrightleftharpoons[\text{CHCl}_3, 0^\circ\text{C}]{\text{sonotrode}} R^1S-S-R^2$			
Entry	Substrates ^a	Reaction outcome ^b	Equilibration time
1	AA + BB	Clean exchange (HPLC)	1 h (24 h) ^c
2	BB + CC	Sluggish reactivity (HPLC)	1 h
3	BB + DD	Clean exchange (NMR)	3.5 h (22 h) ^c
4	BB + EE	Clean exchange (NMR)	1 h (17 h) ^c
5	BB + FF	Side product formation (HPLC)	3 h
6	AA + BB + EE	Clean exchange (HPLC), formation of 6 compounds	1.5 h

^a Reaction conditions: equimolar quantities of the substrates (10 mM in CHCl₃), Suslick cell under an Ar atmosphere, sonotrode irradiation as specified in Table 1. ^b All reactions were monitored by HPLC(MS) or NMR spectroscopy (see the ESI). ^c Values in parentheses correspond to reactions under “ultrasonic bath” conditions.




Fig. 3 HPLC monitoring of the metathesis of disulfides **AA** and **BB** under sonotrode conditions in chloroform. (a) Influence of starting material concentration (2 mM, 10 mM, 50 mM). (b) Influence of switching off the ultrasound source (period: 20–100 min) in argon, air or in argon with addition of 40 mM of radical scavenger TEMPO ((2,2,6,6-tetramethylpiperidin-1-yl)oxyl) after 20 min.

does not stop the reaction. Whether the reaction is carried out in argon or air does not significantly affect the rate while ultrasound irradiation generates sufficient radicals, but during the “resting period” (Fig. 3b, 20–100 minutes) dioxygen supposedly leads to quenching of persistent radicals, thus slowing down the reaction. Addition of stoichiometric radical scavenger TEMPO

after 20 minutes inhibits the reaction, both in the presence and absence of ultrasound irradiation, providing further support for a radical mechanism.¹⁰

In conclusion, we have discovered a sonochemical method for generating dynamic mixtures of disulfides. Although this reaction is somewhat limited regarding solvents and substrates, we believe that it could be a practical alternative to the existing methods, particularly in scenarios where short equilibration times, base-free reaction conditions or a new vector for dynamic covalent orthogonality¹¹ are needed.

This work was supported by the Deutsche Forschungsgemeinschaft (Emmy-Noether grant DE1830/2-1) and the Evangelisches Studienwerk Villigst (PhD fellowship to U. F. F.). We thank Dr Konstantin Amsharov for providing access to a photoreactor.

Notes and references

- ‡ In CHBr₃, significant formation of side products was observed after extended reaction times (e.g. one hour), indicating that the reaction should be terminated after equilibrium is reached. In contrast to CHCl₃, CHBr₃ was not a suitable additive in solvent THF (see the ESI,† page S30).
- § We found that the reaction kinetics vary to some extent between different batches of solvent. However, the sigmoidal shape of the curves and the general trends remain unaffected.
- ¶ We were unable to identify possible adducts of $\cdot\text{CHCl}_2$ radicals to the starting materials, which could be due to the fact that only small quantities of such compounds are formed.

- (a) C. B. Anfinsen, *Science*, 1973, **181**, 223; (b) P. Dopieralski, J. Ribas-Arino, P. Anjukandi, M. Krupicka, J. Kiss and D. Marx, *Nat. Chem.*, 2013, **5**, 685.
- S. Otto, R. L. E. Furlan and J. K. M. Sanders, *J. Am. Chem. Soc.*, 2000, **122**, 12063.
- (a) P. T. Corbett, J. Leclaire, L. Vial, K. R. West, J.-L. Wietor, J. K. M. Sanders and S. Otto, *Chem. Rev.*, 2006, **106**, 3652; (b) Y. Jin, C. Yu, R. J. Denman and W. Zhang, *Chem. Soc. Rev.*, 2013, **42**, 6634; (c) S. P. Black, J. K. M. Sanders and A. R. Stefankiewicz, *Chem. Soc. Rev.*, 2014, **43**, 1861.
- R. Singh and G. M. Whitesides, *J. Am. Chem. Soc.*, 1990, **112**, 1190.
- (a) R. Caraballo, M. Rahm, P. Vongvilai, T. Brinck and O. Ramström, *Chem. Commun.*, 2008, 6603; (b) A. M. Belenguer, T. Frišić,

- G. M. Day and J. K. M. Sanders, *Chem. Sci.*, 2011, **2**, 696; (c) A. M. Belenguer, G. I. Lampronti, D. J. Wales and J. K. M. Sanders, *J. Am. Chem. Soc.*, 2014, **136**, 16156; (d) B. Rasmussen, A. Sorensen, H. Gotfredsen and M. Pittelkow, *Chem. Commun.*, 2014, **50**, 3716.
- 6 F. Dénès, M. Pichowicz, G. Povie and P. Renaud, *Chem. Rev.*, 2014, **114**, 2587.
- 7 (a) S. Ji, W. Cao, Y. Yu and H. Xu, *Angew. Chem., Int. Ed.*, 2014, **53**, 6781; (b) S. Ji, W. Cao, Y. Yu and H. Xu, *Adv. Mater.*, 2015, **27**, 7740.
- 8 J. Li, C. Nagamani and J. S. Moore, *Acc. Chem. Res.*, 2015, **48**, 2181.
- 9 V. Misik and P. Riesz, *J. Phys. Chem.*, 1994, **98**, 1634.
- 10 A. Studer and D. P. Curran, *Angew. Chem., Int. Ed.*, 2016, **55**, 58.
- 11 (a) A. Wilson, G. Gasparini and S. Matile, *Chem. Soc. Rev.*, 2014, **43**, 1948; (b) A. G. Orrillo, A. M. Escalante and R. L. E. Furlan, *Chem. Commun.*, 2008, 5298; (c) Z. Rodriguez-Docampo and S. Otto, *Chem. Commun.*, 2008, 5301; (d) M. von Delius, E. M. Geertsema, D. A. Leigh and A. M. Z. Slawin, *Org. Biomol. Chem.*, 2010, **8**, 4617; (e) S. Diemer, M. Kristensen, B. Rasmussen, S. Beeren and M. Pittelkow, *Int. J. Mol. Sci.*, 2015, **16**, 21858.

