# Chem Soc Rev



## **REVIEW ARTICLE**

**View Article Online** 

## Chiral metal nanoparticle-catalyzed asymmetric C-C bond formation reactions

Cite this: Chem. Soc. Rev., 2014, 43, 1450

Tomohiro Yasukawa, Hiroyuki Miyamura and Shū Kobayashi\*

Received 10th August 2013 DOI: 10.1039/c3cs60298b

www.rsc.org/csr

Chiral ligand-modified metal nanoparticles possess an attractive potential for application in asymmetric synthesis. This article focuses on chiral-nanoparticle-catalyzed asymmetric C-C bond formation reactions and discusses the nature of the active species.

### 1. Introduction

Rapid progress has been made in the field of homogeneous chiral metal complexes and many types of difficult transformations have been realized, including asymmetric C-C bond formation, to construct complicated molecular structures with high selectivity. 1,2 However, the application of these homogeneous catalysts to large-scale industrial processes is still limited.<sup>3-9</sup> This is because industry generally prefers heterogeneous catalysts because of their great advantages, such as low cost, reusability, avoidance of metal contamination of products, and the availability for reaction integration such as in continuous flow systems and tandem reactions. 10-14 One of the common strategies to convert homogeneous catalysts to heterogeneous catalysts is immobilization

Department of Chemistry, School of Science, The University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113-0033, Japan. E-mail: shu\_kobayashi@chem.s.u-tokyo.ac.jp; Fax: +81-3-5684-0634

of chiral ligands on inorganic or organic supports followed by introduction of metal salts to form "immobilized chiral metal complexes."15-19 While this strategy has already been investigated for several reactions, these systems have drawbacks, such as complicated preparation of monomeric ligands and individual fabrication of heterogeneous polymers. In addition, these catalysts themselves are not robust because of metal leaching and the instability of immobilized ligands, and it is difficult to revive their activity when the catalysts are deactivated.

Metal nanoparticles (NPs) in catalysts are of great interest in both academia and industry, not only due to their very large surface areas, producing high catalytic activity, but also due to their quantum size effect that sometimes shows high activity and unique selectivity.20-23 Recently, several reports have shown that homogeneous metal complex-catalyzed reactions can be conducted by using immobilized metal NP catalytic systems, with the advantages of the heterogeneous catalysts mentioned above.<sup>24</sup> It might indicate that metal NPs possess the potential



Tomohiro Yasukawa

Tomohiro Yasukawa was born in Kanagawa, Japan, in 1987 and studied at the University of Tokyo, receiving his MSc degree in 2012 working under the direction of Prof. Shū Kobayashi. At present, he is a PhD student in the group of Prof. Shū Kobayashi. His research interests include development of chiral metal nanoparticle catalysts, polymer catalysts immobilized environmentally benign organic synthesis.



Hiroyuki Miyamura

Hiroyuki Miyamura was born in 1982 in Saitama, Japan. He received his BS degree in pharmaceutical sciences in 2004 from The University of Tokyo. He received his MSc and PhD degrees at the same University in 2006 and 2009 under the supervision of Professor Shū Kobayashi. In 2009 just after receiving PhD, he was appointed assistant professor in the Department of chemistry at The University of Tokyo. In 2010, he received the winner of The Reaxys PhD Prize.

His research interest includes metal nanocluster catalysis and reaction integration such as tandem reaction and flow synthesis.

**Review Article** Chem Soc Rev




Fig. 1 Functionalized chiral NP catalysts.

for application in asymmetric synthesis, namely, the use of chiral-ligand-modified metal NPs (chiral NPs) as asymmetric catalysts (Fig. 1).

Chiral NPs themselves have attracted the attention of many researchers in various fields as they show characteristic physical properties<sup>25-28</sup> such as circular dichroism (CD) and metal-based electronic transitions. Furthermore, it may be possible to use them not only for asymmetric catalysis, but also as chiral nematic liquid crystals, <sup>29–32</sup> in chiral recognition, <sup>33–36</sup> etc.

Chiral NPs have been mainly prepared by two types of methods:<sup>28</sup> direct synthesis of NPs in the presence of a chiral modifier as a stabilizer (Fig. 2a) or ligand exchange from stabilized NPs (Fig. 2b). In the latter method, it is possible that chiral NPs form in situ simply by combining chiral modifiers and metal NPs in the reaction medium and then using them directly as chiral catalysts.

The first example of heterogeneous asymmetric catalysis using this concept was reported by Orito and co-workers in 1979, demonstrating asymmetric hydrogenation of methyl



Shū Kobayashi

Shū Kobayashi studied at the University of Tokyo, receiving his PhD in 1988 working under the direction of Professor T. Mukaiyama. Following an initial period as assistant professor, he was promoted to lecturer then associate professor at Science University of Tokyo (SUT). In 1998, he moved to the Graduate School of Pharmaceutical Sciences, the University of Tokyo, as full professor. In 2007, he was appointed to his current position

as professor of organic chemistry in the Department of Chemistry, Faculty of Science, The University of Tokyo. He has held various visiting professorships, including the Universite Louis Pasteur, Strasbourg (1993), Kyoto University (1995), Nijmegen University (1996), Philipps-University of Marburg (1997), Paris-Sud (2010). Professor Kobayashi has wide-ranging research interests that include the development of new synthetic methods and novel catalysts, organic reactions in water, solid-phase synthesis, total synthesis of biologically interesting compounds, and organometallic chemistry.




Fig. 2 (a) Direct synthesis of chiral NPs; (b) synthesis of chiral NPs via ligand exchange

Scheme 1 Orito-type reactions.

pyruvate or methyl benzovlformate catalyzed by cinchonidinemodified Pt on carbon (Scheme 1). 37,38 Although the first Oritotype reaction was reported more than 30 years ago, this type of reaction is still being studied. 39-43 Recently, studies were carried out to improve the catalyst system<sup>44-61</sup> and to reveal its detailed mechanism. 62-67 It is generally accepted that the chemisorption layer, through the formation of individual 1:1 diastereomeric complexes from substrates and chiral modifiers, is the key to inducing chirality in this reaction. 62 Very recently, Maeda, Baiker and co-workers proposed a possible intermediate and complex between a substrate and a modifier formed by multiple hydrogen bonds. 64 Based on this chiral NP system, other types of chiral NP catalysts for asymmetric hydrogenation reactions, for example nickel-boride NPs, 68 Pd NPs, 69-85 Ru NPs, 86-90 Rh NPs, 88,91-96 Ir NPs, 88,97-99 and Fe NPs, 100 have been widely investigated. Several reviews illustrate examples of heterogeneous asymmetric hydrogenation reactions. 17,101-104

On the other hand, asymmetric C-C bond formation reactions catalyzed by chiral NPs are a challenging topic and less well developed, judging from the mechanism of the corresponding homogeneous metal complex system. The mechanism of chiral induction is more complicated and multicoordination of both chiral modifier and substrates to NPs is required in transition states. Moreover, this type of multicoordination may easily cause the metal to leach from supports or the decomposition of NPs to form molecular complexes. For these reasons, the nature of the active species should be examined carefully because it is also possible that such a homogeneous molecular complex is a real active species. 105-108 However, lack of fair observation and

Chem Soc Rev **Review Article** 

discussion about heterogeneity is sometimes found. Widely accepted tests to confirm heterogeneity, for example, the mercurypoisoning test, hot filtration test (filtrate transfer test), and three-phase test, are usually conducted. 105

In this review, we focus on chiral NP-catalyzed asymmetric C-C bond formation. Although the detailed mechanism and active species are still unclear, some reports describe the specific nature of chiral NPs catalysts, which showed different catalyst behavior from the corresponding homogeneous molecular complex catalysts.

### 2. Palladium nanoparticles

#### 2.1 Asymmetric allylic alkylation

Palladium-catalyzed asymmetric allylic alkylation reactions are one of the most powerful tools to construct asymmetric C-C bonds and have been widely investigated in homogeneous Pd complex systems. 109,110 Several groups have reported that chiral palladium NPs catalyzed this reaction using various chiral modifiers and, interestingly, the assumed nature of the active species was different for each of the catalyst systems.

In general, stabilization methods for metal NPs are crucial to develop efficient catalysts. In this context, Kobayashi and co-workers developed microencapsulated (MC) catalysts. 111-114 Microcapsules have been used for coating and isolating substances until their activity is required, and their applications in medicine and pharmacy have been extensively studied. 115 This microencapsulation technique was applied to the immobilization of metal NP catalysts into polymers. That is, catalysts would be physically enveloped by polymer backbones and, simultaneously, immobilized by the interaction between the  $\pi$  electrons of the benzene rings of the polystyrene used as a polymer support and vacant orbitals of the catalysts (metal NPs) (Fig. 3).

In 2002, Kobayashi et al. stabilized Pd(0) by the microencapsulation technique, and the thus prepared MC Pd(0) (MC Pd(PPh<sub>3</sub>)) was applied to a catalytic asymmetric allylation reaction (Scheme 2). 116 The reaction of 1,3-diphenyl-2-propen-1-yl




Fig. 3 Microencapsulation technique.

$$\begin{array}{c} OCO_{2}Et \\ Ph \end{array} + \begin{array}{c} CO_{2}R \\ CO_{2}R \end{array} \xrightarrow{\begin{array}{c} MC \ Pd(PPh_{3}) \ (20 \ mol\%) \\ \textbf{2} \ (20 \ mol\%) \\ BSA, \ KOAc \\ MeCN, \ reflux \end{array}} \begin{array}{c} RO_{2}C \\ Ph \end{array} \xrightarrow{\begin{array}{c} CO_{2}R \\ Ph \end{array}} \begin{array}{c} Ph \\ Ph \end{array}$$

Scheme 2 MC Pd-catalyzed asymmetric allylation

ethyl carbonate (1.0 equiv.) with dimethyl malonate (3.0 equiv.) was performed in the presence of MC Pd(PPh<sub>3</sub>) (20 mol%), chiral phosphine-oxazoline 2 (20 mol%), bis(trimethylsilyl)acetamide (BSA) (3.0 equiv.), and potassium acetate (0.10 equiv.) under reflux conditions in acetonitrile. The allylation adduct (R)-3 was obtained in 87% yield with 83% ee.

In 2004, Gómez, Philippot, Chaudret and co-workers reported asymmetric allylic alkylation reactions catalyzed by chiral diphosphite 4 with xylose-backbone-modified colloidal Pd NPs (Schemes 3 and 4). 117 The reactivity of a molecular Pd complex (referred to as **Mol. 4)**, which was generated in situ by the reaction of  $[Pd(C_3H_5)Cl]_2$ with 4, was compared with that of NP catalysts. A significant difference was observed (Table 1). In the presence of a NP catalyst (Pd4), the reaction mainly proceeded with one enantiomer of the racemic 8 and, consequently, a very high degree of kinetic resolution was demonstrated, while no kinetic resolution was observed in the presence of **Mol. 4**. The relative rates of alkylation of the two enantiomers of 8 in two catalyst systems

1/2 [Pd<sub>2</sub>(dba)<sub>3</sub>] + L 
$$H_2$$
 (3 bar)  $H_2$  (3 bar)  $H_2$  (7 PdL, L = 4–7  $H_2$  (3 bar)  $H_2$  (7 PdL, L = 4–7  $H_2$  (8 bar)  $H_2$  (8 bar)  $H_2$  (9 bar)  $H_$ 

Scheme 3 Synthesis of chiral Pd NPs

Scheme 4 Chiral diphosphite ligands.

 $\begin{tabular}{ll} \textbf{Table 1} & Asymmetric allylic alkylation of $\it{rac-8}$ with dimethyl malonate by a Pd catalyst \\ \end{tabular}$ 

**Review Article** 

OAc Ph + CO <sub>2</sub> R						₹ + (S)- <b>8</b> `Ph
Entry	Catalyst	$Pd/L^{\textstyle *}$	Time (h)	Conv. (%)	ee of 3 (%)	ee of 8 (%)
1	Pd4	1/0.2	24	56	97	89
2	Pd4	1/0.2	168	59	97	89
3	Pd4	1/1.05	168	61	97	89
4	Mol. 4	1/1	1.5	95	90	0

were studied. The kinetic preference for the *R*-8 was only a factor of 2 in the **Mol.** 4 system, whereas it was clearly higher (12–20) in the **Pd4** system.

In 2007, the same group reported detailed studies of this reaction and showed higher sensitivity of NP catalyst systems upon adjustment of the metal, the chiral modifier and the substrate than that of molecular catalysts. 118 For example: (i) Pd5 showed a remarkable difference in stability and decomposed into molecular species, while Pd4 did not, and (ii) alkyl-substituted allyl acetates were not alkylated by using Pd4 or Pd6, while the corresponding molecular catalyst systems gave high conversions. In 2008, Diéguez, Gómez, Leeuwen and co-workers reported modified Pd NPs systems with chiral oxazolinyl-phosphite ligands 7a-7e for the same reaction. 119 The nature of the active species was carefully studied by continuous-flow membrane reactor experiments, transmission electron microscopy observations, classical poisoning experiments, and kinetic measurements. The results proved the molecular nature of the true catalysts, which were leached, as measured by inductively coupled plasma atomic emission spectrometry (ICP-AES) analysis.

In 2005, Felpin and Landais reported a heterogeneous Pd catalyst with a chiral modifier that promoted asymmetric allylic alkylation. 120 Pd/C and (R)-2,2'-bis(diphenylphosphino)-1,1'binaphthyl (BINAP) 9 were used as catalysts for the reaction of racemic 8 with diethylmalonate in water, and high enantioselectivity (80% ee) was achieved in spite of low yield (21%). Subsequently, in 2007, Baiker and co-workers reported 9 modified Pd/Al<sub>2</sub>O<sub>3</sub> catalyst systems for the reaction of racemic 8 with the sodium salt of dimethyl malonate (Scheme 5). 121 Reductive treatment of Pd/Al<sub>2</sub>O<sub>3</sub> in hydrogen before the reaction was required for obtaining reproducibility and, as a result, good chemoselectivity (94%) and moderate enantioselectivity (~60% ee) were obtained. The enantioselectivity of this system was independent of the reaction temperature (60 and 120 °C), while the corresponding homogeneous complex system afforded poor enantioselectivity to give the opposite enantiomer of the product (R-3) at 60  $^{\circ}$ C or above. In a report from the same group in 2008, the enantioselectivity of this system was improved (82% ee) by changing the chiral modifier to a ferrocenyl phosphine ligand 10. 122 As in the case of the Pd/Al<sub>2</sub>O<sub>3</sub>-BINAP system, no dependence of enantioselectivity on temperature was observed (60 and 120 °C), while the enantioselectivity was decreased by increasing the reaction temperature from 85% ee at 60  $^{\circ}\text{C}$  to 55% ee at 120  $^{\circ}\text{C}$  in the

OAc 
$$Ph$$
 Ph CO<sub>2</sub>Me  $O$  CO<sub>2</sub>

 $\begin{tabular}{ll} Scheme 5 & Asymmetric allylic alkylation of {\it rac-8} with dimethyl malonate by Pd catalyst. \end{tabular}$ 

corresponding homogeneous complex system. Here, the same enantiomer of the product (S-3) was produced. To confirm the metal leaching in the reaction mixture, the catalytic activity in the filtrate with additional substrates was examined after the first reaction was conducted at 120  $^{\circ}$ C for 6 h and the catalyst was removed by centrifugation. The reaction was continued for a further 6 h and gave the additional product in 19% yield, while the yield in the blank experiment, which was conducted in the absence of Pd/Al<sub>2</sub>O<sub>3</sub>, was 10%.

In 2010, the same group reported a detailed study of the nature of the active species in the Pd/Al<sub>2</sub>O<sub>3</sub>-BINAP system and emphasized heterogeneity. 123 The oxidation state of Pd was investigated by in situ X-ray absorption near-edge structure analysis of the reaction mixture. They did not observe oxidized Pd in the supernatant solution and hence proposed that the molecules in the reaction mixture, such as the solvent and sodium dimethyl malonate, were able to reduce Pd atoms on the surface of Pd NPs and to maintain the reduced state during the reaction. They also pointed out that addition of a chlorinated solvent, such as chloroform, favored dissolution of metallic Pd to form homogeneous catalysis. After the first reaction at 120 °C for 3 h, the reaction mixture was cooled. The catalyst was removed by centrifugation and the supernatant with additional sodium malonate did not give further conversion at room temperature over 18 h in the case of pure THF as solvent, while a small conversion was observed in a similar control experiment in the presence of chloroform.

#### 2.2 Asymmetric cross-coupling reactions

Palladium-catalyzed cross-coupling reactions are one of the biggest breakthroughs in transition metal-catalyzed C–C bond formation reactions. <sup>124</sup> Not only have homogeneous catalysts been developed, but also many heterogeneous catalyst systems. <sup>107,108,125</sup> Among these systems, a couple of examples of asymmetric cross-coupling systems were reported recently.

In 2008, Fujihara and co-workers succeeded in preparing the optically active mono- and bisphosphine-modified chiral Pd NPs and applying these NPs to asymmetric Suzuki-Miyaura coupling reactions (Scheme 6). The zero oxidation state of the chiral Pd NPs was confirmed by X-ray photoelectron

Chem Soc Rev **Review Article** 

$$R + ArB(OH)_{2} \xrightarrow{(S)-9-Pd NPs (0.1 mol\%) \\ KF or Ba(OH)_{2} (3 equiv)}$$

$$DME/H_{2}O, -7 - 25 °C \qquad Ar$$

$$11a: R = Me$$

$$11b: R = OMe$$

$$12a: Ar = 1-naphthyl$$

$$12b: Ar = (2-Me)C_{6}H_{4}$$

$$11c: R = OEt$$

$$4 examples$$

$$42-90\% yield$$


$$55-74\% ee$$

Scheme 6 Asymmetric Suzuki-Miyaura coupling reactions catalyze by chiral Pd NPs

spectroscopy (XPS) analysis, and CD spectra of the chiral Pd NPs showed negative Cotton effects. It was demonstrated that (S)-BINAP-modified chiral Pd NPs could catalyze the coupling reactions with naphthyl halides and arylboronic acids to afford axially chiral biaryl compounds in good enantioselectivity at low temperature (-7 to 25 °C), although reported syntheses of binaphthalenes catalyzed by a chiral Pd-bisphosphine complex required high temperatures to obtain acceptable yields. The latter showed low enantioselectivity.

In 2009, Yamashita and co-workers reported the BINAPmodified core-shell structured bimetallic Fe/Pd NP-catalyzed asymmetric Suzuki-Miyaura coupling reaction. 127 The Fe/Pd NPs with an Fe-rich core and a Pd-rich shell were synthesized by thermal decomposition of Fe(CO)<sub>5</sub> and subsequent reduction of Pd(acac)<sub>2</sub> in the presence of oleic acid and oleylamine, and BINAP-modified NPs were formed through a simple ligand exchange procedure. X-ray absorption measurements revealed that most Pd atoms existed in a metallic form while Fe atoms existed in an oxidized form. The optically active BINAP-modified Fe/Pd NPs showed catalytic activity for the coupling reaction between halide 11b and boronic acid 12a with moderate enantioselectivity (up to 48% ee). The NPs were recovered easily by applying an external magnet and could be reused, while maintaining identical enantioselectivity without an external addition of chiral modifier. A hot filtration test was conducted and the reaction hardly occurred in the filtrate after removal of the catalyst.

In 2010, Glorius and co-workers reported the first successful application of chiral N-heterocyclic carbene (NHC) as a chiral modifier of magnetite supported Pd NPs (Scheme 7) in the asymmetric α-arylation of ketones with aryl halides (Scheme 8). 128 Chiral Pd NPs were prepared by addition of a NHC precursor and a base in the presence of Fe<sub>3</sub>O<sub>4</sub>/Pd. A zero-oxidation state of Pd was confirmed, even after surface modification, by XPS analysis. The presence of a NHC-modified surface was supported by differences between spectra of the free salt 13 and that of Fe<sub>3</sub>O<sub>4</sub>/Pd/13 in attenuated total reflection infrared spectroscopy analysis. Various ketones and aryl halides could be employed for



Scheme 7 Preparation of Fe<sub>3</sub>O<sub>4</sub>/Pd NPs modified by chiral NHC 13

Scheme 8 The asymmetric  $\alpha$ -arylation of ketones with aryl halides

asymmetric α-arylation reactions, including intramolecular reactions, to afford the corresponding α-arylated ketones in moderate-to-good enantioselectivity. The paramagnetic catalyst could be recovered using a magnet and then reused five times without significant loss of activity and selectivity. The heterogeneous nature of active species was indicated by several experiments, such as the hot filtration test, mercury-poisoning test and trace metal analysis. Moreover, dramatically low selectivity was exhibited by several homogeneous complex systems, which afforded the racemic α-arylated product, biphenyl, and other side products. The same research group also developed a homogeneous Pd catalyst system with quinine for this asymmetric α-arylation reaction and achieved excellent substrate generality and enantioselectivity. 129 A combination of Fe<sub>3</sub>O<sub>4</sub>/Pd and quinine as a chiral modifier was also examined. The heterogeneous nature of the real active species was indicated by the mercury-poisoning test and hot filtration test. The results emphasized the important role of chiral NHC for both stability and activity of the Pd NPs as catalysts.

In 2011, Glorius and co-workers further developed immobilized Pd catalysts with NHC. 130 From magnetite-supported NHC 14, a molecular Pd catalyst 15 was prepared with Pd(OAc)2, while a Pd NP catalyst 16 was also prepared by the reduction of K<sub>2</sub>PdCl<sub>4</sub> (Scheme 9). These catalysts were characterized by XPS analysis to confirm the formation of a Pd complex or Pd(0), respectively, and the catalytic activities of these two Pd catalysts were tested in the asymmetric allylation of 4-nitrobenzaldehyde with allyltributyltin (Scheme 10). The NP catalyst 16 provided improved allylation results for both yield and enantioselectivity, and, surprisingly, the product was obtained with the opposite absolute configuration to the one formed with the molecular catalyst 15.

## Rhodium nanoparticles

#### Hydroformylation 3.1

Asymmetric hydroformylation of olefins using chiral Rh complexes in the presence of syngas is a useful method to synthesize optically active aldehydes with excellent regioselectivity and enantioselectivity.<sup>131</sup> Despite extensive development of homogeneous catalysis and recent advances in Rh NP catalysts, 132 heterogeneous asymmetric hydroformylation is still a challenging topic. 133

In 2000, Anderson and co-workers reported styrene hydroformylation over chiral-ligand-modified Rh/SiO2·Al2O3. Chiral induction was achieved by using DIOP 17 or chiraphos 18 (Scheme 11) as a chiral modifier although selectivity was very Review Article Chem Soc Rev

Scheme 9 Preparation of a molecular catalyst and a Pd NP catalyst on  ${\rm Fe_3O_4}.$ 

Scheme 10 Asymmetric allylation of 4-nitrobenzaldehyde with allyltributyltin.

Scheme 11 Chiral diphosphine ligands

low (up to 9% ee).  $^{134}$  In 2006, Li and co-workers reported the asymmetric hydroformylation of styrene or vinyl acetate catalyzed by BINAP-modified Rh/SiO<sub>2</sub> (Scheme 12).  $^{135}$  To achieve chiral induction, an optimized ratio of Rh and modifier was close to 1.0 and, as a result, up to 72% ee and 100% selectivity of a branched aldehyde for the hydroformylation was obtained, although conversion was very low (<10%). It was noted that the DIOP 17-modified Rh/SiO<sub>2</sub> catalyst showed higher turnover frequency (TOF) for the hydroformylation of vinyl acetate (128 h<sup>-1</sup>) than that of unmodified Rh/SiO<sub>2</sub> (90 h<sup>-1</sup>), while other diphosphine ligands decreased the catalytic activity.  $^{31}$ P magic angle spinning nuclear magnetic resonance (MAS NMR) analysis

Scheme 12 Asymmetric hydroformylation of olefins by BINAP-Rh/SiO<sub>2</sub>

$$[Rh(\mu - OMe)(cod)]_2 + L \xrightarrow{H_2 (3 \text{ bar})} [Rh_x(THF)_y L_z]$$
 
$$(0.2 \text{ equiv}) \xrightarrow{THF, \text{ rt}} RhL, L = 4 \text{ or } 19$$
 
$$L = 4 \text{ or } 19$$

Scheme 13 Synthesis of chiral Rh NPs 4 and 19.

Scheme 14 Asymmetric hydroformylation of styrene by Rh NPs

and IR spectra of adsorbed CO indicated the existence of a strong interaction between the P atoms of the chiral modifier and the surface Rh sites. In 2008, the same group reported a different method to prepare modified Rh NPs, by one-pot chemical reduction of aqueous rhodium chloride dispersed in toluene in the presence of amphiphilic tetraoctylammonium bromide and (R)-BINAP. These BINAP-modified Rh NPs could also be immobilized on SiO<sub>2</sub> in a vacuum by adsorption; however, no significant improvement of catalytic performance was observed.

In 2008, Axet, Claver and Philippot reported that carbohydratederived diphosphite-ligand-modified colloidal Rh NPs (Scheme 13) catalyzed the asymmetric hydroformylation of styrene. 137 With the addition of an excess amount of the ligand, high conversion, high regioselectivity, and moderate enantioselectivity were observed in the presence of Rh19 NPs (Scheme 14). The catalytic activities of molecular complexes, which were prepared from [Rh(acac)(CO)<sub>2</sub>] and the corresponding chiral ligands, were examined. The enantioselectivities in the molecular systems were slightly lower ( $\sim$ 10% ee difference) than those in the respective colloidal systems. As results of experiments carried out with dilute solutions and poisoning tests were not conclusive, in situ high-pressure NMR spectroscopic studies under catalytic conditions in the presence of Rh4 NPs were conducted and the well-known hydridorhodium diphosphite complex [RhH(CO)<sub>2</sub>(4)] was detected after a reaction time of 45 h. The results indicated that molecular species generated from NPs are real active species. Given the difference in enantioselectivity between the molecular complex system and the NPs system, the possibility that some activity comes from NPs cannot be excluded.

#### 3.2 Pauson-Khand reaction

The Pauson-Khand reaction is a [2+2+1] cycloaddition of an alkyne, an alkene, and carbon monoxide to construct a Chem Soc Rev Review Article

crotonaldehyde (2.5 equiv) 
$$Co_2Rh_2$$
 on charcoal (Co and Rh: ~7.5 mol%)  $20$  (11 mol%)  $Ph_2P$ ,  $PPh_2$  75–94% yield, 51–87% ee (S,S)-BDPP  $20$ 

Scheme 15 Co/Rh NP-catalyzed asymmetric Pauson–Khand type reactions.

cyclopentanone skeleton. In early studies of this reaction, a dicobalt octacarbonyl complex was used as the catalyst, 138,139 but further developments revealed that several other transition metal complexes 138,140 or cobalt NPs 141 could be used for this reaction. Moreover, the use of heterobimetallic NP catalysts has attracted more attention due to their high catalytic activity. 141,142 In 2005, Park and Chung reported charcoal-immobilized Co/Rh heterobimetallic NP-catalyzed asymmetric Pauson-Khand-type reactions with a chiral modifier. 143 In the presence of the Co/Rh NP catalyst and chiral diphosphine modifier 20, the reactions of several envnes with crotonaldehyde as an alternative CO source proceeded, and the corresponding bicyclic cyclopentanones were obtained in high yields with moderateto-good enantioselectivities (Scheme 15). The catalyst could be recovered by filtration and then reused five times with the addition of a new chiral modifier without significant loss of yield and enantioselectivity. Although elemental analysis of the reaction mixture by ICP-AES after three cycles detected 3.9 ppm of Co and 0.3 ppm of Rh species, results of the mercurypoisoning test supported a heterogeneous nature of the active species as Hg(0) completely eliminated further catalysis after the addition of Hg(0) to the reaction mixture.

#### 3.3 1,4-Addition reaction

Since the development of the rhodium-catalyzed asymmetric 1,4-addition of boron compounds to  $\alpha,\beta$ -unsaturated carbonyl compounds by Hayashi and Miyaura,  $^{144}$  both academic research  $^{145-147}$  and large-scale processes  $^{148-150}$  (the largest example being 20 kg) of this reaction have been widely demonstrated; however, due to the high cost of Rh, reusable and robust chiral heterogeneous Rh catalysts are required.

In 2012, Kobayashi and co-workers developed chiral Rh NPs and Rh/Ag bimetallic NPs, which were immobilized on a mixture of polystyrene-based copolymer with a cross-linking moiety and carbon black (PI/CB Rh and PI/CB Rh/Ag), and used them in the asymmetric 1,4-addition of boronic acids to  $\alpha,\beta$ -unsaturated ketones (Scheme 16). In the presence of PI/CB Rh and (S)-BINAP as a chiral modifier, a significant amount of Rh leaching was observed during the reaction between a cyclic substrate and phenylboronic acid. When changing the chiral modifier to optically active chiral diene 21, Rh leaching in a crude mixture was dramatically suppressed to the detection limit of ICP-AES, and the desired product was obtained in high yield and excellent enantioselectivity. The low catalytic activity of PI/CB

Scheme 16 Rh/Ag NP-catalyzed asymmetric 1,4-addition reactions.

Rh towards acyclic substrates was improved after the formation of bimetallic NPs with Ag; various substrates were applicable and the desired β-arylated ketones were obtained in high yields and high enantioselectivities by using PI/CB Rh/Ag and chiral modifiers 21 or 22, without leaching of the metals. Scanning transmission electron microscopy (STEM) analysis and energy dispersive X-ray spectroscopy (EDS) mapping showed that Ag formed alloy NPs with Rh that could prevent the aggregation of Rh NPs, to enhance the catalytic activity. PI/CB Rh/Ag could be recovered by filtration and then reused. High yields and high enantioselectivities could be maintained for eight reactions. The deactivated catalyst, after repeated use, could be restored fully simply by heating. Fourteen cycles were demonstrated. Furthermore, a hot filtration test confirmed that after the reaction the filtrate did not catalyze the 1,4-addition reaction, indicating that the reaction is not catalyzed by homogeneous leached species. A one-pot aerobic oxidation-asymmetric 1,4-addition reaction of an allyl alcohol and an arylboronic acid was also demonstrated by combining with PI/CB Au as an aerobic oxidation catalyst.

## 4. Other metal nanoparticles

### 4.1 Nanocrystalline magnesium oxide

Examples of asymmetric catalysis by chiral modifiers and nontransition metal NPs are rather limited. In 2005, Choudary et al. reported several asymmetric reactions using aerogel-prepared nanocrystalline MgO (NAP-MgO), which is commercially available and possesses a large surface area (590 m<sup>2</sup> g<sup>-1</sup>) and a threedimensional polyhedral structure. 152 The NAP-MgO was used for asymmetric Henry reactions with aldehydes or α-ketoesters in the presence of (S)-BINOL 23 (Scheme 17) and asymmetric Michael reactions with nitroalkanes in the presence of optically active chiral diamine 24 (Scheme 18). In both reactions, high yields and high enantioselectivities were achieved for several substrates, and NAP-MgO could be reused five times after heating at 250 °C for 1 h without loss of activity and enantioselectivity. The control experiments using silylated MgO or protected hydroxyls of 23 indicated that the hydrogen bond interaction between the hydroxy groups of 23 and the hydroxy groups of partially hydrated MgO are essential for obtaining high enantioselectivity. In 2006, the same group reported that Review Article Chem Soc Rev

NAP-MgO (0.125 g)
$$R^{1} = H \text{ or } CO_{2}Et$$

NAP-MgO (0.125 g)
$$23 \text{ (14 mol%)}$$

$$THF, -78 °C$$

$$R^{1} = H \text{ or } CO_{2}Et$$

NAP-MgO (0.125 g)
$$R^{1} = H \text{ or } CO_{2}Et$$

14 examples
$$70-95\% \text{ yield}$$

$$60-98\% \text{ ee}$$

(S)-BINOL 23

Scheme 17 NAP-MgO-catalyzed asymmetric Henry reactions

Scheme 18 NAP-MgO-catalyzed asymmetric Michael reactions with nitroalkanes.

Scheme 19 NAP-MgO-catalyzed asymmetric aldol reactions

Scheme 20 NAP-MgO-catalyzed asymmetric Michael reactions with malonates.

NAP-MgO systems modified with chiral diamine **24** were applicable for asymmetric aldol reactions with moderate yields and enantioselectivities (Scheme 19).<sup>153</sup>

Kantam and co-workers also investigated the NAP-MgO-mediated asymmetric Michael reactions with malonates. The desired products were obtained in high yields and high enantioselectivities in the presence of optically active chiral diamine 25 (Scheme 20). <sup>154</sup>

### 4.2 Copper nanoparticles

In 2008, Kantam and co-workers found that nanocrystalline CuO (nano-CuO) could also be utilized for the asymmetric aldol reactions of acetone with arylaldehydes in the presence

Scheme 21 Nano-CuO-catalyzed asymmetric aldol reactions.

of chiral diamine 25 as a chiral modifier (Scheme 21). Nano-CuO had a higher large surface area (136 m $^2$  g $^{-1}$ ) and smaller crystal size (7–9 nm) and showed higher activity and enantioselectivity compared with commercial CuO. Nano-CuO could be reused for four cycles without significant loss of activity, and no structure or morphology change was observed after the reaction, as confirmed by X-ray diffraction (XRD) analysis. The filtrate, which was obtained after removal of the solids, did not promote the reaction, and the absence of copper in the filtrate was confirmed by atomic absorption spectrometry studies.

#### 4.3 Gold nanoparticles

Although gold NPs as catalysts, such as in aerobic oxidation, have been extensively studied,  $^{156-160}$  asymmetric catalysis with gold NPs had not been reported until very recently. In 2013, Toste, Somorjai and co-workers investigated enantioselective heterogeneous Au NP catalysts for the asymmetric cyclopropanation reaction by immobilization of a chiral self-assembled monolayer (SAM) on the surface of a mesoporous silica support.  $^{161}$  After amino acids, as chiral modifier, were attached to the surface of mesoporous SiO<sub>2</sub> (MCF-17) through the OH terminal, Au ions were introduced, and the encapsulated Au ions were reduced to NPs by exposure to  $H_2$  (Fig. 4).

The obtained Au NPs catalyst (Au@SAM/MCF) was used for the intermolecular cyclopropanation reaction of styrene with propargyl pivalate in the presence of an oxidizer (Scheme 22). The desired product 26 was obtained in good diastereoselectivity and moderate enantioselectivity, when peptide 28 was immobilized on the surface of mesoporous SiO2. The same catalyst could also be used for the asymmetric intramolecular cyclopropanation reaction although, there, moderate conversion and moderate enantioselectivity were observed (Scheme 23). The corresponding homogeneous system using AuCl<sub>3</sub> and unsupported diproline in the intermolecular reaction produced low yield and no enantioselection. While ICP analysis showed no leaching of Au ions to the solution phase due to the repulsion forces between hydrophilic catalyst and hydrophobic solvent, in situ near-edge X-ray adsorption fine structure measurements carried out under reaction conditions proved that the formation




Fig. 4 Au NPs encapsulated in chiral SAM/mesoporous MCF-17 support.

Chem Soc Rev **Review Article** 

Scheme 22 Au@SAM/MCF-17-catalyzed asymmetric intermolecular cyclopropanation.

Scheme 23 Au@SAM/MCF-17-catalyzed asymmetric intramolecular cyclopropanation

of Au(III) ions, which were generated from Au(0) NPs by oxidation with PhICl<sub>2</sub>, were the active species. Spectroscopic measurements revealed that the enantioselectivity was correlated with the stability of SAM, by the formation of a hydrogen bonding network. Although, in this study, the active species was proved to be not the NPs themselves, the advantages of the surrounding chiral SAM for the formation of a mesoscale enantioselective catalyst were noted.

### 5. Conclusion

The recent reports that we reviewed clearly showed the potential of chiral NP catalysts not only for asymmetric hydrogenation, but also for various types of asymmetric C-C bond formation reactions. Chiral NPs can be easily generated from metal precursors and chiral modifiers. The obtained NPs can be immobilized on several insoluble supports, such as metal oxides, polymers and magnetic NPs, to form heterogeneous catalysts. It was demonstrated that such catalysts could be recovered by simple operations and reused several times, without significant loss of activities and selectivities. The enantioselectivities in chiral NP catalyst systems were comparable to those in the corresponding homogeneous catalytic systems, and some chiral NP catalyst systems showed different selectivities compared with homogeneous catalytic systems. These facts indicate that chiral NP catalysts are not just "immobilized homogeneous catalysts;" they should be considered as distinct active species. The choice of the chiral modifier is a critical key not only to control enantioselectivity, but also to stabilize the NP catalyst itself. A wrong choice of chiral modifier can lead to the decomposition of NP catalyst to a corresponding homogeneous catalyst.

Given these aspects, we should be careful when discussing the real active species. Although the heterogeneity was confirmed by several traditional experiments, such as the poisoning test,

hot filtration test, three-phase test, and ICP analysis in a reaction mixture, it is still unclear how the chiral environment is constructed around the NPs and how chiral modifiers and substrates interact with the surface of NPs or any other active species.

It is important to clarify the chiral active species and the mechanism of specific selectivities in chiral NP-catalyzed reactions. This is an emerging area of research. Further studies should lead to achievement of more efficient and unique chiral NP-catalyzed reactions.

### Acknowledgements

Our work written in this article was partially supported by a Grant-in-Aid for Science Research from the Japan Society for the Promotion of Science (JSPS), Global COE Program, The University of Tokyo, MEXT Japan, NEDO, and Japan Science Technology Agency (JST). T.Y. thanks the JSPS for a Research Fellowship for Young Scientists.

### References

- 1 Asymmetric Synthesis More Methods and Applications, ed. M. Christmann and S. Bräse, Wiley-VCH John Wiley & Sons, Hoboken, 2012.
- 2 Catalytic Asymmetric Synthesis, ed. I. Ojima, John Wiley, Hoboken, N.J., 3rd edn, 2010.
- 3 Asymmetric Catalysis on Industrial Scale: Challenges, Approaches and Solutions, ed. H. U. Blaser and H.-J. Federsel, Wiley-VCH, Weinheim, 2010.
- 4 G. P. Howell, Org. Process Res. Dev., 2012, 16, 1258-1272.
- 5 C. A. Busacca, D. R. Fandrick, J. J. Song and C. H. Senanayake, Adv. Synth. Catal., 2011, 353, 1825-1864.
- 6 N. B. Johnson, I. C. Lennon, P. H. Moran and J. A. Ramsden, Acc. Chem. Res., 2007, 40, 1291-1299.
- 7 H. Shimizu, I. Nagasaki, K. Matsumura, N. Sayo and T. Saito, Acc. Chem. Res., 2007, 40, 1385-1393.
- 8 V. Farina, J. T. Reeves, C. H. Senanayake and J. J. Song, Chem. Rev., 2006, 106, 2734-2793.
- 9 M. Breuer, K. Ditrich, T. Habicher, B. Hauer, M. Kesseler, R. Sturmer and T. Zelinski, Angew. Chem., Int. Ed., 2004, 43,
- 10 P. Barbaro, F. Liguori, N. Linares and C. M. Marrodan, Eur. J. Inorg. Chem., 2012, 3807-3823.
- 11 C. Lucarelli and A. Vaccari, Green Chem., 2011, 13, 1941–1949.
- 12 B. Pugin and H.-U. Blaser, Top. Catal., 2010, 53, 953-962.
- 13 L. Yin and J. Liebscher, Chem. Rev., 2006, 107, 133-173.
- 14 M. Butters, D. Catterick, A. Craig, A. Curzons, D. Dale, A. Gillmore, S. P. Green, I. Marziano, J.-P. Sherlock and W. White, Chem. Rev., 2006, 106, 3002-3027.
- 15 D. Zhao and K. Ding, ACS Catal., 2013, 3, 928-944.
- 16 A. F. Trindade, P. M. P. Gois and C. A. M. Afonso, Chem. Rev., 2009, 109, 418-514.
- 17 M. Heitbaum, F. Glorius and I. Escher, Angew. Chem., Int. Ed., 2006, 45, 4732-4762.

Review Article Chem Soc Rev

- 18 P. McMorn and G. J. Hutchings, Chem. Soc. Rev., 2004, 33, 108–122.
- 19 S. Bräse, F. Lauterwasser and R. E. Ziegert, *Adv. Synth. Catal.*, 2003, **345**, 869–929.
- 20 N. Yan, C. Xiao and Y. Kou, Coord. Chem. Rev., 2010, 254, 1179–1218.
- 21 Nanoparticles and Catalysis, ed. D. Astruc, Wiley-VCH, Weinheim, 2008.
- 22 D. Astruc, F. Lu and J. R. Aranzaes, Angew. Chem., Int. Ed., 2005, 44, 7852–7872.
- 23 D. L. Fedlheim and C. A. Foss, *Metal Nanoparticles: Synthesis, Characterization, and Applications*, CRC Press, 2001.
- 24 H. Cong and J. A. Porco, ACS Catal., 2012, 2, 65-70.
- 25 Y. Xia, Y. Zhou and Z. Tang, Nanoscale, 2011, 3, 1374-1382.
- 26 A. Guerrero-Martínez, J. Lorenzo Alonso-Gómez, B. Auguié, M. Magdalena Cid and L. M. Liz-Marzán, *Nano Today*, 2011, 6, 381–400.
- 27 C. Noguez and I. L. Garzón, Chem. Soc. Rev., 2009, 38, 757–771.
- 28 C. Gautier and T. Bürgi, ChemPhysChem, 2009, 10, 483-492.
- 29 U. Shivakumar, J. Mirzaei, X. Feng, A. Sharma, P. Moreira and T. Hegmann, *Liq. Cryst.*, 2011, **38**, 1495–1514.
- 30 W. Hu, H. Zhao, L. Shan, L. Song, H. Cao, Z. Yang, Z. Cheng, C. Yan, S. Li, H. Yang and L. Guo, *Liq. Cryst.*, 2010, 37, 563–569.
- 31 H. Qi, J. O'Neil and T. Hegmann, J. Mater. Chem., 2008, 18, 374–380.
- 32 H. Qi and T. Hegmann, J. Mater. Chem., 2006, 16, 4197-4205.
- 33 Y. Sun, L. Zhang and H. Li, New J. Chem., 2012, 36, 1442-1444.
- 34 M. Zhang and B.-C. Ye, Anal. Chem., 2011, 83, 1504-1509.
- 35 C.-C. You, S. S. Agasti and V. M. Rotello, *Chem.-Eur. J.*, 2008, **14**, 143–150.
- 36 Y. X. Wang, X. L. Yin, M. H. Shi, W. Li, L. Zhang and J. L. Kong, *Talanta*, 2006, **69**, 1240–1245.
- 37 Y. Orito, S. Imai and S. Niwa, Nippon Kagaku Kaishi, 1979, 1118–1120.
- 38 Y. Orito, S. Imai, S. Niwa and G. H. Nguyen, *J. Synth. Org. Chem., Jpn.*, 1979, 37, 173–174.
- 39 G. Kyriakou, S. K. Beaumont and R. M. Lambert, *Langmuir*, 2011, 27, 9687–9695.
- 40 F. Zaera, J. Phys. Chem. C, 2008, 112, 16196-16203.
- 41 H.-U. Blaser and M. Studer, Acc. Chem. Res., 2007, 40, 1348–1356.
- 42 M. Studer, H. U. Blaser and C. Exner, *Adv. Synth. Catal.*, 2003, 345, 45-65.
- 43 M. von Arx, T. Mallat and A. Baiker, *Top. Catal.*, 2002, **19**, 75–87.
- 44 G. Szőllősi, Z. Makra, M. Fekete, F. Fülöp and M. Bartók, *Catal. Lett.*, 2012, **142**, 889–894.
- 45 E. Schmidt, C. Bucher, G. Santarossa, T. Mallat, R. Gilmour and A. Baiker, *J. Catal.*, 2012, **289**, 238–248.
- 46 S. Sano, M. J. Beier, T. Mallat and A. Baiker, *J. Mol. Catal. A: Chem.*, 2012, 357, 117–124.
- 47 B. Li, X. Li, Y. Ding and P. Wu, Catal. Lett., 2012, 142, 1033–1039.

48 A. Indra and G. K. Lahiri, Chem.-Eur. J., 2012, 18, 6742-6745.

- 49 M. J. Beier, J.-M. Andanson, T. Mallat, F. Krumeich and A. Baiker, ACS Catal., 2012, 2, 337–340.
- 50 B. Török, A. Kulkarni, R. DeSousa, K. Satuluri, M. Török and G. K. S. Prakash, *Catal. Lett.*, 2011, **141**, 1435–1441.
- 51 G. Szőllősi, Z. Makra, F. Fueloep and M. Bartók, Catal. Lett., 2011, 141, 1616–1620.
- 52 B. Li, X. Li, H. Wang and P. Wu, *J. Mol. Catal. A: Chem.*, 2011, 345, 81–89.
- 53 N. Erathodiyil, H. Gu, H. Shao, J. Jiang and J. Y. Ying, *Green Chem.*, 2011, **13**, 3070–3074.
- 54 Z. Chen, Z. Guan, M. Li, Q. Yang and C. Li, *Angew. Chem.*, *Int. Ed.*, 2011, **50**, 4913–4917.
- S. Böttcher, C. Hoffmann, K. Räuchle and W. Reschetilowski, ChemCatChem, 2011, 3, 741–748.
- 56 M. U. Azmat, Y. Guo, Y. Guo, Y. Wang and G. Lu, J. Mol. Catal. A: Chem., 2011, 336, 42–50.
- 57 H. Wang, X. Li, Y. M. Wang and P. Wu, *ChemCatChem*, 2010, 2, 1303–1311.
- 58 S. Nowag, X.-S. Wang, J. Keilitz, A. Thomas and R. Haag, *ChemCatChem*, 2010, 2, 807–811.
- 59 C. Mondelli, C. Bucher, A. Baiker and R. Gilmour, J. Mol. Catal. A: Chem., 2010, 327, 87–91.
- 60 J. Keilitz, S. Nowag, J.-D. Marty and R. Haag, Adv. Synth. Catal., 2010, 352, 1503–1511.
- 61 S. Cserényi, G. Szőllősi, K. Szöri, F. Fülöp and M. Bartók, *Catal. Commun.*, 2010, **12**, 14–19.
- 62 G. Goubert and P. H. McBreen, *ChemCatChem*, 2013, 5, 683–685.
- 63 F. Meemken, N. Maeda, K. Hungerbühler and A. Baiker, *ACS Catal.*, 2012, **2**, 464–467.
- 64 F. Meemken, N. Maeda, K. Hungerbühler and A. Baiker, *Angew. Chem., Int. Ed.*, 2012, **51**, 8212–8216.
- N. Maeda, S. Sano, T. Mallat, K. Hungerbühler and A. Baiker, J. Phys. Chem. C, 2012, 116, 4182–4188.
- 66 N. Maeda, K. Hungerbühler and A. Baiker, *J. Am. Chem. Soc.*, 2011, **133**, 19567–19569.
- 67 E. Schmidt, A. Vargas, T. Mallat and A. Baiker, *J. Am. Chem. Soc.*, 2009, **131**, 12358–12367.
- 68 K. Molvinger, M. Lopez and J. Court, *Tetrahedron Lett.*, 1999, **40**, 8375–8378.
- 69 Z. Makra, G. Szőllősi and M. Bartók, *Catal. Today*, 2012, 181, 56-61.
- 70 N. Győrffy and A. Tungler, J. Mol. Catal. A: Chem., 2011, 336, 72-77.
- 71 T. Sugimura and H. Ogawa, Chem. Lett., 2010, 232-233.
- 72 T. Y. Kim and T. Sugimura, J. Mol. Catal. A: Chem., 2010, 327, 58-62.
- 73 N. Győrffy, A. Tungler and M. Fodor, *J. Catal.*, 2010, **270**,
- 74 S. K. Beaumont, G. Kyriakou, D. J. Watson, O. P. H. Vaughan, A. C. Papageorgiou and R. M. Lambert, *J. Phys. Chem. C*, 2010, 114, 15075–15077.
- 75 D. J. Watson, R. J. B. R. J. Jesudason, S. K. Beaumont, G. Kyriakou, J. W. Burton and R. M. Lambert, *J. Am. Chem. Soc.*, 2009, **131**, 14584–14589.

76 G. Szőllősi, Z. Nemeth, K. Hernadi and M. Bartók, *Catal. Lett.*, 2009, **132**, 370–376.

Chem Soc Rev

- 77 S. Li, C. Chen, E. Zhan, S.-B. Liu and W. Shen, *J. Mol. Catal. A: Chem.*, 2009, **304**, 88–94.
- 78 S. C. Mhadgut, M. Török, S. Dasgupta and B. Török, *Catal. Lett.*, 2008, **123**, 156–163.
- 79 S. Li, E. Zhan, Y. Li, Y. Xu and W. Shen, *Catal. Today*, 2008, **131**, 347–352.
- 80 E. Zhan, S. Li, Y. Xu and W. Shen, *Catal. Commun.*, 2007, **8**, 1239–1243.
- 81 A. I. McIntosh, D. J. Watson and R. M. Lambert, *Langmuir*, 2007, 23, 6113–6118.
- 82 A. I. McIntosh, D. J. Watson, J. W. Burton and R. M. Lambert, *J. Am. Chem. Soc.*, 2006, **128**, 7329–7334.
- 83 R. Strobel, F. Krumeich, W. J. Stark, S. E. Pratsinis and A. Baiker, *J. Catal.*, 2004, 222, 307–314.
- 84 A. Tungler, Y. Nitta, K. Fodor, G. Farkas and T. Mathe, *J. Mol. Catal. A: Chem.*, 1999, **149**, 135–140.
- 85 A. Tungler, T. Mathe, T. Tarnai, K. Fodor, G. Toth, J. Kajtar, I. Kolossvary, B. Herenyi and R. A. Sheldon, *Tetrahedron: Asymmetry*, 1995, 6, 2395–2402.
- 86 L. Ye, H. Lin, H. Zhou and Y. Yuan, *J. Phys. Chem. C*, 2010, **114**, 19752–19760.
- 87 H.-y. Jiang, H. Chen and R.-x. Li, *Catal. Commun.*, 2010, **11**, 584–587.
- 88 A. Gual, C. Godard, K. Philippot, B. Chaudret, A. Denicourt-Nowicki, A. Roucoux, S. Castillón and C. Claver, *ChemSusChem*, 2009, 2, 769–779.
- 89 J. Wang, H. Feng, R. Qin, H. Fu, M. Yuan, H. Chen and X. Li, *Tetrahedron: Asymmetry*, 2007, **18**, 1643–1647.
- 90 S. Jansat, D. Picurelli, K. Pelzer, K. Philippot, M. Gómez, G. Muller, P. Lecante and B. Chaudret, *New J. Chem.*, 2006, 30, 115–122.
- 91 E. G. Bilé, E. Cortelazzo-Polisini, A. Denicourt-Nowicki, R. Sassine, F. Launay and A. Roucoux, *ChemSusChem*, 2012, 5, 91–101.
- 92 Y. Huang, S. Xu and V. S. Y. Lin, *ChemCatChem*, 2011, 3, 690-694.
- 93 E. G. Bilé, A. Denicourt-Nowicki, R. Sassine, P. Beaunier, F. Launay and A. Roucoux, *ChemSusChem*, 2010, 3, 1276–1279.
- 94 F. Hoxha, N. van Vegten, A. Urakawa, F. Krurneich, T. Mallat and A. Baiker, *J. Catal.*, 2009, **261**, 224–231.
- 95 F. Hoxha, T. Mallat and A. Baiker, J. Catal., 2007, 248, 11-19.
- 96 O. J. Sonderegger, G. M. W. Ho, T. Burgi and A. Baiker, J. Catal., 2005, 230, 499–506.
- 97 H.-y. Jiang, B. Sun, X.-x. Zheng and H. Chen, *Appl. Catal.*, *A*, 2012, **421**, 86–90.
- 98 C. Yang, H. Jiang, J. Feng, H. Fu, R. Li, H. Chen and X. Li, *J. Mol. Catal. A: Chem.*, 2009, **300**, 98–102.
- 99 H.-y. Jiang, C.-f. Yang, C. Li, H.-y. Fu, H. Chen, R.-x. Li and X.-j. Li, *Angew. Chem., Int. Ed.*, 2008, 47, 9240–9244.
- 100 J. F. Sonnenberg, N. Coombs, P. A. Dube and R. H. Morris, *J. Am. Chem. Soc.*, 2012, **134**, 5893–5899.
- 101 K. V. S. Ranganath and F. Glorius, *Catal. Sci. Technol.*, 2011, **1**, 13–22.

- 102 P. Barbaro, V. Dal Santo and F. Liguori, *Dalton Trans.*, 2010, 39, 8391–8402.
- 103 S. Roy and M. A. Pericas, Org. Biomol. Chem., 2009, 7, 2669–2677.
- 104 T. Mallat, E. Orglmeister and A. Baiker, *Chem. Rev.*, 2007, 107, 4863–4890.
- 105 R. H. Crabtree, Chem. Rev., 2011, 112, 1536-1554.
- 106 L. D. Pachón and G. Rothenberg, Appl. Organomet. Chem., 2008, 22, 288–299.
- 107 M. Pagliaro, V. Pandarus, R. Ciriminna, F. Beland and P. D. Cara, *ChemCatChem*, 2012, 4, 432–445.
- 108 J. Durand, E. Teuma and M. Gómez, *Eur. J. Inorg. Chem.*, 2008, 3577–3586.
- 109 B. M. Trost, Org. Process Res. Dev., 2012, 16, 185-194.
- 110 B. M. Trost and D. L. VanVranken, *Chem. Rev.*, 1996, **96**, 395–422.
- 111 S. Kobayashi and S. Nagayama, *J. Am. Chem. Soc.*, 1998, 120, 2985–2986.
- 112 R. Akiyama and S. Kobayashi, *Chem. Rev.*, 2009, **109**, 594–642.
- 113 S. Kobayashi and H. Miyamura, *Chem. Rec.*, 2010, **10**, 271–290.
- 114 S. Kobayashi and H. Miyamura, *Aldrichimica Acta*, 2013, **46**, 3–19.
- 115 M. Donbrow, *Microcapsules and Nanoparticles in Medicine and Pharmacy*, CRC Press, Boca Raton, FL, 1991.
- 116 R. Akiyama and S. Kobayashi, *Angew. Chem., Int. Ed.*, 2001, 40, 3469–3471.
- 117 S. Jansat, M. Gómez, K. Philippot, G. Muller, E. Guiu, C. Claver, S. Castillón and B. Chaudret, *J. Am. Chem. Soc.*, 2004, **126**, 1592–1593.
- 118 I. Favier, M. Gómez, G. Muller, M. R. Axet, S. Castillon, C. Claver, S. Jansat, B. Chaudret and K. Philippot, *Adv. Synth. Catal.*, 2007, **349**, 2459–2469.
- 119 M. Diéguez, O. Pamies, Y. Mata, E. Teuma, M. Gómez, F. Ribaudo and P. W. N. M. van Leeuwen, *Adv. Synth. Catal.*, 2008, **350**, 2583–2598.
- 120 F. X. Felpin and Y. Landais, *J. Org. Chem.*, 2005, **70**, 6441–6446.
- 121 S. Reimann, T. Mallat and A. Baiker, *J. Catal.*, 2007, 252, 30–38.
- 122 S. Reimann, T. Mallat and A. Baiker, *J. Catal.*, 2008, 254, 79–83
- 123 S. Reimann, J.-D. Grunwaldt, T. Mallat and A. Baiker, *Chem.-Eur. J.*, 2010, **16**, 9658–9668.
- 124 Palladium-Catalyzed Coupling Reactions: Practical Aspects and Future Developments, ed. Á. Molnár, Wiley-VCH, Weinheim, 2013.
- 125 M. Lamblin, L. Nassar-Hardy, J.-C. Hierso, E. Fouquet and F.-X. Felpin, *Adv. Synth. Catal.*, 2010, 352, 33–79.
- 126 K. Sawai, R. Tatumi, T. Nakahodo and H. Fujihara, *Angew. Chem., Int. Ed.*, 2008, **47**, 6917–6919.
- 127 K. Mori, Y. Kondo and H. Yamashita, *Phys. Chem. Chem. Phys.*, 2009, **11**, 8949–8954.
- 128 K. V. S. Ranganath, J. Kloesges, A. H. Schäfer and F. Glorius, *Angew. Chem., Int. Ed.*, 2010, **49**, 7786–7789.

129 C. Richter, K. V. S. Ranganath and F. Glorius, Adv. Synth.

Catal., 2012, **354**, 377–382.

**Review Article** 

- 130 K. V. S. Ranganath, A. H. Schaefer and F. Glorius, *Chem-CatChem*, 2011, 3, 1889–1891.
- 131 J. Klosin and C. R. Landis, Acc. Chem. Res., 2007, 40, 1251–1259.
- 132 Y. Yuan, N. Yan and P. J. Dyson, *ACS Catal.*, 2012, 2, 1057–1069.
- 133 A. C. B. Neves, M. J. F. Calvete, T. M. V. D. Pinho e Melo and M. M. Pereira, *Eur. J. Org. Chem.*, 2012, 6309–6320.
- 134 J. M. Coronado, F. Coloma and J. A. Anderson, J. Mol. Catal. A: Chem., 2000, 154, 143–154.
- 135 D. Han, X. Li, H. Zhang, Z. Liu, J. Li and C. Li, J. Catal., 2006, 243, 318–328.
- 136 D. Han, X. Li, H. Zhang, Z. Liu, G. Hu and C. Li, *J. Mol. Catal. A: Chem.*, 2008, **283**, 15–22.
- 137 M. R. Axet, S. Castillon, C. Claver, K. Philippot, P. Lecante and B. Chaudret, *Eur. J. Inorg. Chem.*, 2008, 3460–3466.
- 138 J. Blanco-Urgoiti, L. Anorbe, L. Perez-Serrano, G. Dominguez and J. Perez-Castells, *Chem. Soc. Rev.*, 2004, 33, 32-42.
- 139 S. T. Ingate and J. Marco-Contelles, *Org. Prep. Proced. Int.*, 1998, **30**, 121–143.
- 140 H.-W. Lee and F.-Y. Kwong, Eur. J. Org. Chem., 2010, 789–811.
- 141 J. H. Park, K.-M. Chang and Y. K. Chung, *Coord. Chem. Rev.*, 2009, 253, 2461–2480.
- 142 J. H. Park and Y. K. Chung, Dalton Trans., 2008, 2369-2378.
- 143 K. H. Park and Y. K. Chung, Adv. Synth. Catal., 2005, 347, 854–866.
- 144 Y. Takaya, M. Ogasawara, T. Hayashi, M. Sakai and N. Miyaura, *J. Am. Chem. Soc.*, 1998, **120**, 5579–5580.

- 145 P. Tian, H.-Q. Dong and G.-Q. Lin, ACS Catal., 2012, 2, 95–119.
- 146 H. J. Edwards, J. D. Hargrave, S. D. Penrose and C. G. Frost, *Chem. Soc. Rev.*, 2010, **39**, 2093–2105.
- 147 T. Hayashi and K. Yamasaki, Chem. Rev., 2003, 103, 2829-2844.
- 148 G. P. Howell, Org. Process Res. Dev., 2012, 16, 1258-1272.
- 149 J. Magano and J. R. Dunetz, *Chem. Rev.*, 2011, **111**, 2177–2250.
- 150 S. Brock, D. Hose, J. Moseley, A. Parker, I. Patel and A. Williams, *Org. Process Res. Dev.*, 2008, **12**, 496–502.
- 151 T. Yasukawa, H. Miyamura and S. Kobayashi, J. Am. Chem. Soc., 2012, 134, 16963–16966.
- 152 B. M. Choudary, K. V. S. Ranganath, U. Pal, M. L. Kantam and B. Sreedhar, *J. Am. Chem. Soc.*, 2005, **127**, 13167–13171.
- 153 B. M. Choudary, L. Chakrapani, T. Ramani, K. V. Kumar and M. L. Kantam, *Tetrahedron*, 2006, **62**, 9571–9576.
- 154 M. L. Kantam, K. V. S. Ranganath, K. Mahendar, L. Chakrapani and B. M. Choudary, *Tetrahedron Lett.*, 2007, **48**, 7646–7649.
- 155 M. L. Kantam, T. Ramani, L. Chakrapani and K. V. Kumar, Tetrahedron Lett., 2008, 49, 1498–1501.
- 156 Y. Zhang, X. Cui, F. Shi and Y. Deng, *Chem. Rev.*, 2012, **112**, 2467–2505.
- 157 M. Stratakis and H. Garcia, Chem. Rev., 2012, 112, 4469-4506.
- 158 T. Tsukuda, H. Tsunoyama and H. Sakurai, *Chem.-Asian J.*, 2011, **6**, 736-748.
- 159 A. Corma and H. Garcia, Chem. Soc. Rev., 2008, 37, 2096–2126.
- 160 A. Haruta, Chem. Rec., 2003, 3, 75-87.
- 161 E. Gross, J. H. Liu, S. Alayoglu, M. A. Marcus, S. C. Fakra, F. D. Toste and G. A. Somorjai, *J. Am. Chem. Soc.*, 2013, 135, 3881–3886.