



Cite this: *Dalton Trans.*, 2014, **43**, 14514

Received 25th July 2014,  
Accepted 13th August 2014

DOI: 10.1039/c4dt02265c

www.rsc.org/dalton

## The preparation and structure of $\text{Ge}_3\text{F}_8$ – a new mixed-valence fluoride of germanium, a convenient source of $\text{GeF}_2^\dagger$

Andrew L. Hector, Andrew Jolleys, William Levason,\* David Pugh and Gillian Reid

The new binary mixed-valence fluoride of germanium,  $\text{Ge}_3\text{F}_8$ , has been obtained by heating  $\text{GeF}_4$  with powdered Ge in an autoclave (390 K/4 bar/48 h). The structure contains pyramidal  $\text{Ge}^{\text{II}}\text{F}_3$  and octahedral  $\text{Ge}^{\text{IV}}\text{F}_6$  units, linked by fluoride bridges. The new compound is the missing member of the series  $(\text{GeF}_2)_n\text{-GeF}_4$  ( $n = 2, 4$ , or 6). Sublimation of  $(\text{GeF}_2)_n\text{-GeF}_4$  *in vacuo* provides a convenient source of  $\text{GeF}_2$  in ca. 30% overall yield.

Although germanium is technologically very important both as the element and in oxide or chalcogenide compounds, with key applications in electronics, ceramics and optics,<sup>1</sup> its chemistry was neglected for many years compared with those of silicon and tin. It is now a very active area of main group chemistry and, in addition to the extensive chemistry of  $\text{Ge}(\text{IV})$ ,<sup>2</sup> recent work has identified a large and complex coordination chemistry of  $\text{Ge}(\text{II})$ ,<sup>2,3</sup> the latter contrasting with the limited coordination chemistry of  $\text{Si}(\text{II})$ .<sup>2</sup> In Group 14 as well as the common tetrahalides  $\text{MX}_4$  ( $\text{M} = \text{Si}, \text{Ge}, \text{Sn}; \text{X} = \text{F-I}$ ),<sup>4</sup> there are dihalides  $\text{MX}_2$  ( $\text{M} = \text{Ge}, \text{Sn}$ ), and the subhalides,  $\text{GeBr}$  and  $\text{SnBr}$ .<sup>5</sup> Of these, the chemistry of  $\text{GeF}_2$  has been very little explored since it is not readily available commercially and its preparation by repeatedly passing  $\text{GeF}_4$  over heated germanium, is both inconvenient and time consuming,<sup>6</sup> while the alternative method, involving the reaction of Ge with anhydrous HF in an autoclave, is hazardous. Both routes also require special equipment.<sup>7</sup> A number of intermediate halides have also been identified.<sup>2</sup> The latter are of two types; the most common are those with element–element (E–E) bonds, including  $\text{Si}_2\text{F}_6$ ,  $\text{Si}_2\text{Cl}_6$ ,  $\text{Si}_3\text{Cl}_8$ ,  $\text{Si}_6\text{Cl}_{14}$ ,  $\text{Ge}_2\text{Cl}_6$  and  $\text{Ge}_5\text{Cl}_{12}$ ,<sup>8</sup> with structures analogous to the corresponding alkanes. Much

rarer, and limited to Ge and Sn, are a second group of mixed-valence materials, including  $\text{Sn}_3\text{F}_8$ ,  $\text{Ge}_5\text{F}_{12}$ , and  $\text{Ge}_7\text{F}_{16}$ , which are without direct E–E bonds, but are fluoride-bridged and contain distinct environments attributable to  $\text{M}^{\text{II}}$  and  $\text{M}^{\text{IV}}$  centres.<sup>9–11</sup>

We are currently developing new routes for electrodeposition of p-block materials from non-aqueous media, using reagents including halometallate anions as the p-block element source,<sup>12</sup> and have recently reported the electrochemistry of  $[\text{GeX}_3]^-$  ( $\text{X} = \text{Cl}, \text{Br}$  or  $\text{I}$ ) and  $[\text{GeCl}_6]^{2-}$  in  $\text{CH}_2\text{Cl}_2$  solution.<sup>13</sup> During the course of this work we have extended our studies to the fluoride systems. We report here the preparation and characterisation of a new binary, mixed-valence fluoride of germanium and its use to provide a convenient route to  $\text{GeF}_2$ .

Depending upon the experimental conditions, repeatedly passing  $\text{GeF}_4$  at low pressure over heated germanium yields either  $\text{GeF}_2$ ,<sup>6</sup> or mixed valence  $\text{Ge}^{\text{II}}\text{-Ge}^{\text{IV}}$  fluorides.<sup>10,11,14</sup> Two of the latter identified by single crystal X-ray diffraction (XRD) studies are  $\text{Ge}_5\text{F}_{12}^\dagger$  and  $\text{Ge}_7\text{F}_{16}$ ,<sup>10,11</sup> which are members of the series  $(\text{GeF}_2)_n\text{-GeF}_4$ .<sup>14</sup> These flow reactions are inconvenient and low yielding, hence we have investigated the reduction of  $\text{GeF}_4$  with Ge powder in an autoclave under modest pressure (390 K/4 bar/48 h, see ESI†). Initial attempts at temperatures <370 K resulted in little reaction, but on increasing the temperature to 390 K/48 h, much of the  $\text{GeF}_4$  was consumed (as indicated by the drop in pressure), and upon opening the autoclave in a glove-box, a mass of white microcrystalline material was found on the cooler lid. The crystals are extremely moisture sensitive, converting into a pool of liquid immediately on exposure to air. Single crystal X-ray diffraction data were collected from one of the small crystals and the structure solution identified this product as  $\text{Ge}_3\text{F}_8$ , the missing third member of the series  $(\text{GeF}_2)_n\text{-GeF}_4$ , with  $n = 2$ . Unit cell measurements on several other crystals confirmed these as the same compound. Powder X-ray diffraction (PXRD) data were also collected on the bulk material and that showed smaller amounts of  $\text{Ge}_5\text{F}_{12}$  and  $\text{Ge}_7\text{F}_{16}$ , as well as traces of  $\text{GeF}_2$  were also present. The simulated and experimental powder XRD data from this mixture are shown in the ESI.†

Chemistry, University of Southampton, Southampton SO17 1BJ, UK.


E-mail: wxl@soton.ac.uk; Tel: +44 (0)2380 593792

† Electronic supplementary information (ESI) available: Experimental details for the syntheses of  $\text{Ge}_3\text{F}_8$  and  $\text{GeF}_2$ , and the PXRD data for all the products. Further details of the crystal structure investigation may be obtained from Fachinformationszentrum Karlsruhe, 76344 Eggenstein-Leopoldshafen, Germany (fax: (+49)7247-808-666; e-mail: crysdata@fiz-karlsruhe.de, [http://www.fiz-karlsruhe.de/request\\_for\\_deposited\\_data.html](http://www.fiz-karlsruhe.de/request_for_deposited_data.html)) on quoting CSD number 427896. See DOI: 10.1039/c4dt02265c




Sublimation of the mixture (390 K/0.5 mm) gave ~30% yield of  $\text{GeF}_2$  (based on elemental Ge used in the first step), which was identified by PXRD (see ESI†). Some involatile orange material (*cf.* ref. 6) was also formed.

Germanium difluoride has a polymeric chain structure based upon trigonal pyramidal  $\text{GeF}_3$  units ( $\text{Ge-F} = 1.79(2)$ ,



**Fig. 1** The  $\text{Ge}^{\text{II}}$  and  $\text{Ge}^{\text{IV}}$  units in  $\text{Ge}_3\text{F}_8$  with ellipsoids drawn at the 50% level. Selected bond lengths (Å) and angles ( $^\circ$ ):  $\text{Ge2-F1} = 1.855(1)$ ,  $\text{Ge2-F2} = 1.767(1)$ ,  $\text{Ge2-F3} = 1.782(1)$ ,  $\text{Ge1-F1} = 2.010(1)$ ,  $\text{Ge1-F4}^{\text{ii}} = 1.938(1)$ ,  $\text{Ge1-F4} = 1.980(1)$ ,  $\text{F1-Ge1-F4} = 83.28(5)$ ,  $\text{F4-Ge-F4}^{\text{ii}} = 86.35(3)$ ,  $\text{F4}^{\text{ii}}-\text{Ge1-F1} = 85.42(6)$ . Symmetry codes: (i)  $1-x, 1-y, 1-z$ ; (ii)  $0.5-x, -0.5+y, 1.5-z$ ; (iii)  $x, -1+y, z$ .




**Fig. 2** The  $\text{Ge}_3\text{F}_8$  structure viewed along: (a) the  $b$  axis to observe the sheets, and (b) the  $a$  axis, showing the connectivity within the sheets.

$1.91(2)$ ,  $2.09(2)$  Å), with a distant fourth fluoride at  $2.57(2)$  Å that cross-links the chains.<sup>15</sup> The new preparation is a convenient way to obtain  $\text{GeF}_2$  in useful quantity for further studies of its coordination and organometallic chemistry.

The single crystals of the mixed-valence  $\text{Ge}_3\text{F}_8$  are isomorphous with  $\text{Sn}_3\text{F}_8$ ,<sup>9</sup> adopting the monoclinic space group  $P2_1/n$ . The structure is composed (Fig. 1) of slightly distorted  $\text{GeF}_6$  octahedra with four terminal  $\text{Ge-F}$  bonds ( $1.767(1)$ ,  $1.782(1)$  Å), and two slightly longer  $\text{Ge-F}$  bonds ( $1.855(1)$  Å) that are involved in bridging to the  $\text{Ge}^{\text{II}}$  units. The germanium (ii) core environment is trigonal pyramidal, composed of one terminal ( $\text{Ge-F} = 1.938(1)$  Å) and two bridging ( $\text{Ge-F} = 1.980(1)$ ,  $2.010(1)$  Å) fluorides, one linked to  $\text{Ge}^{\text{IV}}$  and one to a second  $\text{Ge}^{\text{II}}$  centre. There are also longer  $\text{Ge}^{\text{II}}\cdots\text{F}$  contacts ( $2.56$  Å), and if these are included, the germanium(ii) geometry is a distorted saw-horse shape, reminiscent of  $\text{GeF}_2$ . Overall, the packing is best considered as sheets in the (101) planes (Fig. 2a), with each sheet being made up of puckered chains of  $\text{GeF}_3$  units along [010] connected together by the  $\text{GeF}_6$  octahedra (Fig. 2b).

Considering the structures of  $\text{Ge}_5\text{F}_{12}$ <sup>10</sup> and  $\text{Ge}_7\text{F}_{16}$ ,<sup>11</sup> the same basic building blocks are present (trigonal pyramidal  $\text{GeF}_3$  and octahedral  $\text{GeF}_6$ ), but as the F/Ge ratio declines, the



**Fig. 3** (a) View of the structure of  $\text{Ge}_5\text{F}_{12}$ .<sup>10</sup> (b) View of the structure of  $\text{Ge}_7\text{F}_{16}$ .<sup>11</sup>



structures become more distorted to maintain the germanium coordination numbers. In  $\text{Ge}_5\text{F}_{12}$ , if we ignore the distant fourth fluoride at 2.44 Å, the  $\text{GeF}_3$  trigonal pyramids ( $\text{Ge}-\text{F} = 1.80(2), 1.99(2), 2.20(2)$  Å) form corrugated sheets in (001), based on  $\text{GeF}_6$  octahedra linked to dimers of two corner-linked  $\text{GeF}_3$  pyramids. As a consequence of the 4 : 1  $\text{Ge}^{\text{II}} : \text{Ge}^{\text{IV}}$  constitution, the  $\text{GeF}_6$  units are linked to four dimers (rather than two as in  $\text{Ge}_3\text{F}_8$ ) (Fig. 3a). The structure of  $\text{Ge}_7\text{F}_{16}$  is complicated in that there are seven distinct germanium sites,<sup>11</sup> but again, the building blocks are trigonal pyramidal  $\text{GeF}_3$  and octahedral  $\text{GeF}_6$  units. The structure is best described as chains of  $\text{GeF}_3$  pyramids along [001] with side chains of four  $\text{GeF}_3$  units terminated by a  $\text{GeF}_6$  octahedron attached to every second  $\text{GeF}_3$  of the main chain (Fig. 3b).

In conclusion, the missing member of the unique series of mixed-valence germanium fluorides  $(\text{GeF}_2)_n\text{-GeF}_4$  ( $n = 2, 4$ , or 6) has been obtained by reaction of  $\text{GeF}_4$  and Ge powder under modest pressure and temperature, its structure determined and the structural relationships within the series established. Sublimation of the  $(\text{GeF}_2)_n\text{-GeF}_4$  *in vacuo* provides a convenient route to the previously rather inaccessible  $\text{GeF}_2$ . Further work to explore the chemistry of  $\text{GeF}_2$  formed by this route is underway and will be reported in due course.

## Acknowledgements

We thank EPSRC for support (EP/1033394/1 and EP/1010890/1). The SCFED Project (<http://www.scfed.net>) is a multidisciplinary collaboration of British universities investigating the fundamental and applied aspects of supercritical fluids.

## Notes and references

‡  $\text{Ge}_5\text{F}_{12}$  was originally formulated as  $\text{Ge}_2\text{F}_5$ ,<sup>14</sup> the correct formula subsequently being established from the crystal structure determination.<sup>10</sup>

- (a) D. D. Vaughn II and R. E. Schaak, *Chem. Soc. Rev.*, 2013, **42**, 2861; (b) S. Raoux, W. Welnic and D. Ielmini, *Chem. Rev.*, 2010, **110**, 240; (c) D. V. Talapin, J. S. Lee, M. V. Kovalenko and E. V. Shevchenko, *Chem. Rev.*, 2010, **110**, 389.
- (a) J. Parr, in *Comprehensive Coordination Chemistry II*, ed. J. A. McCleverty and T. J. Meyer, Elsevier, Oxford, 2004, vol. 3, p. 545; (b) W. Levason, G. Reid and W. Zhang, *Coord. Chem. Rev.*, 2011, **255**, 1319; (c) N. N. Greenwood and A. Earnshaw, *Chemistry of the Elements*, Butterworth, Oxford, 2nd edn, 1997; (d) R. S. Ghadwal, H. W. Roesky, S. Merkel, J. Henn and D. Stalke, *Angew. Chem., Int. Ed.*, 2009, **48**, 5683; (e) A. C. Filippou, O. Chernov and G. Schnakenburg, *Angew. Chem., Int. Ed.*, 2009, **48**, 5687.
- For lead references on  $\text{Ge}(\text{II})$  complexes see: (a) A. J. Arduengo III, H. V. Rasika Dias, J. C. Calabrese and F. Davidson, *Inorg. Chem.*, 1993, **32**, 1541; (b) P. A. Rugar, V. N. Staroverov, P. J. Ragona and K. M. Baines, *J. Am. Chem. Soc.*, 2007, **129**, 15138; (c) F. Cheng, A. L. Hector, W. Levason, G. Reid, M. Webster and W. Zhang, *Angew. Chem., Int. Ed.*, 2009, **48**, 5152; (d) F. Cheng, J. M. Dyke, F. Ferrante, A. L. Hector, W. Levason, G. Reid, M. Webster and W. Zhang, *Dalton Trans.*, 2010, **39**, 847; (e) F. Cheng, A. L. Hector, W. Levason, G. Reid, M. Webster and W. Zhang, *Inorg. Chem.*, 2010, **49**, 752; (f) J. England and K. Wieghardt, *Inorg. Chem.*, 2013, **52**, 10067; (g) P. A. Rugar, M. C. Jennings and K. M. Baines, *Organometallics*, 2008, **27**, 5043; (h) S. M. I. Al-Rafia, M. R. Momeni, R. McDonald, M. J. Ferguson, A. Brown and E. Rivard, *Angew. Chem., Int. Ed.*, 2013, **52**, 6390.
- A. K. Wolf, J. Glinnemann and M. U. Schmidt, *CrystEngComm*, 2008, **10**, 1364.
- (a) A. Schnepf and R. Köppe, *Z. Anorg. Allg. Chem.*, 2002, **628**, 2914; (b) C. Schrenk, R. Köppe, I. Schellenberg, R. Pöttgen and A. Schnepf, *Z. Anorg. Allg. Chem.*, 2009, **635**, 1541.
- N. Bartlett and K. C. Yu, *Can. J. Chem.*, 1961, **39**, 80.
- E. L. Muetterties, *Inorg. Chem.*, 1962, **1**, 342.
- (a) E. Hengge, in *Halogen Chemistry*, ed. V. Gutmann, Academic Press, NY, 1967, vol. 2, p. 169; (b) D. Shriver and W. L. Jolly, *J. Am. Chem. Soc.*, 1958, **80**, 6692; (c) I. R. Beattie, P. J. Jones, G. Reid and M. Webster, *Inorg. Chem.*, 1998, **37**, 6032.
- M. F. A. Dove, R. King and T. J. King, *J. Chem. Soc., Chem. Commun.*, 1973, 944.
- J. C. Taylor and P. W. Wilson, *J. Am. Chem. Soc.*, 1973, **95**, 1834.
- J. Köhler and J.-H. Chang, *Z. Anorg. Allg. Chem.*, 1997, **623**, 596.
- (a) P. N. Bartlett, D. Cook, C. H. de Groot, A. L. Hector, R. Huang, A. Jolleys, G. P. Kissling, W. Levason, S. J. Pearce and G. Reid, *RSC Adv.*, 2013, **3**, 15645; (b) P. N. Bartlett, D. A. Cook, M. W. George, A. L. Hector, J. Ke, W. Levason, G. Reid, D. C. Smith and W. Zhang, *Phys. Chem. Chem. Phys.*, 2014, **16**, 9202.
- P. N. Bartlett, C. Y. Cummings, W. Levason, D. Pugh and G. Reid, *Chem. – Eur. J.*, 2014, **20**, 5019.
- G. P. Adams, J. L. Margrave and P. W. Wilson, *J. Inorg. Nucl. Chem.*, 1973, **33**, 1301.
- (a) J. Trotter, M. Akhtar and N. Bartlett, *J. Chem. Soc. A*, 1966, 30; (b) G. Denes, *J. Solid State Chem.*, 1989, **78**, 52.

