Faraday Discussions

Cite this: *Faraday Discuss.*, 2014, **168**, 185

PAPER

View Article Online
View Journal | View Issue

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Reaction kinetics and isotope effect of water formation by the surface reaction of solid H_2O_2 with H atoms at low temperatures

Yasuhiro Oba,* Kazuya Osaka, Naoki Watanabe, Takeshi Chigai and Akira Kouchi

Received 26th November 2013, Accepted 17th January 2014
DOI: 10.1039/c3fd00112a

We performed laboratory experiments on the formation of water and its isotopologues by surface reactions of hydrogen peroxide (H_2O_2) with hydrogen (H) atoms and their deuterated counterparts (D_2O_2, D) at 10–30 K. High-purity H_2O_2 (>95%) was prepared *in situ* by the codeposition of molecular oxygen and H atoms at relatively high temperatures (45–50 K). We determined that the high-purity H_2O_2 solid reacts with both H and deuterium (D) atoms at 10–30 K despite the large activation barriers (~2000 K). Moreover, the reaction rate for H atoms is approximately 45 times faster than that for D atoms at 15 K. Thus, the observed large isotope effect indicates that these reactions occurred through quantum tunneling. We propose that the observed HDO/H_2O ratio in molecular clouds might be a good tool for the estimation of the atomic D/H ratio in those environments.

Introduction

Water (H_2O) is the predominant solid constituent of icy layers of submicron-sized interstellar grains. Because of the potential importance of H_2O for chemical evolution in molecular clouds (MCs), elucidating the formation mechanism of H_2O in those environments is important. Although H_2O formation is possible by gas phase reactions at low temperatures,[1] the observed large abundance of H_2O cannot be explained only by the gas-phase synthesis.[2] Therefore, it is generally accepted that grain-surface reactions are crucial for producing H_2O in MCs.

Tielens and Hagen[3] proposed that H_2O formation is initiated by hydrogenation of atomic oxygen (O), molecular oxygen (O_2), and ozone (O_3), and is completed by the following reactions:

$$OH + H \rightarrow H_2O, \tag{1}$$

N19W8, Kita-ku, Sapporo, Hokkaido, Japan. E-mail: oba@lowtem.hokudai.ac.jp; Fax: +81 11 706 7142; Tel: +81 11 706 5477

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

$$OH + H_2 \rightarrow H_2O + H, \quad (2)$$

$$H_2O_2 + H \rightarrow H_2O + OH. \quad (3)$$

Since reaction (1) is a radical–radical reaction, it should proceed immediately once reactants encounter each other on the surface. This reaction has been studied experimentally by several groups.[4–7] In contrast to reaction (1), reactions (2) and (3) have large activation barriers (>2000 K) in the gas phase.[8,9] However, despite such a large barrier, these two reactions were proposed to contribute significantly to H_2O formation in dense MCs.[10] Since reactions having such large barriers do not occur thermally in MCs, reactions (2) and (3) require quantum tunneling. The quantum tunneling rate k_q is expressed by the following equation, assuming a rectangular activation barrier with a height E_a and width a:[11]

$$k_q \approx \nu_0 \exp[-(2a/\hbar)(2mE_a)^{1/2}], \quad (4)$$

where ν_0 and m represent the frequency of harmonic motion and the mass of the reaction, respectively. Since temperature is not included in the equation, k_q does not depend on the reaction temperature. Further details about quantum tunneling reactions have been described elsewhere.[11–14]

We have recently studied reaction (2) experimentally by the codeposition of nonenergetic OH with H_2 and isotopologues such as OD, HD, and D_2 on a substrate and determined that the reactions occur at 10 K.[14] In addition, significant isotope effects were observed, and reactions of OH and OD abstracting a D atom from HD and D_2 were approximately ten times slower than those abstracting an H atom from H_2 and HD. This isotope effect can be explained by the difference in the effective mass of tunneling reactions.[14]

A number of research studies have been conducted on reaction (3). In these previous studies, O_2 was used as an initial reactant rather than hydrogen peroxide (H_2O_2), which was produced by the successive hydrogenation of O_2 as follows:

$$O_2 + H \rightarrow HO_2, \quad (5)$$

$$HO_2 + H \rightarrow H_2O_2. \quad (6)$$

For example, Miyauchi *et al.*[15] exposed solid O_2 layers to H or D atoms at 10 K and determined that (i) the rate of O_2 hydrogenation (reaction (5)) is equal to that of O_2 deuteration, (ii) the rate of reaction (3) is slower than that of reaction (5), and (iii) the rate of reaction (3) is eight times faster than that of the following isotopically substituted reaction (7):

$$D_2O_2 + D \rightarrow D_2O + OD. \quad (7)$$

Miyauchi *et al.* considered that the rate difference between reactions (3) and (7) would be due to the isotope effect of quantum tunneling.[15] However, in the O_2 hydrogenation experiments, the formation of both H_2O_2 and H_2O occurs in the sample solid. Furthermore, the parent O_2 molecule is IR-inactive. Thus, multiparameter fittings, which often cause significant errors, are necessary to obtain the rates for reactions (3) and (5). Moreover, recent studies suggested that H_2O may form by another exothermic reaction in typical experimental conditions for O_2 hydrogenation:[16,17]

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

$$OH + OH \rightarrow H_2O + O. \quad (8)$$

In addition to reaction (3), OH is expected to form by the following pathway:

$$HO_2 + H \rightarrow H_2O_2^* \rightarrow 2OH, \quad (9)$$

where $H_2O_2^*$ is a reaction intermediate. The reaction of two OH also yields H_2O_2 on the surface:

$$OH + OH \rightarrow H_2O_2. \quad (10)$$

The branching ratio of barrierless reactions (8) to (10) was determined experimentally[17] to be 1 to 4 at 40 K and theoretically[18] to be 1 to 9 on a cold substrate. In any case, reactions (8) and (10) may compete during O_2 hydrogenation experiments to some extent, making it more difficult to obtain reliable kinetic parameters for reaction (3). In the case of CO hydrogenation experiments, a similar problem occurred because both formaldehyde (H_2CO) and methanol (CH_3OH) were produced in a single experiment.[19–23] However, additional experiments using H_2CO as an initial reactant have enabled us to better understand the reaction kinetics and isotope effects of CO and H_2CO hydrogenation.[13,24] Similarly, the use of H_2O_2 as an initial reactant is desirable for studying the kinetics and isotope effects of reaction (3). However, because of difficulty in using pure H_2O_2, to date such an experiment has not been performed.

In the present study, we performed experimental studies on the formation of H_2O *via* reaction (3) and its isotope effect using high-purity (>95%) solid H_2O_2 and D_2O_2.

Experimental

Apparatus and experimental conditions for water formation

All experiments were performed using the Apparatus for SUrface Reaction in Astrophysics (ASURA) system. The ASURA primarily comprises a main chamber and an atomic source. An aluminum (Al) substrate was mounted at the center of the main chamber and all reactions were performed on the substrate at 10–30 K. Hydrogen (H) and deuterium (D) atoms were produced by the dissociation of H_2 and D_2 molecules, respectively, in a microwave discharge plasma, and were cooled by multiple interactions with the inner wall of the aluminum pipe, which was cooled to 100 K. We confirmed that the formed H and D atoms were well thermalized to the pipe temperature.[25] Further details of the ASURA have been described elsewhere.[25,26]

The fluxes of H and D atoms were not directly measured in the present experimental setup; they were estimated by comparing the effective rates of CO hydrogenation and deuteration with those reported by Hidaka *et al.*,[27] which were obtained under the same experimental conditions. Briefly, amorphous H_2O ice (a-H_2O) with a thickness of approximately ten monolayers (ML; $\sim 10^{15}$ molecules cm^{-2}) was produced by vapor deposition on the substrate at 15 K, followed by the deposition of CO with a thickness of ~0.8 ML. The column density of H_2O and CO was calculated from the peak area and the previously published band strengths, as described by Hidaka *et al.*[27] The band strengths for the CO stretching of CO and OH

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

stretching of H_2O are 2.0×10^{-16} and 1.1×10^{-17} cm molecules^{-1}, respectively.[28] The obtained effective rates were 1.4 and 0.21 min^{-1} for the hydrogenation and deuteration of CO, respectively (see Results section below for the determination of effective rate constants). These values are a factor of 3.3 and 6.4 larger than the effective rates of CO hydrogenation and deuteration, respectively, compared to those reported by Hidaka *et al.*,[27] whose fluxes of both H and D atoms were 2.6×10^{14} atoms cm^{-2} s^{-1}. Assuming that the surface density of H and D atoms correlates linearly with their fluxes, the fluxes of H and D atoms in the present study were estimated to be 8.7×10^{14} and 1.7×10^{15} atoms cm^{-2} s^{-1}, respectively, which corresponds to a D and H atom flux ratio of ~2. The variations in the H and D fluxes are expected to be less than 10% during and between each experiment.

Approximately 1 ML of solid H_2O_2 or its deuterated counterpart D_2O_2 was produced on the substrate by the procedures shown in the next section. H_2O_2 and D_2O_2 were exposed to H (D) atoms at 10–30 K. Reaction products were monitored *in situ* by a reflection absorption Fourier Transform Infrared (FTIR) spectrometer with a resolution of 4 cm^{-1} in the spectral range from 700 to 4000 cm^{-1}. The column density of H_2O_2 and D_2O_2 was calculated from the peak area and previously published band strengths.[15] The band strengths used were 2.1×10^{-17} and 1.5×10^{-17} cm molecule^{-1} for the OH and OD bending bands at 1385 and 1039 cm^{-1}, respectively.

Experiments were also performed on an amorphous D_2O ice (a-D_2O; ~30 ML) vapor-deposited at 10 K. The band strength used was 1.3×10^{-16} cm molecule^{-1} for the OD stretching band.[15]

Preparation of high-purity solid hydrogen peroxide

In previous studies,[29,30] high-purity H_2O_2 (>97%) has been prepared by distilling commercially available H_2O_2 solution under vacuum. The distilled H_2O_2 was introduced into a reaction substrate through a transfer line made with nonreactive materials such as glass to avoid the catalytic decomposition of H_2O_2 on metal surfaces. However, in a typical apparatus, which comprises a stainless steel chamber and gas lines, it is not easy to obtain pure H_2O_2 using this procedure without significant modification.

Thus, we produced high-purity solid H_2O_2 *in situ* by the codeposition of H atoms with O_2 molecules on a substrate at relatively high temperatures. In our previous study, we determined that H_2O_2-rich ice tends to form by the O_2/H codeposition at high temperatures (>30 K) and with an increasing proportion of O_2 relative to H atoms.[31] For example, when O_2 and H were codeposited with an O_2/H ratio of $\sim 2 \times 10^{-3}$ at 20 K, the main product was H_2O with a small amount of H_2O_2 ($H_2O/H_2O_2 \sim 5$). When the same experiment was performed with an O_2/H ratio of $\sim 9 \times 10^{-3}$ at 40 K, the main product was H_2O_2 with ~15% contamination of H_2O. We further extended this experiment for the production of solid H_2O_2 with higher purity suitable for studying the kinetics and the isotope effect of reaction (3). One of the major advantages of this sample preparation method is that it is possible to study the isotope effect of reaction (3) using different isotopes (*i.e.*, H and D), which was not possible in previous studies.[15,32]

Gaseous O_2 was introduced into the main chamber through a capillary plate. From the pressure inside the main chamber, the O_2 flux was estimated to be 1.0×10^{14} molecules cm^{-2} s^{-1}, which is two to four orders of magnitude larger than

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

that used by Oba *et al.*[31] The H atoms were codeposited with O_2 onto the substrate at 45–50 K. In the present experiments, pure H_2O_2 solid and the solid on a-D_2O were used as reactants. The amount of solid H_2O_2 was ~1 ML. After the formation of solid H_2O_2, the microwave was switched off, the supply of all gases (H_2 and O_2) was stopped, and the substrate temperature was raised to 70 K to remove residual O_2 from the sample solid. H_2O contamination in the solid H_2O_2 was <5%, which was confirmed by the IR spectrum of the product (Figure 1a). We believe that this small amount of contamination does not significantly impact the kinetics of reaction (3). We confirmed this by a temperature-programmed desorption experiment wherein little O_2 remained on the substrate after the sample treatment. The produced H_2O_2 was then cooled to the desired temperatures (10–30 K) for hydrogenation or deuteration experiments. Moreover, when D atoms were used, high-purity D_2O_2 was formed (Figure 1b).

Results

H_2O_2 + H and H_2O_2 + D

Figure 2 shows IR absorption spectra of solid H_2O_2 (top) and H_2O (bottom) for comparison and the difference spectra of 1 ML pure solid H_2O_2 after H atom exposure for up to 10 min at 15 K (middle). In the difference spectra, the peaks below and above the baseline represent decreases of initial reactant and increase of reaction products, respectively. With increasing H atom fluence, the peak intensity for the OH bending of H_2O_2 at 1385 cm^{-1} decreased, and new peaks appeared at 3000–3600, 2850, and 1660 cm^{-1}. The peak at 1660 cm^{-1} is attributable to the OH bending of H_2O formed by reaction (3). Based on the peak position of OH stretching bands of solid H_2O_2 and H_2O (Figure 2), the H_2O band at 3000–3600 cm^{-1} has a substantial overlap with that of H_2O_2, and the peak shape would be attributable to the sum of H_2O_2 decrease and H_2O increase. In addition, H_2O_2 has another strong peak at 2827 cm^{-1}, which is often assigned to the $\nu_2 + \nu_6$ combination band.[33] If the amount of H_2O_2 decreases after H atom exposure, the intensity of the peak should also decrease. However, a peak was observed slightly above the baseline at 2850 cm^{-1} after H atom exposure, unlike other peaks at ~3300 and 1385 cm^{-1} (Figure 2). These apparently contradictory observations will be discussed later in the Discussion section. Two small peaks



Fig. 1 FTIR spectra of (a) H_2O_2 and (b) D_2O_2 produced at 45 K.

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.



Fig. 2 FTIR spectra of H_2O_2 (top) and pure H_2O (bottom) and the variations in the difference spectra of H_2O_2 after exposure to H atoms for up to 10 min at 15 K (middle).

appeared at 3697 and 3721 cm^{-1} after H atom exposure (Figure 2), which are attributed to the 3- and 2-coordinated water molecules, respectively.[34]

Figure 3 shows the difference spectra of 1 ML H_2O_2 after D atom exposure for up to 120 min at 15 K. If H_2O_2 reacts with D atoms, HDO is expected to form as a main product by the following reaction:

$$H_2O_2 + D \rightarrow HDO + OH. \tag{11}$$

After D atom exposure to H_2O_2, the peak area for OH stretching and bending bands decreased. New peaks appeared at 3490, 2527, and 1492 cm^{-1}, which are



Fig. 3 Variations in the difference spectra of H_2O_2 after exposure to D atoms for up to 120 min at 15 K.

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

typically observed for solid HDO at low temperatures.[35,36] This result clearly indicates that HDO was formed by reaction (11) at 15 K. Although the peak area for OH stretching and bending bands decreased after D atom exposure, the $\nu_2 + \nu_6$ combination band at 2860 cm^{-1} increased slightly. Very small peaks appeared at 2722 and 3693 cm^{-1} after D atom exposure (Figure 3), the former of which is probably derived from the dangling OD bond of HDO.[37] The latter peak could be the dangling OH bond of HDO; however, the assignment is uncertain because of low S/N.

Figure 4 plots variations in the column density of solid H_2O_2 normalized to the unexposed initial amount after exposure to H or D atoms. We fitted the plots in Figure 4 to the following single-exponential decay function to obtain the kinetic parameters for reactions (3) and (11):

$$\Delta[H_2O_2]_t/[H_2O_2]_0 = A(e^{-k_n[X]t} - 1), \tag{12}$$

where A is a saturation value, t is the H or D atom exposure time, k_n is the rate constant of reaction (n), and [X] is the number density of X atoms (X = H or D) on the surface. Unfortunately, it is difficult to measure [X] in the present experiment; thus, the product k_n[X] is obtained as a fitting parameter for equation (12). Hereafter, k_n[X] is denoted as the effective rate constant k_n' in the present study. We assume that during exposure, [X] is independent of time and is governed mainly by the balance between the flux of X atoms, the sticking coefficient of the impinging atoms, and the loss of atoms by X–X recombination. We obtained $k_3' = 7.2 \times 10^{-1}$ and $k_{11}' = 3.2 \times 10^{-2}$ min^{-1} at 15 K, and the k_3'/k_{11}' ratio was 23.

In the present study, we do not determine the absolute yields of reaction products with the decrease in the column density of reactants, because the band strengths of the products (H_2O or HDO) have been reported only for a transmission method. Using these reported values may cause a large error (<50%)[31] when those band strengths are used in a reflection method. However, it does not



Fig. 4 Variations in the column density of H_2O_2 normalized to the initial amount as a function of exposure time of H or D atoms at 15 K. Solid lines are single-exponential decay fits to the plots.

affect the values of effective rate constants because the term of band strength is not included in equation (12). In addition, a portion of the reaction products could desorb from the substrate upon formation,[38,39] making the interpretation of the yields of products more difficult.

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

D_2O_2 + H and D_2O_2 + D

Figure 5 shows variations in the difference spectra of pure solid D_2O_2 after exposure to H atoms on the Al substrate at 15 K. With increasing D atom fluence, the peak intensity of OD stretching and bending bands at 2467 and 1045 cm^{-1}, respectively, decreased, and new peaks appeared at 3455, 2587, and 1477 cm^{-1}. By comparing the peak positions with those given in the literature,[36] we determined that these new peaks are attributable to HDO, indicating that solid D_2O_2 reacted with H atoms to yield HDO at 15 K:

$$D_2O_2 + H \rightarrow HDO + OD. \quad (13)$$

The peak intensity of the $\nu_2 + \nu_6$ combination band for D_2O_2 (2126 cm^{-1})[33] increased slightly, which is opposite to the behavior that would be observed if D_2O_2 was consumed by reaction (13). The dangling OH band of HDO was observed at 3694 cm^{-1} while the dangling OD band was not confirmed probably due to low S/N (Figure 5).

Figure 6 shows variations in the difference spectra after D atom exposure to D_2O_2 for up to 120 min at 15 K. This experiment was performed to study reaction (7):

$$D_2O_2 + D \rightarrow D_2O + OD. \quad (7)$$

This reaction has been studied in previous O_2 deuteration experiments.[15,32] In the present study, with increasing D atom fluence, new peaks appeared at 2562 and



Fig. 5 Variations in the difference spectra of D_2O_2 after exposure to H atoms for up to 10 min at 15 K.

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.



Fig. 6 Variations in the difference spectra of D_2O_2 after exposure to D atoms for up to 120 min at 15 K.

1220 cm^{-1}, concurrent with the decrease of D_2O_2 (Figure 6). The new peaks can be attributed to the OD stretching and bending of D_2O, which clearly indicates that reaction (7) occurred to yield D_2O on the surface at 15 K. A small peak was observed after exposure to D atoms at 2727 cm^{-1}, which is assigned to the 3-coordinated dangling OD band of D_2O.[34] A 2-coordinated dangling OD band (at 2748 cm^{-1})[34] was not firmly identified, probably due to the low S/N of the spectrum. The peak intensity of the $\nu_2 + \nu_6$ combination band for D_2O_2 increased as well (Figure 6).

Figure 7 shows variations in the column density of solid D_2O_2 normalized to the unexposed initial amount after H or D atom exposure at 15 K. The relative abundance of D_2O_2 by reaction (13) reaches a saturation value of approximately



Fig. 7 Variations in the column density of D_2O_2 normalized to the initial amount as a function of H or D atom exposure times at 15 K. Solid lines are single-exponential decay fits to the plots.

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

−0.2 after a few minutes. In contrast, the decrease of D_2O_2 by reaction (7) was much slower, reaching the almost same saturation value after 150 min. By fitting the plots in Figure 7 into single exponential decay function (12) where $[H_2O_2]$ is replaced with $[D_2O_2]$, we obtained $k_{13}' = 8.9 \times 10^{-1}$ and $k_7' = 2.3 \times 10^{-2}$ min^{-1}, and the k_{13}'/k_7' ratio was 38.

Reactions on amorphous D_2O ice

Hydrogenation and deuteration of solid H_2O_2 and D_2O_2 were also performed on vapor-deposited a-D_2O with a thickness of ~30 ML at 15 K. Figure 8 shows an IR spectrum of 1 ML solid H_2O_2 produced on a-D_2O and the difference spectra after H atom exposure for up to 5 min. The peak area of the $\nu_2 + \nu_6$ combination band before H atom exposure was larger by a factor of two than that on the Al substrate, although the peak area of other bands (OH stretching and bending) was almost equal.

With increasing H atom fluence, the intensities of peaks at 3233, 2852, and 1392 cm^{-1} decreased and new peaks appeared at 3422 and 1635 cm^{-1}. These observations clearly indicate that H_2O was formed by reaction (13). Notably, the peak intensity for the $\nu_2 + \nu_6$ combination band at 2852 cm^{-1} decreased after H atom exposure. This behavior is straightforward since H_2O_2 was consumed by reaction (13); however, interestingly, this is opposite to the result for the same reactions of pure H_2O_2 on the Al substrate described above. In addition to reaction (13), we confirmed in separate experiments that reactions (11), (13), and (7) occur on a-D_2O at 15 K. Two small peaks appeared at 3697 and 3719 cm^{-1} after exposure to H atoms (Figure 8), which are attributed to the 3- and 2-coordinated dangling OH bands, respectively.[34]

We determined the effective rate constants with statistical errors for each reaction (Table 1) by fitting the attenuation of H_2O_2 and D_2O_2 into a single exponential decay function (12). As a general trend, values of k' are larger for reactions on a-D_2O.



Fig. 8 Variations in the difference spectra of H_2O_2 produced on a-D_2O for up to 5 min at 15 K.

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Table 1 Effective rate constants with statistical errors determined in the present study at 15 K

Reaction	Reaction number	Substrate	Effective rate (min^{-1})	E_a (K)[a]	Reduced mass (u)
H_2O_2 + H	3	Al	$7.2 \pm 0.6 \times 10^{-1}$	2508	0.97
		a-D_2O	$9.9 \pm 0.6 \times 10^{-1}$		
H_2O_2 + D	11	Al	$3.2 \pm 0.1 \times 10^{-2}$	2355	1.89
		a-D_2O	$4.0 \pm 0.6 \times 10^{-2}$		
D_2O_2 + H	13	Al	$8.9 \pm 1.1 \times 10^{-1}$	2540	0.97
		a-D_2O	$9.2 \pm 0.6 \times 10^{-1}$		
D_2O_2 + D	7	Al	$2.3 \pm 0.2 \times 10^{-2}$	2384	1.89
		a-D_2O	$2.2 \pm 0.8 \times 10^{-2}$		

[a] Taquet *et al.*[40]

The column densities of H_2O formed on the Al and a-D_2O by reaction (3) were calculated using the band strength of the OH-bending at 1635 cm^{-1} (1.2×10^{-17} cm $molecule^{-1}$).[28] We determined that the H_2O yield on a-D_2O was approximately two orders of magnitude larger than that on the Al at the same H fluence, although the amount of H_2O_2 consumption on H atom exposure was identical for a-D_2O and Al (Figure 9).

Temperature dependence of reaction kinetics

Reaction (3) was studied using pure H_2O_2 at 10, 20, and 30 K as well as at 15 K. Reaction (3) occurred at all temperatures. We obtained kinetic parameters for reaction (3) (k_3') at each temperature following the procedures described above. Figure 10 shows variations in the relative abundance of pure H_2O_2 after H atom exposure at 10–30 K. The temperature dependence shows that the saturation value of H_2O_2 becomes larger with increasing temperature up to 20 K; however, at



Fig. 9 Variations in the column densities of H_2O (circle) and H_2O_2 (square) normalized to the initial H_2O_2 amount obtained after H atom exposure to H_2O_2 at 15 K. Open and filled symbols represent experimental results on a-D_2O and Al substrate, respectively.

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.



Fig. 10 Variations in the column density of pure H_2O_2 normalized to the initial amount after exposure to H atoms at 10 (square), 15 (circle), 20 (triangle), and 30 (diamond) K. Lines are single-exponential decay fits to the plots.

30 K, the reaction is very slow and the saturation value is much less than that below 20 K. The effective rate constant is the largest at 10 K and decreases with increasing temperature. These features were also obtained for reactions on a-D_2O ice (Figure 11).

Discussion

Quantum tunneling and isotope effect

The reaction of H_2O_2 with H atoms and that of their deuterated counterparts has a large activation barrier (>2000 K) in the gas phase.[9,40,41] Therefore, these reactions are expected to proceed through quantum tunneling at 10–30 K even on the



Fig. 11 Temperature dependence of the effective rate constant for reaction (3): filled circle and open square symbols represent the rate for pure H_2O_2 and H_2O_2 on a-D_2O, respectively.

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

surface, as mentioned earlier. A tunneling reaction strongly depends on the transmission mass of the activation barrier of reaction, as can be seen in equation (4). Theoretical studies proposed that H_2O formation by reaction (3) initiates from the formation of an intermediate by H atom addition to one O atom in H_2O_2, followed by the cleavage of the O–O bond.[9,41] In general, the tunneling mass in the two-body addition reaction is described by the reduced mass μ.[12,13] As shown in Table 1, the mass dependence of the reaction rate is evident; the lighter mass results in a faster reaction rate. Thus, we conclude that the reactions of H_2O_2/D_2O_2 with H/D atoms proceed by quantum tunneling.

The ratios of the effective rate constants between reactions (3) and (11) (k_3'/k_{11}') are 23 and 25 for pure H_2O_2 and H_2O_2 on a-D_2O, respectively. Since the effective rate constant k_n' is expressed as k_n[X] where X = H or D, the ratio of the effective rate constants is not equal to the ratio of the actual reaction rate constants if [H] $\neq$ [D]. Assuming that the value of [X] has a linear correlation with the flux of X atoms in the present experiment, the ratios of k_3 to k_{11} (k_3/k_{11}) become 44 and 48 for pure H_2O_2 and H_2O_2 on a-D_2O, respectively, because the D atom flux is approximately twice that of the H atom flux. Similarly, the k_{13}/k_7 ratio was calculated to be 75 and 81 for pure D_2O_2 and D_2O_2 on a-D_2O, respectively. Taquet *et al.*[40] calculated a tunneling probability P_n through an Eckart potential barrier for reaction (n). In the present study, if we compare the P_3/P_{11} ratio with the k_3/k_{11} and the P_{13}/P_7 ratio with the k_{13}/k_7, we obtain $k_3/k_{11} = \sim 2 \times P_3/P_{11}$ and $k_{13}/k_7 = \sim 3 \times P_{13}/P_7$. These differences are not surprising because the tunneling probability is significantly affected by the shape of the potential barrier, and it is very difficult to determine the tunneling rate with an accurate barrier for surface reactions. In fact, Taquet *et al.*[40] noted that, although the Eckart model provides a significant improvement over square barriers, which were typically used for this type of calculation, the tunneling probabilities of some reactions at low temperatures can be underestimated (or overestimated).

We obtained the k_3/k_7 ratios of 61 and 88 for pure H_2O_2/D_2O_2 and H_2O_2/D_2O_2 on a-D_2O, respectively. These values are about one order of magnitude larger than those reported by Miyauchi *et al.*, who determined the k_3'/k_7' ratio to be 8 in their O_2 hydrogenation/deuteration experiments at 10 K. Since the flux of H atoms was the same with that of D atoms in the previous experiment,[15] the k_3'/k_7' ratio is certainly equal to the k_3/k_7 ratio on the assumption that the ratio of surface number densities between H and D atoms is same as that of the fluxes. The large difference may arise for several reasons. First, multi-parameter fittings may cause significant errors of k_3 and k_7 in their work. Second, H_2O and D_2O would form in O_2 hydrogenation/deuteration *via* separate pathways: OH + OH $\rightarrow$ H_2O + O and OD + OD $\rightarrow$ D_2O + O, respectively, which are both barrierless reactions. This would lead to an overestimation of the k_3/k_7 ratio. In the present study, these uncertainties are removed because H_2O_2 or D_2O_2 are used as the initial reactants. Thus, we believe that our calculated value of k_3/k_7 is more reliable than that obtained in O_2 hydrogenation/deuteration experiments.

Next, we compare the rates of reactions (3) and (5). The k_5/k_3 ratio in the O_2 hydrogenation experiment was reported to be 3.3.[15] Based on the fact that reaction (5) is barrierless and reaction (3) has a large barrier (>2000 K),[9] the value obtained by Miyauchi *et al.* seems to be rather small. Subsequently, we calculated the k_5/k_3 ratio using k_3' obtained in the present study. The value of k_5' reported by Miyauchi *et al.* (12.8 min^{-1})[15] was not used for this calculation

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

because their conditions such as atom fluxes and sample amounts were significantly different from ours. Instead, we used the value of k_5' (5.2 min^{-1}) obtained in the following experiment: solid O_2 (~3 ML) on an amorphous D_2O ice (30 ML) was exposed to H atoms (flux: 2×10^{14} atoms cm^{-2} s^{-1}) at 10 K.[42] After the correction of the flux difference, we obtained the k_5/k_3 ratio of ~15, which is approximately five times larger than that reported by Miyauchi *et al.*[15] Although the sample compositions and the fluxes of H atoms in these two experiments do not exactly match, we believe that the present result would better represent the k_5/k_3 ratio compared to previous studies.

Notably, the observed isotope effect in the hydrogen/deuterium addition to H_2O_2 (k_3/k_{11}) is much larger than that for one of the astrochemically-important tunneling surface reactions: the hydrogenation/deuteration of CO. Hidaka *et al.*[27] reported that the rate of CO hydrogenation (k_H) was larger than CO deuteration (k_D), with a k_H/k_D ratio of 12.5. This isotope effect is approximately four times smaller than that of the hydrogenation/deuteration of H_2O_2 observed in the present study. This large difference is rather surprising because the barrier height for the hydrogenation/deuteration of H_2O_2 is similar to that of CO.[40,43] We believe that the difference might have been due to accumulation of multiple factors such as differences in residence time of H/D atoms on H_2O_2 and CO and the shape of the potential barriers.

As mentioned previously, we assume that the surface density of H atoms is the same as that of D atoms when the flux is equal. However, in a series of experiments under high H and D flux conditions, this may not always be true. Even under the same flux conditions for H and D atoms, the surface density of D atoms ([D]) may become larger than that of H atoms ([H]). Since H atoms can diffuse faster on the surface than D atoms, the recombination probability for H atoms is higher than that for D atoms,[12] resulting in a lower value of [H] than that of [D] even under the same fluxes. In that case, the ratio of [H]/[D] becomes smaller than 0.5, yielding larger values of k_3/k_{11} and k_{13}/k_7 than those reported in the present study. Further studies related to the surface densities of H and D atoms are necessary.

Dependence of reaction efficiency on temperature and type of substrate

The effective rate constants for reaction (3) decrease with increasing substrate temperature (Figure 11). This is opposite to the typical Arrhenius-type behavior where the reaction rate has a positive correlation with temperature. In addition, because reaction (3) occurred through quantum tunneling, the reaction rate should have lesser dependence on the surface temperatures. The decrease of the effective rate can be explained well by the decrease in the number density of H atoms on the surface with increasing temperature because the effective rate k_3' is expressed by k_3[H]. This trend was also observed in CO hydrogenation and deuteration, H_2CO hydrogenation, H–D substitution of CH_3OH, and O_2 hydrogenation on low-temperature surfaces.[11,42]

The difference of the effective rate constant between reactions of pure H_2O_2 and H_2O_2 on a-D_2O is the largest at 10 K and decreases with increasing temperature (Figure 11). The large difference observed at lower temperatures (<15 K) may be explained by the difference in the number density of H atoms on the surface because a-D_2O would have a much larger surface area compared to planar Al

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

substrate, as reported for CO hydrogenation by Hidaka *et al.*[44] who determined that the surface area for a-H_2O with thickness equivalent to ~17 ML is approximately nine times larger than that for crystalline H_2O. In contrast, at elevated temperatures where the residence time of H atoms on the surface is very short even on a-D_2O, the difference could little be affected by the density of H atoms, which can lead to similar effective rate constant values under both conditions (Figure 11).

In Figure 9, H_2O yield for the sample of pure H_2O_2 on Al substrate is lower than that for H_2O_2 on a-D_2O, although the decrease of H_2O_2 is approximately the same for both samples. This difference may be partly explained by immediate desorption of H_2O formed as a product at the reaction preferentially occurred on the Al substrate. This behavior is also observed in O_2 + D experiments on amorphous silicates and a-H_2O, where the yield of D_2O is higher for reactions on a-H_2O.[39] The heat of reaction (3) (285 kJ mol^{-1} = 2.9 eV) is partly partitioned to the reaction product H_2O and OH. This energy would be sufficient for H_2O to desorb from the substrate (Al or a-D_2O). For the H_2O_2 on a-D_2O sample, interaction of H_2O_2 and H_2O product with the a-D_2O surface is much stronger than that with the Al surface because of hydrogen bonding. Therefore, the reaction heat can dissipate into a-D_2O more easily, and desorption at the reaction surface may be reduced. Furthermore, since a-D_2O has a large surface area because of pores and cracks, the desorbed H_2O can be retrapped on the surface of a-D_2O. This type of H_2O relaxation and trap has been demonstrated both experimentally[45–47] and theoretically[48,49] to occur during the photodesorption of H_2O from bulk a-H_2O. In addition to H_2O, OH should also form by reaction (3) and can be the source of another H_2O formation. Accordingly, H_2O and OH trapped on the surface of a-D_2O may explain the larger column density of the formed H_2O than that on the Al substrate (Figure 9).

As shown in Figure 1, solid H_2O_2 produced on the Al substrate has representative strong absorptions at three positions: 1385 (ν_2), 2827 ($\nu_2 + \nu_6$), and 3326 cm^{-1} (ν_1).[33] Similarly, for solid H_2O_2 on a-D_2O, these three peaks are slightly shifted on the spectrum at 1392, 2852, and 3233 cm^{-1} (Figure 8). The difference in the peak position for each band is not very large (<25 cm^{-1}). However, the peak area of the $\nu_2 + \nu_6$ combination band was larger by a factor of two for the sample on a-D_2O, although the peak area of other bands (ν_1 and ν_2) was identical on both substrates. A relatively strong peak at ~2850 cm^{-1} for solid H_2O_2 is typically assigned to the $\nu_2 + \nu_6$ combination band;[33,50] however, some studies considered that this assignment is disputable.[51,52] Ignatov *et al.*[51] proposed that this peak is attributable to an OH valence band of the H_2O_2 molecule placed in the unusual surface environment. The present results imply that the peak intensity at ~2850 cm^{-1} might correlate with the configuration of H_2O_2 on the substrate. In addition, the fact that H_2O_2 forms hydrogen bonds more on a-D_2O than on Al could result in the different IR features at ~2850 cm^{-1}. In other words, the number of hydrogen bonds between H_2O_2 and surrounding H_2O may constrain the peak intensity at ~2850 cm^{-1}. If this is the case, it could explain the contradictory behavior of the peak intensity after exposure to atoms (*e.g.* Figures 2 and 8). More detailed experimental and theoretical studies are necessary for the precise peak assignment. Thus, we propose that care should be taken when this IR peak is used for quantification of solid H_2O_2 in various environments.

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Dangling OH (OD) bands of reaction products

We confirmed that the formed solid H_2O and its isotopologues (HDO and D_2O) show the dangling OH (OD) bands (*e.g.*, Figure 2). The presence of dangling OH bands in the IR spectrum of solid H_2O typically indicates that it is amorphous and has a microporous structure.[34] However, we have reported that dangling OH bands are not observed in solid H_2O formed by codeposition of O_2 and H atoms at 10 K and that the formed ice is amorphous but has a compact structure.[31] These different results suggest that the formed ice structure depends on experimental conditions and that the properties of the reaction products are strikingly different between each study (multilayers with O_2 contaminants *vs.* very thin layers with little contaminants).

Astrophysical implication

Deuterated water (HDO and D_2O) has been identified in the gas phase with HDO/H_2O and D_2O/H_2O ratios up to the order of 10^{-2} and 10^{-3}, respectively, toward protostars.[53–57] In contrast, deuterated water was not clearly identified in the solid phase; only the upper limit of HDO was determined (HDO/H_2O < 0.2%–2%).[58,59] Despite the detection of deuterated water in the gas phase only, it is reasonable to consider that deuterated water is also produced by surface reactions on interstellar grains at very low temperatures.

The following reactions are possible to yield water isotopologues from H_2O_2 and its isotopologues:

$$H_2O_2 + H \rightarrow H_2O + OH, \tag{3}$$

$$H_2O_2 + D \rightarrow HDO + OH, \tag{11}$$

$$D_2O_2 + H \rightarrow HDO + OD, \tag{13}$$

$$D_2O_2 + D \rightarrow D_2O + OD, \tag{7}$$

$$HDO_2 + H \rightarrow HDO + OH, \tag{14a}$$

$$HDO_2 + H \rightarrow H_2O + OD, \tag{14b}$$

$$HDO_2 + D \rightarrow HDO + OD, \tag{15a}$$

$$HDO_2 + D \rightarrow D_2O + OH. \tag{15b}$$

We did not study reactions (14) and (15) because of difficulty in producing high-purity HDO_2. Two types of reaction products are possible for reactions (14) and (15). Since the reduced mass of a reaction to produce the intermediate does not depend on the type of products, we expect that reactions (14a) and (14b) occur to the same extent statistically. In fact, the tunneling probability for reaction (14a) is calculated to be identical with that of reaction (14b).[40] The same is true for reactions (15a) and (15b). Thus, for simplicity, we made a rough assumption based on the obtained results that H atom addition reactions such as reactions (3) and (13) occur faster by a factor of 50 than D atom addition reactions such as reactions (7) and (11). In addition, we assume that the atomic D/H ratio is

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

constant and the surface diffusion rates of H and D atoms are the same. Under these assumptions, we determined that the HDO/H_2O ratio was almost identical to the atomic D/H ratio, while the D_2O formation was negligible ($\sim 10^{-6}$ that of H_2O). On the other hand, assuming no isotope effect on the reactions, the HDO/H_2O ratio becomes twice as large as the atomic D/H ratio; thus, quantum tunneling would cause a decrease of the HDO/H_2O ratio in this reaction pathway. In other words, if quantum tunneling corrections are not included in a chemical model, the obtained result should overestimate the value of the HDO/H_2O ratio.

We extend this discussion to other water formation pathways, *i.e.*, reactions (1) ($OH + H \rightarrow H_2O$) and (2) ($OH + H_2 \rightarrow H_2O + H$). For reaction (2), we have experimentally determined the relative efficiency of reactions for all possible isotopologues;[14] H atom abstraction reactions are approximately ten times more efficient than D atom abstraction reactions. Assuming that OH and OD are formed by surface reactions O + H and O + D, respectively, HD/H_2 ratio is 10^{-5}, and no D_2 is present, the HDO/H_2O ratio was very much consistent with the atomic D/H ratio.

In contrast to the former two reaction pathways, the reactions of OH or OD with H or D atoms do not have an activation barrier. Thus, the product ratio would strongly depend on the atomic D/H ratio. Since there are two possible reactions to yield HDO ($OH + D \rightarrow HDO$ and $OD + H \rightarrow HDO$), statistically the HDO/H_2O ratio is twice as large as the atomic D/H ratio. With regard to the formation of D_2O, this reaction pathway is the most favorable among the three reaction pathways; the D_2O/H_2O ratio is statistically the square of the value of the D/H ratio. For example, the D_2O/H_2O ratio is 10^{-4} if the atomic D/H is 10^{-2}. This value is two to four orders of magnitude higher than that estimated by other pathways under the same assumptions. Therefore, the OD + D reaction is the only possible pathway to effectively yield D_2O in MCs.

Moreover, it is important to note that, on grain surfaces, the deuterium fractionation of water occurs only during the formation by surface reactions at the typical temperature of MCs ($\sim$10 K). This is supported by the fact that H_2O does not react with D atoms at <15 K[60] and thermal H–D exchange does not occur at <100 K,[61] which prevents the D-enrichment by H–D substitution with other deuterium-enriched species after the formation of H_2O on the grains. The reactivity of water with D atoms at low temperatures differs from that of organic species such as CH_3OH,[25,60] H_2CO,[13,27] and CH_3NH_2[62] where H–D exchange occurs for these molecules by reacting with D atoms at temperatures as low as 10 K.

Based on the present and previous experimental results for water formation, we suggest that the HDO/H_2O ratio might be a key parameter to estimate the atomic D/H ratio during the formation of water. Namely, the following relationship is roughly derived from experimental studies: $(D/H)_{atom} \leq (HDO/H_2O) \leq 2(D/H)_{atom}$ or $1/2(HDO/H_2O) \leq (D/H)_{atom} \leq (HDO/H_2O)$, where $(D/H)_{atom}$ and (HDO/H_2O) represent the atomic D/H on grains and the HDO/H_2O ratio formed by surface reactions, respectively. Note that these relationships were derived with reference to experimental studies for surface reactions. In addition to surface reactions, deuterium fractionation of water by gas phase reactions may also be possible.[63] Moreover, energetic processes induced by UV and cosmic rays might cause hydrogen isotopic fractionation of water during its decomposition and interactions with other ice components, both of which may modify the HDO/H_2O ratio. Therefore, we propose that further collaborative theoretical and

experimental studies on surface and gas phase chemistries are necessary to construct a complete chemical model regarding the evolution of the water D/H ratio in MCs.

Acknowledgements

The authors thank Drs H. Hidaka and T. Hama for fruitful discussions at the earlier stages of manuscript preparation. We also thank an anonymous referee for providing constructive comments. This work is partly supported by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science. Y.O. has received funding from the Kurita Water and Environment Foundation.

References

1 E. Herbst and W. Klempere, *Astrophys. J.*, 1973, **185**, 505–533.
2 T. I. Hasegawa, E. Herbst and C. M. Leung, *Astrophys. J. Suppl.*, 1992, **82**, 167–195.
3 A. Tielens and W. Hagen, *Astron. Astrophys.*, 1982, **114**, 245–260.
4 F. Dulieu, L. Amiaud, E. Congiu, J. H. Fillion, E. Matar, A. Momeni, V. Pirronello and J. L. Lemaire, *Astron. Astrophys.*, 2010, **512**, 5.
5 K. Hiraoka, T. Miyagoshi, T. Takayama, K. Yamamoto and Y. Kihara, *Astrophys. J.*, 1998, **498**, 710–715.
6 D. P. Jing, J. He, M. Bonini, J. R. Brucato and G. Vidali, *J. Phys. Chem. A*, 2013, **117**, 3009–3016.
7 D. P. Jing, J. He, J. Brucato, A. De Sio, L. Tozzetti and G. Vidali, *Astrophys. J.*, 2011, **741**, 5.
8 R. Atkinson, D. L. Baulch, R. A. Cox, J. N. Crowley, R. F. Hampson, R. G. Hynes, M. E. Jenkin, M. J. Rossi and J. Troe, *Atmos. Chem. Phys.*, 2004, **4**, 1461–1738.
9 H. Koussa, M. Bahri, N. Jaidane and Z. Ben Lakhdar, *THEOCHEM*, 2006, **770**, 149–156.
10 H. M. Cuppen and E. Herbst, *Astrophys. J.*, 2007, **668**, 294–309.
11 N. Watanabe and A. Kouchi, *Prog. Surf. Sci.*, 2008, **83**, 439–489.
12 T. Hama and N. Watanabe, *Chem. Rev.*, 2013, **113**, 8783–8839.
13 H. Hidaka, M. Watanabe, A. Kouchi and N. Watanabe, *Astrophys. J.*, 2009, **702**, 291–300.
14 Y. Oba, N. Watanabe, T. Hama, K. Kuwahata, H. Hidaka and A. Kouchi, *Astrophys. J.*, 2012, 749.
15 N. Miyauchi, H. Hidaka, T. Chigai, A. Nagaoka, N. Watanabe and A. Kouchi, *Chem. Phys. Lett.*, 2008, **456**, 27–30.
16 H. M. Cuppen, S. Ioppolo, C. Romanzin and H. Linnartz, *Phys. Chem. Chem. Phys.*, 2010, **12**, 12077–12088.
17 Y. Oba, N. Watanabe, A. Kouchi, T. Hama and V. Pirronello, *Phys. Chem. Chem. Phys.*, 2011, **13**, 15792–15797.
18 T. Lamberts, H. M. Cuppen, S. Ioppolo and H. Linnartz, *Phys. Chem. Chem. Phys.*, 2013, **15**, 8287–8302.
19 G. W. Fuchs, H. M. Cuppen, S. Ioppolo, C. Romanzin, S. E. Bisschop, S. Andersson, E. F. van Dishoeck and H. Linnartz, *Astron. Astrophys.*, 2009, **505**, 629–639.
20 C. Pirim and L. Krim, *Chem. Phys.*, 2011, **380**, 67–76.

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

21 N. Watanabe and A. Kouchi, *Astrophys. J.*, 2002, **571**, L173–L176.
22 N. Watanabe, T. Shiraki and A. Kouchi, *Astrophys. J.*, 2003, **588**, L121–L124.
23 N. Watanabe, A. Nagaoka, T. Shiraki and A. Kouchi, *Astrophys. J.*, 2004, **616**, 638–642.
24 H. Hidaka, N. Watanabe, T. Shiraki, A. Nagaoka and A. Kouchi, *Astrophys. J.*, 2004, **614**, 1124–1131.
25 A. Nagaoka, N. Watanabe and A. Kouchi, *J. Phys. Chem. A*, 2007, **111**, 3016–3028.
26 N. Watanabe, A. Nagaoka, H. Hidaka, T. Shiraki, T. Chigai and A. Kouchi, *Planet. Space Sci.*, 2006, **54**, 1107–1114.
27 H. Hidaka, A. Kouchi and N. Watanabe, *J. Chem. Phys.*, 2007, 126.
28 P. A. Gerakines, W. A. Schutte, J. M. Greenberg and E. F. van Dishoeck, *Astron. Astrophys.*, 1995, **296**, 810–818.
29 M. J. Loeffler and R. A. Baragiola, *J. Phys. Chem. A*, 2011, **115**, 5324–5328.
30 R. G. Smith, S. B. Charnley, Y. J. Pendleton, C. M. Wright, M. M. Maldoni and G. Robinson, *Astrophys. J.*, 2011, **743**, 13.
31 Y. Oba, N. Miyauchi, H. Hidaka, T. Chigai, N. Watanabe and A. Kouchi, *Astrophys. J.*, 2009, **701**, 464–470.
32 S. Ioppolo, H. M. Cuppen, C. Romanzin, E. F. van Dishoeck and H. Linnartz, *Astrophys. J.*, 2008, **686**, 1474–1479.
33 J. A. Lannon, F. D. Verderam and R. W. Anderson, *J. Chem. Phys.*, 1971, **54**, 2212–2223.
34 V. Buch and J. P. Devlin, *J. Chem. Phys.*, 1991, **94**, 4091–4092.
35 J. P. Devlin, *J. Mol. Struct.*, 1990, **224**, 33–43.
36 D. F. Hornig, H. F. White and F. P. Reding, *Spectrochim. Acta*, 1958, **12**, 338–349.
37 J. P. Devlin, *J. Chem. Phys.*, 2000, **112**, 5527–5529.
38 R. T. Garrod, V. Wakelam and E. Herbst, *Astron. Astrophys.*, 2007, **467**, 1103–1115.
39 H. Chaabouni, M. Minissale, G. Manico, E. Congiu, J. A. Noble, S. Baouche, M. Accolla, J. L. Lemaire, V. Pirronello and F. Dulieu, *J. Chem. Phys.*, 2012, **137**, 234706.
40 V. Taquet, P. S. Peters, C. Kahane, C. Ceccarelli, A. Lopez-Sepulcre, C. Toubin, D. Duflot and L. Wiesenfeld, *Astron. Astrophys.*, 2013, **550**, 23.
41 B. A. Ellingson, D. P. Theis, O. Tishchenko, J. Zheng and D. G. Truhlar, *J. Phys. Chem. A*, 2007, **111**, 13554–13566.
42 Y. Oba, N. Miyauchi, T. Chigai, H. Hidaka, N. Watanabe and A. Kouchi, in *Physics and Chemistry of Ice 2010*, Hokkaido University Press, 2011, Y. Furukawa, G. Sazaki, T. Uchida, and N. Watanabe, Eds., 361–368.
43 D. E. Woon, *Astrophys. J.*, 2002, **569**, 541–548.
44 H. Hidaka, N. Miyauchi, A. Kouchi and N. Watanabe, *Chem. Phys. Lett.*, 2008, **456**, 36–40.
45 T. Hama, M. Yokoyama, A. Yabushita, M. Kawasaki, S. Andersson, C. M. Western, M. N. R. Ashfold, R. N. Dixon and N. Watanabe, *J. Chem. Phys.*, 2010, **132**, 8.
46 A. Yabushita, T. Hama, M. Yokoyama, M. Kawasaki, S. Andersson, R. N. Dixon, M. N. R. Ashfold and N. Watanabe, *Astrophys. J.*, 2009, **699**, L80–L83.
47 K. I. Oberg, H. Linnartz, R. Visser and E. F. van Dishoeck, *Astrophys. J.*, 2009, **693**, 1209–1218.

Open Access Article. Published on 22 May 2014. Downloaded on 11/16/2025 10:53:12 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

48 S. Andersson and E. F. van Dishoeck, *Astron. Astrophys.*, 2008, **491**, 907–916.
49 C. Arasa, S. Andersson, H. M. Cuppen, E. F. van Dishoeck and G. J. Kroes, *J. Chem. Phys.*, 2010, **132**, 12.
50 A. Engdahl, B. Nelander and G. Karlstrom, *J. Phys. Chem. A*, 2001, **105**, 8393–8398.
51 S. K. Ignatov, A. G. Razuvaev, P. G. Sennikov and O. Schrems, *THEOCHEM*, 2009, **908**, 47–54.
52 P. G. Sennikov, S. K. Ignatov and O. Schrems, *ChemPhysChem*, 2005, **6**, 392–412.
53 H. M. Butner, S. B. Charnley, C. Ceccarelli, S. D. Rodgers, J. R. Pardo, B. Parise, J. Cernicharo and G. R. Davis, *Astrophys. J.*, 2007, **659**, L137–L140.
54 C. Ceccarelli, C. Dominik, E. Caux, B. Lefloch and P. Caselli, *Astrophys. J.*, 2005, **631**, L81–L84.
55 A. Coutens, C. Vastel, E. Caux, C. Ceccarelli, S. Bottinelli, L. Wiesenfeld, A. Faure, Y. Scribano and C. Kahane, *Astron. Astrophys.*, 2012, **539**, 12.
56 B. Parise, E. Caux, A. Castets, C. Ceccarelli, L. Loinard, A. Tielens, A. Bacmann, S. Cazaux, C. Comito, F. Helmich, C. Kahane, P. Schilke, E. van Dishoeck, V. Wakelam and A. Walters, *Astron. Astrophys.*, 2005, **431**, 547–554.
57 C. Vastel, C. Ceccarelli, E. Caux, A. Coutens, J. Cernicharo, S. Bottinelli, K. Demyk, A. Faure, L. Wiesenfeld, Y. Scribano, A. Bacmann, P. Hily-Blant, S. Maret, A. Walters, E. A. Bergin, G. A. Blake, A. Castets, N. Crimier, C. Dominik, P. Encrenaz, M. Gerin, P. Hennebelle, C. Kahane, A. Klotz, G. Melnick, L. Pagani, B. Parise, P. Schilke, V. Wakelam, A. Baudry, T. Bell, M. Benedettini, A. Boogert, S. Cabrit, P. Caselli, C. Codella, C. Comito, E. Falgarone, A. Fuente, P. F. Goldsmith, F. Helmich, T. Henning, E. Herbst, T. Jacq, M. Kama, W. Langer, B. Lefloch, D. Lis, S. Lord, A. Lorenzani, D. Neufeld, B. Nisini, S. Pacheco, J. Pearson, T. Phillips, M. Salez, P. Saraceno, K. Schuster, X. Tielens, F. van der Tak, M. H. D. van der Wiel, S. Viti, F. Wyrowski, H. Yorke, P. Cais, J. M. Krieg, M. Olberg and L. Ravera, *Astron. Astrophys.*, 2010, **521**, 5.
58 E. Dartois, W. F. Thi, T. R. Geballe, D. Deboffle, L. d'Hendecourt and E. van Dishoeck, *Astron. Astrophys.*, 2003, **399**, 1009–1020.
59 B. Parise, T. Simon, E. Caux, E. Dartois, C. Ceccarelli, J. Rayner and A. Tielens, *Astron. Astrophys.*, 2003, **410**, 897–904.
60 A. Nagaoka, N. Watanabe and A. Kouchi, *Astrophys. J.*, 2005, **624**, L29–L32.
61 J. P. Devlin and V. Buch, *J. Chem. Phys.*, 2007, **127**, 4.
62 Y. Oba, T. Chigai, Y. Osamura, N. Watanabe and A. Kouchi, *Meteorit. Planet. Sci.*, 2014, **49**, 117–132.
63 H. Roberts, E. Herbst and T. J. Millar, *Astron. Astrophys.*, 2004, **424**, 905–917.