



Cite this: *Chem. Commun.*, 2015, 51, 6682

Correction: A FRET based pH probe with a broad working range applicable to referenced ratiometric dual wavelength and luminescence lifetime read out

Robert J. Meier,^a Johann M. B. Simbürger,^a Tero Soukka^b and Michael Schäferling^{*c}

DOI: 10.1039/c5cc90142a

www.rsc.org/chemcomm

Correction for 'A FRET based pH probe with a broad working range applicable to referenced ratiometric dual wavelength and luminescence lifetime read out' by Robert J. Meier *et al.*, *Chem. Commun.*, 2015, DOI: 10.1039/c5cc00144g.

Complexes 1 and 4 in Fig. 1 were displayed incorrectly in the original manuscript. There was one CH₂ group missing in the binding arms to europium.




Fig. 1 (a) Synthesis of the molecular FRET system consisting of a sensitized europium chelate Eu7D (**1**) containing an isothiocyanate function and CNF (**3**) linked by 1,12-aminododecane (**2**). (b) Red luminescence emission of the europium chelate (dashed line) and absorption of CNF at different pH (from 10 to 3, colored lines).

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Institute of Analytical Chemistry, Chemo- and Biosensors, University of Regensburg, 93040 Regensburg, Germany

^b Department of Biochemistry/Biotechnology, University of Turku, 20520 Turku, Finland

^c BAM - Federal Institute for Materials Research and Testing, Division 1.10 Biophotonics, 12489 Berlin, Germany. E-mail: michael.schaeferling@bam.de