PCCP

PAPER



Cite this: Phys. Chem. Chem. Phys., 2015, 17, 19230

Received 17th April 2015, Accepted 17th June 2015

DOI: 10.1039/c5cp02248g

www.rsc.org/pccp

1. Introduction

The coordination of transition metal atoms M, cations M⁺ or polar salts M⁺X⁻ to alkynes and alkenes has been described in terms of the interaction of the π electrons of the latter compounds with the π^* orbitals of the metal atom or ion.^{1,2} Such interactions achieve importance by virtue of their presence in coordination compounds that feature in hydrogenation and polymerization reactions.³ For that reason, we have recently been investigating systematically a series of simple, gas-phase complexes of the type $C_2H_2\cdots MX$ which contain such interactions, where M is a coinage metal atom and X is a halogen atom.^{4–6} The advantage of working in the gas phase is that distortions of the alkyne on complex formation are free from lattice or solvent effects and are therefore intrinsic to the interaction of interest. The technique employed in the synthesis of $C_2H_2\cdots MX$ complexes is laser ablation of the metal M in the



View Article Online View Journal | View Issue

Distortions of ethyne when complexed with a cuprous or argentous halide: the rotational spectrum of $C_2H_2\cdots CuF^{\dagger}$

Daniel P. Zaleski,^a Susanna L. Stephens,^a David P. Tew,^b Dror M. Bittner,^a Nicholas R. Walker*^a and Anthony C. Legon*^b

A new molecule $C_2H_2\cdots$ CuF has been synthesized in the gas phase by means of the reaction of laser-ablated metallic copper with a pulse of gas consisting of a dilute mixture of ethyne and sulfur hexafluoride in argon. The ground-state rotational spectrum was detected by two types of Fourier-transform microwave spectroscopy, namely that conducted in a microwave Fabry–Perot cavity and the chirped-pulse broadband technique. The spectroscopic constants of the six isotopologues ${}^{12}C_2H_2\cdots {}^{63}Cu^{19}F$, ${}^{12}C_2H_2\cdots {}^{65}Cu^{19}F$, ${}^{12}C_2D_2\cdots {}^{65}Cu^{19}F$ and ${}^{12}C_2D_2\cdots {}^{65}Cu^{19}F$ were determined and interpreted to show that the molecule has a planar, T-shaped geometry belonging to the molecular point group $C_{2\nu}$, with CuF forming the stem of the T. Quantitative interpretation reveals that the ethyne molecule is distorted when subsumed into the complex in such manner that the C=C bond lengthens (by δr) and the two H atoms cease to be collinear with the C=C internuclear line. The H atoms move symmetrically away from the approaching Cu atom of CuF, to increase each *=C-H angle by $\delta A = 14.65(2)^{\circ}$, from 180° to 194.65(2)°. *Ab initio* calculations at the explicitly-correlated level of theory CCSD(T)(F12*)/aug-cc-pVTZ lead to good agreement with the experimental geometry. It is shown that similar distortions δr and δA , similarly determined, for four complexes $C_2H_2\cdots MX$ (M = Cu or Ag; X = F, Cl or CCH) are approximately linearly related to the energies D_e for the dissociation process $C_2H_2\cdots MX = C_2H_2 + MX$.

presence of both ethyne and a source of halogen atom X in a preponderating excess of argon gas. The subsequent, rapid supersonic expansion of the mixture into a vacuum cools and stabilizes the product molecules and allows their rotational spectra to be observed. By these methods it has been possible to determine, with all the precision associated with microwave spectroscopy, the geometries of several species^{4–6} $C_2H_2\cdots MX$, where MX is AgCl, AgCCH, or CuCl.

In this article, we report the rotational spectra of six isotopologues of $C_2H_2\cdots Cu$ –F. The complexes were produced by the laser ablation of metallic copper in the presence of a pulse of ethyne–SF₆–Ar gas mixture and their spectra were detected by two versions of the Fourier-transform microwave technique. Interpretation of the spectroscopic constants thereby determined leads to a precise characterization of the distortion that ethyne undergoes when it is coordinated to CuF. The distortions will be compared with those predicted by an *ab initio* calculation conducted at the CCSD(T)(F12*)/AVTZ level of theory Such distortions have now been established, both experimentally and *ab initio*, as accompanying the attachment of C_2H_2 to MX = CuF, CuCl, AgCl and AgCCH (where M is copper or silver, and X = F, Cl or CCH as appropriate). Variations in these distortions with (1) M and (2) X will be considered.

^a School of Chemistry, Newcastle University, Bedson Building, Newcastle-upon-Tyne, NE1 7RU, UK. E-mail: nick.walker@newcastle.ac.uk

^b School of Chemistry, University of Bristol, Cantock's Close, Bristol, BS8 1TS, UK. E-mail: a.c.legon@bristol.ac.uk

[†] Electronic supplementary information (ESI) available. See DOI: 10.1039/ c5cp02248g

2. Experimental and theoretical methods

Experimental data for $C_2H_2\cdots$ CuF and $C_2D_2\cdots$ CuF were recorded on a chirped-pulse Fourier transform microwave spectrometer⁷ operating between 7 and 18.5 GHz. The instrument has been previously described in detail.^{8,9} A gas sample was prepared containing approximately 1% C_2H_2 (or C_2D_2 as appropriate) and 1% SF₆ balanced in 6 bar argon. The gas mixture then pulsed over a copper target rod which was ablated with a Nd:YAG laser. The gas pulse was subsequently expanded into a vacuum chamber. For these experiments, the pulsed nozzle was placed perpendicular to the axis of microwave propagation. During each sample injection cycle, 8 free induction decays (FID's) were measured, each 20 µs in duration. For $C_2H_2\cdots$ CuF a total of 480 k FID's were acquired (16 h), for $C_2D_2\cdots$ CuF the total was 780 k (26 h), and 720 k (24 h) for ¹³C₂H₂-CuF. In each case, Fourier transformation used a digital Kaiser–Bessel window function.

For ${}^{12}C_2H_2\cdots$ CuF, data were also recorded on a Balle–Flygare cavity spectrometer, 10,11 with an equivalent gas sample. These measurements were acquired with the pulsed nozzle placed coaxial with the direction of microwave propagation. Transitions measured in this arrangement appear as Doppler doublets, each normally with a full-width half maximum (FWHM) of 5 kHz (but see Section 3.1); this is compared to a FWHM of approximately 150 kHz for the perpendicular arrangement found in the broadband instrument. Only samples containing isotopes in natural abundance were employed when using this spectrometer.

Geometry optimizations were performed using CCSD(T)(F12*),¹² a coupled-cluster method with single and double excitations, explicit correlation,¹³ and a perturbative treatment of triple excitations.¹⁴ An AVTZ basis set combination was used, by which we mean that the aug-cc-pVTZ basis sets¹⁵ were used for the C, F and H atoms and the aug-cc-pVTZ-PP basis for Cu, in combination with the ECP-10-MDF effective core potential on Cu to account for scalar relativistic effects.^{16,17} The frozen-core approximation was used throughout, and all calculations were performed using the MOLPRO package.¹⁸ The equilibrium dissociation energy D_e for the process $C_2H_2\cdots Cu-F = C_2H_2 + Cu-F$ was also computed at the CCSD(T)-(F12*)/AVTZ level using the counterpoise correction method¹⁹ where, for numerical stability, the CABS singles correction was not included in the correction term. The result was $D_e = 170$ kJ mol⁻¹.

For the free CuF, the bond length was optimised on a series of potential curves including successive corrections for core correlation, computed at the CCSD(T) level by using the cc-pwCVQZ basis, full triples at the frozen-core level with a cc-pVQZ basis and perturbative quadruples using the cc-pVTZ basis, following a similar procedure to that employed by Gauss *et al.*²⁰ Post CCSD(T) calculations were performed using the MRCC program²¹ and yielded the fully corrected bond length $r_e(\text{Cu-F}) = 1.7430$ Å, with a basis set uncertainty of 0.0006 Å. This result is in good agreement with the experimental value 1.74493 Å calculated from equilibrium rotational constants²² of ⁶³Cu¹⁹F and ⁶⁵Cu¹⁹F by using the expression $r_e = \{h/8\pi^2\mu B_e\}^{\frac{1}{3}}$. At the CCSD(T)(F12*)/aug-cc-pVTZ level of theory, as used for C₂H₂…Cu–F, the equilibrium bond length for the free CuF molecule is predicted to be 1.7422(25) Å,

where the error is that estimated (*via* gaussian error propagation) to arise from basis set incompleteness (0.0007 Å), core valence correlation error (0.0021 Å) and higher-order correlation (0.0011 Å).

3. Results

3.1 Determination of spectroscopic constants

For all six isotopologues investigated, only vibrational groundstate, *a*-type R-branch transitions $(J + 1)_{K_{-1}K'_1} \rightarrow J_{K_{-1}K''_1}$ with $K_{-1} = 0$ or 1 of the asymmetric-rotor complex $C_2H_2\cdots Cu$ –F were observed under the experimental conditions described in Section 2. Each transition carried a resolvable nuclear quadrupole hyperfine structure resulting from a single quadrupolar nucleus ⁶³Cu or ⁶⁵Cu (I = 3/2). Transition frequencies of each isotopologue were fitted by means of the program PGOPHER²³ with the following choice of Hamiltonian:

$$H = H_{\rm R} - \frac{1}{6} \mathbf{Q}({\rm Cu}): \nabla \mathbf{E}({\rm Cu}), \tag{1}$$

in which Q(Cu) and $\nabla E(Cu)$ are the Cu nuclear electric quadrupole tensor and the electric field gradient tensor at the Cu nucleus, respectively. H_R is the familiar Hamiltonian for a semi-rigid, asymmetric-rotor molecule and contains both rotational constants and centrifugal distortion constants. The *H* matrix was constructed in the coupled symmetric rotor basis, with the Watson A reduction²⁴ chosen for H_R , In addition to the rotational constants, H_R contains terms involving the five quartic centrifugal distortion constants $\Delta_{f_L} \Delta_{f_K}$, δ_{f_L} , and δ_K .

The rotational constants of the most abundant isotopologue ${}^{12}C_{2}H_{2}\cdots {}^{63}Cu^{19}F$ are large enough that only one group of $J + 1 \rightarrow J$ transitions, namely that with J = 1, could be measured in the frequency range available. A set of internally consistent arguments will be presented to show that the geometry of $C_{2}H_{2}\cdots Cu-F$ is of the planar, T-shaped type in which the non-covalent interaction of the two component molecules involves primarily the Cu atom of CuF and the π bond of ethyne, as shown in Fig. 1. The experimental consequences of such a geometry are as follows:

(1) The molecule is a nearly prolate, planar asymmetric rotor of molecular point group C_{2v} , with the *a* axis coincident with the C_2 axis. The rotational constant A_0 is close in magnitude to the rotational constant B_0 of free ethyne. Any difference of the two rotational constants will provide quantitative information about the extent of any geometrical distortion of the ethyne molecule when subsumed into $C_2H_2\cdots Cu-F$.

(2) The ground-state molecule will have a small positive inertia defect

$$\Delta_0 = I_c^0 - I_b^0 - I_a^0, \tag{2}$$

which is an important criterion of molecular planarity.

(3) A rotation C_2^a exchanges a pair of equivalent protons (I = 1/2) and this endows $K_{-1} = 1$ transitions with a nuclear spin statistical weight of 3 relative to that of 1 for $K_{-1} = 0$ transitions. This effect is evident in the set of $2_{12} \rightarrow 1_{11}$, $2_{02} \rightarrow 1_{01}$, $2_{11} \rightarrow 1_{10}$ transitions, as can be seen from the recording of these transitions shown in Fig. 2. Similar arguments applied to



Fig. 1 The geometry (drawn to scale) of $C_2H_2\cdots Cu-F$ determined from analysis of the rotational spectra of six isotopologues. The atoms are coplanar and the principal inertia axis *a* coincides with the C_2 axis of symmetry. The ethyne subunit is significantly distorted relative to the free molecule, both in respect of the angle *A* and of the distance $r(C \equiv C)$. See text for discussion of the detailed geometry.

the ${}^{12}C_2D_2\cdots Cu{}^{19}F$ and ${}^{13}C_2H_2\cdots Cu{}^{19}F$ isotopologues require the corresponding ratios to be 1:2 and 6:10, respectively, which were indeed as observed.

(4) Isotopic substitution of 63 Cu by 65 Cu should leave A_0 unchanged.

It is evident that the rotational constant A_0 has a central role to play in drawing conclusions about the symmetry and geometry of $C_2H_2\cdots Cu$ -F. Unfortunately, the $(J+1)_{K_{-1}K'_1} \rightarrow J_{K_{-1}K''_1}$ $(K_{-1} = 0 \text{ or } 1)$ transition frequencies have a very weak dependence on A_0 and it was not possible from the measured transition frequencies to determine A_0 independently of the two centrifugal distortion constants Δ_J and Δ_{JK} . (There is insignificant dependence on the remaining centrifugal distortion constants.) This difficulty was circumvented by the following approach.

It has been shown elsewhere that centrifugal distortion constants of molecules similar to $C_2H_2\cdots Cu$ -F can be calculated *ab initio* with good accuracy.⁶ For example, in the closely related species $C_2H_2\cdots Cu$ -Cl calculations at the MP2/cc-pVTZ level of theory reproduced Δ_J and Δ_{JK} to within a few standard deviations of the well-determined experimental values. Here, the five quartic constants $(\Delta_J, \Delta_{JK}, \Delta_K, \delta_J \text{ and } \delta_{JK})$ for $C_2H_2\cdots Cu$ -F were calculated for the optimized geometry at the higher level of theory MP2/aug-cc-pVQZ with the aid of the Gaussian electronic structure package²⁵ for each isotopologue investigated. The calculated constants Δ_J and Δ_{JK} were then fixed in the PGO-PHER²³ fits of observed frequencies to determine the rotational constants λ_{a_0} and C_0 and the Cu nuclear quadrupole coupling constants χ_{aa} (Cu) and χ_{bb} (Cu) – χ_{cc} (Cu) for the isotopologues ¹²C₂H₂···⁶³Cu¹⁹F and ¹²C₂H₂···⁶⁵Cu¹⁹F in the first instance.



Fig. 2 A recording (10 × vertical magnification) of the three $J = 2 \rightarrow 1$ transitions of ${}^{12}C_{2}H_{2}...{}^{63}Cu^{19}F$ and ${}^{12}C_{2}H_{2}...{}^{65}Cu^{19}F$, showing Cu nuclear quadrupole hyperfine structure (upper panel). The downward pointing spectrum is that synthesized by using PGOPHER and the spectroscopic constants given in Table 1 (2 K rotational temperature assumed). This simulation does not include nuclear spin statistics, so the difference between the simulation and observed spectrum shows clearly the presence of nuclear spin statistics in the latter. Even though hyperfine splitting is greater in the two outer transitions $2_{12} \rightarrow 1_{11}$ and $2_{11} \rightarrow 1_{10}$, their statistical weight advantage (3:1) over the central $2_{02} \rightarrow 1_{01}$ transition is evident. The lower panel gives an expanded version of the $2_{12} \rightarrow 1_{11}$ transitions of ${}^{12}C_{2}H_{2}...{}^{63}Cu^{19}F$ (blue) and ${}^{12}C_{2}H_{2}...{}^{65}Cu^{19}F$ (red). The simulated spectrum (with appropriate scaling of intensities) is again downward pointing.

The spectroscopic constants so determined both from measurements made with the Balle-Flygare and the chirped-pulse spectrometers are displayed in Table 1. The residuals of the fit to frequencies collected with the Balle-Flygare spectrometer were poorer than the usual few kHz. This arises because C_2H_2 ···CuF carries four nuclei (H, H, Cu and F) with magnetic moments of significant magnitude, in addition to the electric quadrupolar moment of the Cu nucleus. Moreover, only the lowest *I* transitions were available because of the relatively large rotational constants. The magnetic coupling of these nuclei (Cu, F spin-spin coupling, for example) can lead to further complicated but minor, partially resolved splitting or broadening and/or lineshape distortion of each Cu nuclear quadrupole component and is most significant at the lowest J. Therefore assignment of the magnetic sub-structure was not possible in the observed transitions of C_2H_2 . CuF, particularly those with $K_{-1} = 1$, and hence larger than normal residuals in the fit of the Cu nuclear quadrupole hyperfine structure resulted. When measured with the lower resolution, chirped-pulse instrument, fitted frequencies assigned to Cu quadrupole components gave similar magnitude residuals. Reassuringly, the spectroscopic constants determined independently with the two spectrometers are identical within experimental error.

Measurements for ${}^{13}C_2H_2 \cdots {}^{63,65}Cu^{19}F$, and ${}^{12}C_2D_2 \cdots {}^{63,65}Cu^{19}F$ were made exclusively on the chirped-pulse instrument to conserve the more expensive isotopic materials. The spectroscopic constants

Table 1 Ground-state spectroscopic constants of ${}^{12}C_2H_2$... ${}^{63,65}CuF$ as determined with two different pulsed-jet, Fourier-transform microwave spectrometers (Balle–Flygare and chirped-pulse types)

	Balle–Flygare		Chirped-pulse		
Spectroscopic constant	$^{12}C_2H_2$ ···· ⁶³ CuF	$^{12}\mathrm{C}_{2}\mathrm{H}_{2}\!\cdots^{65}\mathrm{CuF}$	$^{12}\mathrm{C_2H_2}{}^{\cdots}{}^{63}\mathrm{CuF}$	$^{12}C_2H_2$ ··· ⁶⁵ CuF	
A ₀ /MHz	34 749(83)	34 930(270)	35 030(230)	35 020(180)	
B ₀ /MHz	3431.4407(48)	3430.5201(55)	3431.4404(56)	3430.5159(42)	
C_0/MHz	3116.7391(44)	3115.9722(55)	3116.7348(56)	3115.9752(39)	
Δ_{IK}/kHz^a	[38.3]	[38.2]	[38.3]	[38.2]	
$\Delta_{\rm I}/{\rm kHz}^a$	0.61	0.61	0.61	0.61	
$\chi_{aa}(Cu)/MHz$	69.991(47)	64.721(53)	69.979(53)	64.720(40)	
$\{\gamma_{bb}(Cu) - \gamma_{cc}(Cu)\}/MHz$	-79.17(11)	-73.25(12)	-79.17(13)	-73.224(73)	
N^{b}	20	16	18	19	
$\sigma_{ m r.m.s.}/ m kHz^c$	30	33	34	26	

^{*a*} Values calculated at the MP2/aug-cc-pVQZ level of theory and fixed in the fit. ^{*b*} Number of nuclear quadrupole components included in the fit. ^{*c*} Standard deviation of fit.

for these species were obtained in a similar way and are recorded in Table 2. Detailed fits are available as Supplementary Data²⁶ at http://dx.doi.org/xxxxx. The effect of changes in the assumed values of the centrifugal distortion constants were investigated for the isotopologue ¹²C₂H₂...⁶³Cu¹⁹F. Changes of 10% in the assumed values of the distortion constants Δ_J and Δ_{JK} lead to changes in *A* that are small compared with the errors shown in Table 1. The standard deviations in A_0 values are higher than those identified in other B...MX studies because only one $J + 1 \rightarrow J$ transition is available within the bandwidth of the spectrometer in the present work.

It is significant that values of the A_0 for ${}^{12}C_2H_2...{}^{63}Cu^{19}F$ and ${}^{12}C_2H_2...{}^{65}Cu^{19}F$ (Table 1) are equal within experimental error (as required if Cu lies on the *a* axis). These values of A_0 and their associated uncertainties strongly imply that the average of the two results provides the most reliable value for this quantity, namely 34 840(90) MHz (where the quoted error is the range about the mean). This value will be used in geometry determinations set out in Section 3.2, together with the mean value $A_0 = 33100(150)$ MHz for the pair (${}^{13}C_2H_2...{}^{63}Cu^{19}F$, ${}^{13}C_2H_2...{}^{65}Cu^{19}F$) and $A_0 = 25182(26)$ MHz for the pair (${}^{12}C_2D_2...{}^{65}Cu^{19}F$), ${}^{12}C_2D_2...{}^{65}Cu^{19}F$). We note that $A_0 =$ 34 840(90) MHz is smaller by 435(90) MHz than the B_0 value of free ${}^{12}C_2H_2$ (see Table 3 for various properties of isotopologues of the free molecules ethyne 27,28 and cuprous fluoride 22). Similar conclusions (see Table 2) apply to the pair (${}^{13}C_2H_2...{}^{63}Cu^{19}F$, ¹³C₂H₂···⁶⁵Cu¹⁹F), for which the mean decrease from free ¹³C₂H₂ is 464(150) MHz, and the pair (¹²C₂D₂···⁶³Cu¹⁹F, ¹²C₂D₂···⁶⁵Cu¹⁹F), for which a mean decrease of 236(26) MHz relative to *B*₀ of free ¹²C₂D₂ is observed. These decreases, although not well determined, indicate a change in the ethyne geometry when it becomes attached to CuF. Moreover, the very small change in all three rotational constants on substitution of ⁶³Cu by ⁶⁵Cu establishes that the Cu atom lies close to the centre of mass and therefore that the order along the *a* axis is *–Cu–F, where * indicates the centre of the C≡C bond.

The Cu nuclear quadrupole coupling constants are reasonably well determined. The ratio of the values $\chi_{aa}({}^{63}Cu)/\chi_{aa}({}^{65}Cu) =$ 1.0814(16) lies within experimental error of the known ratio²⁹ $Q(^{63}Cu)/Q(^{65}Cu) = 1.0806(3)$ of the nuclear electric quadrupole moments of the copper nuclides. The large difference between $\chi_{bb}(Cu) = -74.58$ MHz and $\chi_{cc}(Cu) = 4.59$ MHz is evidence of a large anisotropy in the electric field gradient at Cu along the b and cprincipal inertia axis directions arising from the facts that (1) the Cu atom is close to the ethyne π bond and (2) the electron distribution within the triple bond $C \equiv C$ is different along the two directions. Indeed, the large difference $\{\chi_{bb}(^{63}Cu) - \chi_{cc}(^{63}Cu)\} =$ -79.17(11) MHz provides convincing evidence of the proximity of Cu to the C \equiv C bond. In the related molecule C₂H₂...Ag–Cl, in which Cl is remote from the ethyne π bond, values of the Cl nuclear quadrupole coupling constants are $\chi_{aa}(^{35}\text{Cl}) = -28.9268(78)$ MHz and $\{\chi_{bb}(^{35}\text{Cl}) - \chi_{cc}(^{35}\text{Cl})\} = -3.256(7)$ MHz.

Table 2 Ground-state spectroscopic constants of ${}^{13}C_2H_2$... ${}^{63,65}CuF$ and ${}^{12}C_2D_2$... ${}^{63,65}CuF$ as determined with a chirped-pulse, pulsed-jet, Fourier-transform microwave spectrometer

Spectroscopic constant	$^{13}\mathrm{C}_{2}\mathrm{H}_{2}\!\cdots^{63}\mathrm{CuF}$	$^{13}\mathrm{C}_{2}\mathrm{H}_{2}\!\cdots^{65}\mathrm{CuF}$	$^{12}\mathrm{C}_{2}\mathrm{D}_{2}\!\cdots^{63}\mathrm{CuF}$	$^{12}\mathrm{C}_{2}\mathrm{D}_{2}\!\cdots^{65}\mathrm{CuF}$
A ₀ /MHz	33 250(240)	32 950(290)	25 208(19)	25 156(32)
B ₀ /MHz	3302.7667(62)	3301.4981(98)	3260.5869(86)	3259.257(14)
C ₀ /MHz	2996.4700(43)	2995.4204(47)	2881.9524(88)	2880.939(11)
Δ_{IK}/kHz^a	[34.7]	[34.7]	[30.9]	[30.8]
Δ_{I}/kHz^{a}	0.56	0.56	0.53	0.53
$\gamma_{aa}(Cu)/MHz$	70.012(59)	64.756(71)	70.051(55)	64.743(56)
$\{\gamma_{bb}(Cu) - \gamma_{cc}(Cu)\}/MHz$	-79.23(17)	-73.28(24)	-80.4(38)	-69.8(39)
N ^b	20	15	17	13
$\sigma_{\rm rms}/\rm kHz^{c}$	33.8	32.2	23.7	22.8

^{*a*} Values calculated at the MP2/aug-cc-pVQZ level of theory and fixed in the fit. ^{*b*} Number of nuclear quadrupole components included in the fit. ^{*c*} Standard deviation of fit.

Table 3	Some spectroscopic constants and	I bond lengths of C ₂ H ₂ and Cu	F referred to in evaluating properties of C ₂ H ₂ ···CuF
---------	----------------------------------	--	--

Quantity constant	$^{12}C_{2}H_{2}$	$^{13}C_{2}H_{2}$	$^{12}C_2D_2$	⁶³ Cu ¹⁹ F	⁶⁵ Cu ¹⁹ F
Ethyne P/MHz	$25274,0602(54)^{a}$	22564 005 ^a	25419 6201 ^a	_	_
Geometry	$r_0(C_2H_2)^b$	$r_0(C_2D_2)^c$	$r_{\rm e}^{d}$		
$r(C \equiv C)/Å$	1.206553(6)	(1.206553)	1.20286(3)	—	—
r(С–H)/Å	1.06238(2)	1.06011(3)	1.06166(6)	_	—
Cuprous fluoride				_	_
B_0/MHz^e		—	—	11325.8910	11245.2404
$B_{\rm e}/{\rm MHz}^e$	_	_	—	11374.2092	11293.0426
$r_0(\text{Cu-F})/\text{\AA}$	—	—	—	1.74865	1.74863^{\prime}
<i>r</i> _e (Cu–F)/Å (experiment)	—	—	—	1.74493 ^g	1.74493^{g}
$r_{\rm e}$ (Cu–F)/Å (<i>ab initio</i> calc.)	_	_	_	$1.7430(3)^{n}$	

^{*a*} Ref. 27. ^{*b*} Calculated by fitting the moments of inertia I_b^0 of ${}^{12}C_2H_2$ and ${}^{13}C_2H_2$ from ref. 27. See text. ^{*c*} Calculated by fitting the moment of inertia I_b^0 of ${}^{12}C_2D_2$ with $r(C \equiv C)$ fixed at 1.206553 Å. ^{*d*} Ref. 28. ^{*e*} Ref. 22. ^{*f*} Calculated from B_0 value by using $r_0 = \{h/8\pi^2\mu B_0\}^{\frac{1}{2}}$, with B_0 estimated from Dunham constants in ref. 22. ^{*g*} Calculated from B_e value by using $r_e = \{h/8\pi^2\mu B_e\}^{\frac{1}{2}}$ with B_e from ref. 22. ^{*h*} Calculated at the CCSD(T)(F12*)/aug-cc-pwCVQZ level of theory, with post-CCSD(T) corrections made for basis set incompleteness, core-valence correlation, and higher-order correlations.

3.2 Molecular geometry

Of the four consequences of a planar geometry of the type shown in Fig. 1, it was indicated in Section 3.1 that three were satisfied by experimental results for $C_2H_2\cdots Cu$ –F. The fourth is that the inertia defect, as defined in eqn (2), should be small and positive. If the Cu atom does lie on the *a* axis, it is reasonable to use the mean I_a^0 value for a pair of isotopologues ${}^mC_2{}^nH_2\cdots {}^{63}Cu^{19}F$ and ${}^mC_2{}^nH_2\cdots {}^{65}Cu^{19}F$ in calculating Δ_0 for both members of the pair. The results are $\Delta_0 = 0.365(38)$ u Å² and 0.366(38) u Å² for the ${}^{12}C_2H_2\cdots {}^{63,65}Cu^{19}F$ pair, 0.373(69) u Å² and 0.293(22) u Å² for the ${}^{12}C_2D_2\cdots {}^{63,65}Cu^{19}F$ pair. The very similar values, observed across a similar range of isotopologues, for each of the related planar molecules $C_2H_2\cdots Cu$ –Cl⁶ and $C_2H_2\cdots Ag$ –Cl⁵ reinforce the conclusion that $C_2H_2\cdots Cu$ –F is also planar.

Thus, all the evidence so far presented is consistent with a geometry of $C_2H_2\cdots$ CuF of the planar, T-shaped type of $C_{2\nu}$ symmetry illustrated in Fig. 1, with the atoms in the indicated order. Quantitative detail of the geometry can be determined experimentally in two ways from the zero-point rotational constants of the six isotopologues of $C_2H_2\cdots$ CuF investigated, namely a partial r_s geometry and an almost complete r_0 geometry. These results will be compared with the full r_e geometry obtained from *ab initio* calculations carried out at the CCSD(T)(F12*/AVTZ) level of theory.

If equilibrium rotational constants (and therefore equilibrium principal moments of inertia) of the various isotopologues of $C_2H_2\cdots$ CuF investigated were available, the equilibrium principalaxis co-ordinates a_C , b_C , a_H , and a_{Cu} could be determined by means of eqn (3)–(5). In view of the molecular planarity and the existence of the symmetry operation C_2^a , the changes I_b and I_a in the equilibrium principal moments of inertia that accompany the double isotopic substitution of X by X' (X = ${}^{12}C$ or H; X' = ${}^{13}C$ or D, respectively) at symmetrically equivalent atoms are related to the equilibrium coordinates a_X and b_X by the expressions put forward by Chutjian³⁰

$$|a_{\rm X}| = (\Delta I_b / \mu_{\rm D})^{\frac{1}{2}},\tag{3}$$

$$|b_{\mathbf{X}}| = \{\Delta I_a / (2\Delta m)\}^{\frac{1}{2}},\tag{4}$$

in which $\mu_{\rm D} = (2\Delta mM)/(M + 2\Delta m)$ is the reduced mass for the double substitution in the parent isotopologue of mass *M* leading to a mass change $2\Delta m$. The equilibrium principal axis co-ordinate $a_{\rm Cu}$ of the atom Cu, which lies on the *a*-axis of $C_2H_2\cdots$ CuF, is given in terms of changes ΔI_b and ΔI_c in equilibrium principal moments of inertia accompanying isotopic substitution at Cu by Kraitchman's equation³¹

$$|a_{\rm Cu}| = \{ (\Delta I_b + \Delta I_c) / 2\mu_{\rm S} \}^{\frac{1}{2}}, \tag{5}$$

where $\mu_{\rm S} = (\Delta m M)/(M + \Delta m)$ is the reduced mass for the substitution. When equilibrium quantities are unavailable, Costain^{32,33} proposed using zero-point ΔI^0 values in place of their equilibrium counterparts ΔI^{e} in eqn (3)–(5) and named the resulting coordinates as substitution or r_s coordinates. The r_s coordinates $a_{\rm C}$, $b_{\rm C}$, $a_{\rm H}$, $b_{\rm H}$ and $a_{\rm Cu}$ so obtained from eqn (3)–(5) are given in Table 4. Each refers to the principal inertia axes of ${}^{12}C_2H_2 \cdots {}^{63}Cu^{19}F$ as the parent molecule and the chosen signs are those that lead to reasonable bond lengths. Note that the mean of A₀ for pairs of ^{63,65}Cu isotopologues, as given in Section 3.1, were used in the calculation of the $b_{\rm C}$ and $b_{\rm H}$ coordinates. The substantial uncertainties in $b_{\rm C}$ and $b_{\rm H}$ result from the relatively large errors assigned to A_0 . A r_s coordinate a_F is not available because fluorine possesses only one stable isotope and so the value of $a_{\rm F}$ given in Table 4 was determined, as recommended by Costain,³² from the other r_s coordinates with the aid of the first moment condition,

$$a_{\rm F} = -\left(\sum_{j} m_{j} a_{j}\right) \middle/ m_{\rm F},\tag{6}$$

in which the sum over j indicates that all atoms but F are included.

The significantly larger magnitude of $a_{\rm H}$ than $a_{\rm C}$ provides experimental evidence that the angular geometry of ethyne is distorted in the manner shown in Fig. 1, that is the H atoms move away from the C \equiv C internuclear line when the complex is formed, with the two equivalent angles $A = \angle \text{H-C} \equiv *$ (see Fig. 1 for definition) exceeding 180°. The value obtained from

Table 4	Various types of	f principal inertia a>	is coordinate and geometry	y determined for C₂H₂···Cu−	۰F
---------	------------------	------------------------	----------------------------	-----------------------------	----

	$r_{\rm s}$ geometry		r_0 geometry		$r_{\rm e}$ geometry ^{<i>a</i>}	
	Coordinate ^b		Coordinate		Coordinate	
Atom	a/Å	b/Å	a/Å	b/Å	a/Å	b/Å
H C Cu F	-1.9764(8) -1.7066(9) 0.142(10) 1.893(33)	$egin{array}{c} \pm 1.663(9) \ \pm 0.62(4) \ 0.0000 \ 0.0000 \end{array}$	-1.9767(4) -1.7059(4) 0.1415(3) 1.8961(6)	$egin{array}{c} \pm 1.6587(1) \ \pm 0.6231(1) \ 0.0000 \ 0.0000 \end{array}$	-1.9762 -1.6947 0.1412 1.8803	$\pm 1.6501 \\ \pm 0.6182 \\ 0.0000 \\ 0.0000$
Distance/angle	r _s geo	metry	r ₀ geomet	ry or angle	r _e geo	metry
$r(C \equiv C)/Å$ r(C-H)/Å r(Cu-*)/Å r(Cu-F)/Å Angle $A/^{\circ}$	1.24 1.07 1.85 1.75 194.5	(4) (7(4) (1) (1(10) (5(6)	1.246 [1.07 1.847 1.754 194.6	61(1) 7041] ^c 74(7) 47(9) 655(2)	1.2. 1.0 1.8 1.7 195	363 697 359 392 5.3

^{*a*} The r_e geometry was optimized at the CCSD(T)(F12*)/aug-cc-pVTZ level of theory. ^{*b*} The errors in r_s coordinates a_x are those given by $\delta a = (0.0015 \text{ Å})/a$, as suggested by Costain in ref. 32. The errors in r_s coordinates b_x are those propagated from the poorly determined rotational constants A_0 . ^{*c*} Assumed value obtained by adding the difference $r_0(C-H) - r_e(C-H) = 0.00072$ Å of the quantities for free ethyne to $r_e(C-H)$ calculated for $C_2H_2\cdots Cu-F$ and given in the final column. See text for discussion.

the $r_{\rm s}$ coordinates in Table 3 is $A = 194.5(6)^{\circ}$ (see Table 4), where the large error is consequent upon the errors in the A_0 values. This distortion is established with greater accuracy in the experimental r_0 and *ab initio* $r_{\rm e}$ geometries discussed later (see below). The bond distances implied by the $r_{\rm s}$ coordinates are also included in Table 4.

When the minimum number of bond lengths and angles required to establish a molecular geometry uniquely is fitted to a sufficient number of zero-point principal moments of inertia by the least-squares method, the result is called a r_0 geometry, *i.e* one obtained when the rovibrational contributions $\varepsilon_a = I_a^0 - I$ $_{a}^{e}$, etc. to the moments of inertia are ignored. A fit of I_{b}^{0} and I_{c}^{0} (values of I_a^0 were excluded because of their relatively low accuracy) of all six isotopologues of C₂H₂···Cu-F implied by the rotational constants B_0 and C_0 reported in Tables 1 and 2 to determine $r(C \equiv *)$, r(*-Cu), r(Cu-F), r(C-H), and $A = \angle H-C \equiv *$ was unsatisfactory, however, because of high correlation between two last-named quantities. To break the correlation, the following approach was used. A good estimate of r_0 (C–H) is given by adding r_{e} (C–H) of C₂H₂···Cu–F from the CCSD(T)(F12*)/AVTZ calculation to the difference r_0 (C–H) – r_e (C–H) = 0.00072 Å for free ethyne^{27,28} (Table 3), with the result $r_0(C-H) = 1.07041$ Å. This value was then fixed in the fit. The geometry and principal axis coordinates were then determined under this assumption with the aid of Kisiel's program STRFIT³⁴ and are included in Table 4. There is excellent agreement with the corresponding $r_{\rm s}$ quantities, given the large error in the $b_{\rm s}$ coordinates propagated through the rotational constants A_0 .

4. Discussion

The new molecule $C_2H_2\cdots Cu$ -F has been synthesized by interaction of the plasma produced by laser ablation of a copper rod with a supersonically-expanded pulse of gas containing SF₆ and C₂H₂ diluted with a large excess of argon. It was detected by means of its ground-state rotational spectrum and was shown to have a planar, T-shaped geometry of C_{2v} symmetry, with the CuF molecule lying along a C_2 axis of the C_2H_2 molecule. Quantitative aspects of the molecular geometry were determined through the investigation of six isotopologues. A significant finding is that the ethyne subunit is considerably distorted by the interaction of its C \equiv C bond with the Cu atom of CuF. The H atoms move from the C \equiv C internuclear line, in the direction away from the approaching Cu atom, so that the * \equiv C-H angle increases from 180° to 194.65(2)°. The C \equiv C bond lengthens by *ca*. 0.04 Å. These distortions are well reproduced in the optimized geometry of $C_2H_2\cdots$ Cu–F obtained from *ab initio* calculations carried out at the CCSD(T)(F12*)/AVTZ level of theory (see Table 4).

Similar distortions have now been observed in the series of molecules $C_2H_2\cdots CuF$, $C_2H_2\cdots AgCl$, $C_2H_2\cdots AgCCH$, and $C_2H_2\cdots$ CuCl,⁶ each of which has the planar T-shaped, C_{2v} geometry with the metal atom adjacent to the ethyne π bond (see Fig. 1). The changes δA in the angle $* \equiv$ C–H and δr in the distance $r(C \equiv C)$ are $[14.65(2)^{\circ}, 4.0(1) \text{ pm}], [7.70(4)^{\circ}, 2.86(4)]$ pm], [5.9(1)°, 2.61(3) pm] and [12.5(2)°, 2.7(3) pm] respectively, along the series. The corresponding energy changes De accompanying dissociation into C₂H₂ and MX have been calculated ab initio [at the CCSD(T)(F12*)/AVTZ or AVDZ levels] to be 170, 97.5, 92 and 148 kJ mol⁻¹, respectively. Fig. 3 displays a plot of each of δA and δr versus D_e for these molecules. The origin has been included as a point in each plot because when the interaction energy $D_{\rm e}$ is zero, presumably there is no distortion of the ethyne subunit. Fig. 3 shows that there is an approximate linear relationship between each of the two types of distortion δA and δr and the strength of the interaction $D_{\rm e}$.

The nature of the ethyne distortions can be understood with the aid of a simple model. The nuclei of each of the $C_2H_2 \cdots M-X$ (M = Cu or Ag, X = F or Cl) molecules all lie in the *ab* principal inertia plane. Let the symmetry plane of one of the ethyne π bonding orbitals lie in this plane, with that of the other π orbital perpendicular to it. When M–X approaches from large separation and takes up its position along a C_2 axis of ethyne (which then becomes the *a* axis of the new molecule), the partial



Fig. 3 Plots of the angular distortion δA (see Fig. 1 for a definition of the angle δA) and the extension δr of the C \equiv C bond of ethyne on formation of the complexes $C_2H_2\cdots MX$ against the energy D_e required for the dissociation reaction $C_2H_2\cdots MX = C_2H_2 + MX$. δA and δr are experimentally determined while the D_e values are calculated *ab initio* at the CCSD(T)(F12*)/AVTZ or AVDZ level of theory. The origin is included as a point under the assumption that zero interaction energy is associated with $\delta A = 0$ and $\delta r = 0$.

positive charge δ + that exists on the M atom in the species M-X (the metal halides have large electric dipole moments in the range ~ 5 or $(6 D)^{35}$ polarizes electron density preferentially from the π orbital (π_{ab}) in the molecular plane. The larger the polarization, the more the π bonding in the ethyne molecule will resemble that of an ethene molecule in which the symmetry plane of its π bonding orbital is perpendicular to the *ab* plane and the C-H bonds lie in the *ab* plane. In the limit of very large polarization of the π_{ab} electrons, when both are completely removed from ethyne to yield the ion $C_2H_2^{2+}$, the distortions would be of the order $\delta A \approx 60^{\circ}$ and $\delta r \approx 134.5 - 120.6 =$ 13.9 pm, where 134.5 and 120.6 pm are the CC distances in the ground state ${}^3\Sigma_g^{\ -}$ of the dication 36 $C_2H_2^{\ 2^+}$ and in ethyne, 27 respectively. tively. Similar distortions have been observed in $C_2H_4 \cdots M-X$ (M = Cu or Ag),^{6,37} in which M-X lies on the C₂ axis of C_2H_4 that is perpendicular to the plane of the C₂H₄ nuclei. Similar arguments can be used to rationalize the distortions in these molecules, but with $C_2 H_4^{2+}$ as the limiting dication.

Acknowledgements

We thank the European Research Council for postdoctoral fellowships awarded to D.P.Z. and S.L.S., and for project funding (CPFTMW-307000). ACL thanks the University of Bristol for a Senior Research Fellowship and Newcastle University for a Faculty of SAgE Visiting Professorship. D.P.Z. thanks Newcastle University for the award of a Faculty of SAgE Research Fellowship. D.P.T. gratefully acknowledges the award by the Royal Society of a University Research Fellowship.

References

1 K. Schroeter, C. A. Schalley, R. Wesendrup, D. Schroder and H. Schwarz, *Organometallics*, 1997, **16**, 986.

- 2 K. Judai, A. S. Worz, S. Abbet, J. M. Antonietti, U. Heiz, A. Del Vitto, L. Giordano and G. Pacchioni, *Phys. Chem. Chem. Phys.*, 2005, 7, 955.
- 3 A. Molnar, A. Sarkany and M. Varga, *J. Mol. Catal. A: Chem.*, 2001, **173**, 185.
- 4 S. L. Stephens, W. Mizukami, D. P. Tew, N. R. Walker and A. C. Legon, *J. Chem. Phys.*, 2012, **137**, 174302.
- 5 S. L. Stephens, D. P. Zaleski, W. Mizukami, D. P. Tew, N. R. Walker and A. C. Legon, *J. Chem. Phys.*, 2014, **140**, 124310.
- 6 S. L. Stephens, D. M. Bittner, V. A. Mikhailov, W. Mizukami,
 D. P. Tew, N. R. Walker and A. C. Legon, *Inorg. Chem.*, 2014,
 53, 10722–10730.
- 7 G. G. Brown, B. C. Dian, K. O. Douglass, S. M. Geyer, S. T. Shipman and B. H. Pate, *Rev. Sci. Instrum.*, 2008, **79**, 053103.
- 8 S. L. Stephens and N. R. Walker, *J. Mol. Spectrosc.*, 2010, **263**, 27–33.
- 9 D. P. Zaleski, S. L. Stephens and N. R. Walker, *Phys. Chem. Chem. Phys.*, 2014, **16**, 25221–25228.
- 10 T. J. Balle and W. H. Flygare, *Rev. Sci. Instrum.*, 1981, 52, 33-45.
- 11 R. C. Batten, G. C. Cole and A. C. Legon, J. Chem. Phys., 2013, 119, 7903–7912.
- 12 C. Hättig, D. P. Tew and A. Köhn, J. Chem. Phys., 2010, 132, 231102.
- 13 C. Hättig, W. Klopper, A. Köhn and D. P. Tew, *Chem. Rev.*, 2011, **112**, 4–74.
- 14 K. Raghavachari, G. W. Trucks, J. A. Pople and M. Head-Gordon, *Chem. Phys. Lett.*, 1989, **157**, 479–483.
- 15 R. A. Kendall, T. H. Dunning and R. J. Harrison, *J. Chem. Phys.*, 1992, **96**, 6796–6806.
- 16 K. A. Peterson and C. Puzzarini, *Theor. Chem. Acc.*, 2005, 114, 283–296.
- 17 M. Dolg, U. Wedig, H. Stoll and H. J. Preuss, *Chem. Phys.*, 1987, 86, 866–872.
- 18 H.-J. Werner, P.J. Knowles, R. Lindh, F.R. Manby and M. Schütz, MOLPRO, version 2009.1, a package of ab initio programs; 2009, see http://www.molpro.net.
- 19 S. F. Boys and F. Bernardi, Mol. Phys., 1970, 19, 553-566.
- 20 M. Heckert, M. Kállay and J. Gauss, *Mol. Phys.*, 2005, 103, 2109.
- 21 M. Kállay, MRCC, a generalized CC/CI program, see http:// www.mrcc.hu.
- 22 T. Okabayashi, E. Yamazaki, T. Honda and M. Tanimoto, J. Mol. Spectrosc., 2001, 209, 66–70.
- 23 PGOPHER, a program for simulating rotational structure, designed by C.M. Western, University of Bristol, version 6.0.202, 2010, available at http://pgopher.chm.bris.ac.uk.
- 24 J. K. G. Watson, J. Chem. Phys., 1968, 46, 1935.
- 25 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin,

V. N. Staroverov, T. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, O. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski and D. J. Fox, *Gaussian 09, Revision B.01*, Gaussian, Inc., Wallingford CT, 2010.

- 26 See ESI† for assignments of all observed transitions and the fits used to determine the spectroscopic constants of six isotopologues of $C_2H_2\cdots Cu$ -F.
- 27 M. Herman, A. Campargue, M. I. El Idrissi and J. Van der Auwera, J. Phys. Chem. Ref. Data, 2003, 32, 921.

- 28 J. Lievin, J. Demaison, M. Herman, A. Fayt and C. Puzzarini, *J. Chem. Phys.*, 2011, **134**, 064119.
- 29 W. Gordy and R. L. Cook, *Microwave Molecular Spectra*, Wiley, New York, 1984, Appendix E.
- 30 A. Chutjian, J. Mol. Spectrosc., 1964, 14, 361.
- 31 J. Kraitchman, Am. J. Phys., 1953, 21, 17.
- 32 C. C. Costain, J. Chem. Phys., 1958, 29, 864.
- 33 C. C. Costain, Trans. Am. Crystallogr. Assoc., 1966, 2, 157.
- 34 Z. Kisiel, J. Mol. Spectrosc., 2003, 218, 58-67.
- 35 NIST Diatomic Spectra Database, www.nist.gov/pml/data/msddi/index.cfm, compiled by F.J. Lovas, E. Tiemann, J.S. Coursey, S.A. Kotochigova, J. Chang, K. Olsen and R.A. Dragoset.
- 36 R. Thissen, J. Delwiche, J. H. Robbe, D. Duflot, J. P. Flament and J. H. D. Eland, *J. Chem. Phys.*, 1993, **99**, 6590.
- 37 S. L. Stephens, D. P. Tew, V. A. Mikhailov, N. R. Walker and A. C. Legon, *J. Chem. Phys.*, 2011, 135, 024315.