



Cite this: *Phys. Chem. Chem. Phys.*,
2015, 17, 20425

DOI: 10.1039/c5cp90128f

www.rsc.org/pccp

Correction: A novel benzoxazine/cyanate ester blend with sea-island phase structures

Xiaodan Li,^{*ab} Xiaoyong Luo^b and Yi Gu^{*b}

Correction for 'A novel benzoxazine/cyanate ester blend with sea-island phase structures' by Xiaodan Li *et al.*, *Phys. Chem. Chem. Phys.*, 2015, DOI: 10.1039/c5cp02426a.

The scale bar shown in the SEM pictures of Fig. 1 is incorrect and should read 10 μm instead of 10 m. The amended version of Fig. 1 is shown below:




Fig. 1 SEM phase morphology of C-BOZ/BADCy blends: the uncured, cured at 120 °C for 10 h and etched in acetone for 0.5 h of the 5/5 blend (a₁, a₂ and a₃), the 7/3 blend (b₁, b₂ and b₃), the 8/2 blend (c₁, c₂ and c₃) and the 9/1 blend (d₁, d₂ and d₃).

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Key Laboratory of Catalysis Science and Technology of Chongqing Education Commission, College of Environment and Biological Engineering, Chongqing Technology and Business University, Chongqing, 400067, P. R. China

^b State Key Laboratory of Polymer Materials Engineering, College of Polymer Sciences and Engineering, Sichuan University, Chengdu 610065, Sichuan, P. R. China.
E-mail: 12345ruby@163.com, guyi@scu.edu.cn

