Cite this: *Phys. Chem. Chem. Phys.*,
2015, **17**, 32574

Correction: Size and shape dependent photoluminescence and excited state decay rates of diamondoids

Robert Richter,^{*a} David Wolter,^a Tobias Zimmermann,^a Lasse Landt,^a
Andre Knecht,^a Christoph Heidrich,^a Andrea Merli,^a Otto Dopfer,^a Philipp Reiß,^b
Arno Ehresmann,^b Jens Petersen,^c Jeremy E. Dahl,^d Robert M. K. Carlson,^d
Christoph Bostedt,^e Thomas Möller,^a Roland Mitric^c and Torbjörn Rander^{*a}

DOI: 10.1039/c5cp90213d

www.rsc.org/pccp

Correction for 'Size and shape dependent photoluminescence and excited state decay rates of diamondoids' by Robert Richter *et al.*, *Phys. Chem. Chem. Phys.*, 2014, **16**, 3070–3076.

The authors would like to make the following correction to their article:

In Fig. 5, the second emission spectrum for the bottom (dia (D_{3d})) is shifted to higher energies and should be as seen in Fig. 1 below.

It has to be noted that by now, spectra with higher resolution are available. These can be found in ref. 1.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

References

- 1 R. Richter, M. I. S. Röhr, T. Zimmermann, J. Petersen, C. Heidrich, R. Rahner, T. Möller, J. E. Dahl, R. M. K. Carlson, R. Mitric, T. Rander and A. Merli, *Phys. Chem. Chem. Phys.*, 2015, **17**, 4739.

^a Institut für Optik und Atomare Physik, Technische Universität Berlin, Hardenbergstr. 36, 10623 Berlin, Germany. E-mail: robert.richter@tu-berlin.de

^b Institute of Physics and Center for Interdisciplinary Nanostructure Science and Technology (CINSaT), University of Kassel, Heinrich-Plett-Str. 40, 34132 Kassel, Germany

^c Universität Würzburg, Institut für Physikalische und Theoretische Chemie, Am Hubland, 97074 Würzburg, Germany

^d Stanford Institute for Materials and Energy Sciences, Stanford University, Stanford, California 94305, USA

^e SLAC National Accelerator Laboratory, Menlo Park, California 94025, USA

Fig. 1 Measured photoluminescence spectra (red) and absorption spectra (black) of selected isomers of the five smallest members of the diamondoid series. The excitation energy used to record the PL is marked with an asterisk and E_{gap} is marked with a blue arrow. PL-spectra of adamantane ($\Delta E = 100$ meV), diamantane ($\Delta E = 100$ meV), other spectra ($\Delta E = 25$ meV, the spectra have been smoothed for presentation purposes).