## Dalton Transactions



**PAPER** 

View Article Online
View Journal | View Issue



**Cite this:** *Dalton Trans.*, 2015, **44**, 3664

Received 4th September 2014, Accepted 4th December 2014 DOI: 10.1039/c4dt02713b

www.rsc.org/dalton

# Mono- and dinuclear metal complexes containing the 1,5,9-triazacyclododecane ([12]ane $N_3$ ) unit and their interaction with DNA†

Alfredo Medina-Molner, Melanie Rohner, Devaraj Pandiarajan and Bernhard Spingler\*

The ability of mononucleating and dinucleating macrocylic polyamines and their novel nickel, copper and zinc complexes to induce the left-handed form of poly d(GC) was evaluated. The influence of the nuclearity, the presence or absence of metals ions, the linker length in the case of dinucleating ligands and the metal ion was determined. Almost all dinuclear metal complexes efficiently induced Z-DNA, the zinc ones being the least and the copper ones the most efficient ones. Additionally, the X-ray structures of three dinuclear metal complexes and one partially protonated ligand could be determined.

#### Introduction

Left-handed Z-DNA is a high-energy form of DNA, it can only be formed by alternating purine-pyrimindine sequences and when additional factors like supercoiling or multiple charged cations are present.1 In plasmids, formation of Z-DNA depends upon torsional stress that is induced by negative supercoiling.<sup>2-4</sup> The twin-supercoiled domain model suggests that the movement of an RNA polymerase along the DNA double helix induces positive supercoil waves ahead of, and negative supercoil waves behind the transcription ensemble.<sup>4</sup> As a consequence, Z-DNA is formed after the passing of the polymerase as the kinetics of the B- to Z-DNA transition is relatively rapid compared to the cellular processes such as transcription. Z-DNA was shown to play a functional role in gene regulation,<sup>5</sup> positioning of the nucleosome<sup>6</sup> and chromatin remodelling.<sup>7</sup> Several research groups have studied the possible correlation between chromosomal breakpoints in human tumours with potential Z-DNA forming sequences.8 Z-DNA-GC repeats were shown to cause large-scale genome deletions in mammalian cells<sup>9</sup> but only small changes in E. Coli. 10 These variations were explained by different DNA repair mechanisms. 11 Vásquez observed that the deletions or rearrange-

University of Zürich, Winterthurerstr. 190, CH 8057 Zürich, Switzerland.

E-mail: spingler@chem.uzh.ch; Fax: +41 44 635 68 03; Tel: +41 44 635 46 56
†Electronic supplementary information (ESI) available: Synthesis of 1,4-butylenebistriflate, ORTEP representation of L4-4TfOH, crystallographic table and various CD spectra/titrations of poly d(GC) with of poly d(GC) with LEt, L3, L4, 1, 2, 3, 4, 5, 6, 7, 11, 12, EDTA, 6 and EDTA, 9 and EDTA. CCDC 1021588–1021591. For ESI see DOI: 10.1039/c4dt02713b

ments generated in mammalian cells are mainly replication independent and are likely to be initiated by repair processing cleavages surrounding the Z-DNA-forming sequence. It was proposed that these types of genomic alterations are related to the chromosome breaks and gene translocations that map near Z-DNA-forming sequences in human leukaemia and lymphoma. Recently, the group of Dröge employed the Z-DNA binding protein ADAR1 as a probe together with a chromatin affinity precipitation. This procedure revealed almost 200 genomic Z-DNA hotspots in human A549 tumor cells. A forth of the hotspots were located in centromeres, but only two in promoter regions.

In contrast with B-DNA, Z-DNA is highly immunogenic and some antibodies can recognise this conformation and these antibodies are not species-specific.<sup>14</sup> For instance, patients with *lupus erythematosis* produce – among other nuclear components – antibodies with high specificity for Z-DNA.<sup>3</sup> Lipps carried out an experiment where Z-DNA antibodies were used to stain ciliated protozoa that have both macronucleus and micronucleus.<sup>15</sup> The micronucleus is only needed for genetic reproduction, whereas the transcription occurs exclusively in the macronucleus. The Z-DNA antibody was only staining the macronucleus, not the micronucleus. This experiment indirectly connected Z-DNA and transcription.

Antibodies were also used in metabolically active mammalian nuclei to detect Z-DNA. The experiments showed a relationship between the presence of Z-DNA and the negative torsional strain of DNA. As for the findings of Lipps' group, the amount of Z-DNA increased considerably as transcription occurred, but was unaffected during DNA replication. <sup>16</sup> De la

Dalton Transactions Paper

Torre used immune-detection of Z-DNA as a marker for active transcription.<sup>14</sup> She proposed this mechanism of detecting transcription as a cheaper and quicker method than the alternative *in situ* assay for active RNA polymerases.

The goal of our project was to design and study small molecules that are able to induce Z-DNA by strong and selective binding to this structural motive. The presence of Z-DNA has been linked to gene regulation in general and transcription in particular. Therefore, Z-DNA is an attractive new target for cancer therapy. Since Z-DNA is distinguished from the other conformations of DNA by its own very characteristic geometry, the recognition must be mainly based on geometrical features and not so much on the sequence of the nucleobases. In Z-DNA, the bases are pointing away from the helix axis. As a consequence, the bases are not as protected from the environment as in B-DNA; for example nitrogen N7 of the purines is highly exposed to the solvent. 18,19


Previously, we have studied the ability of mononuclear complexes to induce Z-DNA. 19,20 Furthermore, we have chosen to explore mono- and dinuclear metal complexes based upon 1,5,9-triazacyclododecane ([12]aneN<sub>3</sub>), which are expected to coordinate to N7 of neighbouring guanosine nucleobases.<sup>21</sup> Linking two [12]aneN3 rings creates dinucleating ligands that are unable to form sandwich complexes in contrast to their smaller [9]aneN3 analogues.22 Lu and co-worker have systematically substituted hydrogen atoms of one or two N-H groups by methyl group(s) of dinucleating [12]aneN<sub>3</sub> ligands.<sup>23</sup> An increasing number of methyl groups dramatically decreased the ability of the corresponding dizinc complexes to hydrolyse RNA model compounds. This observation was explained by steric effects, though it remains unclear, whether the missing hydrogen donor ability of these methyl substituted complexes had also an influence.

A preliminary description of our results with one dinucleating [12]ane $N_3$  ligand L3 and its metal complexes has recently appeared.<sup>24</sup> Here, we discuss the chosen system in detail with an emphasis on the influence of the linker length between the two [12]ane $N_3$  macrocycles.

#### Results and discussion

In this publication, we study the interaction of poly d(GC) with the mononucleating ligands L, LEt and the dinucleating ligands L2, L3, L4 as well as their Ni, Zn and Cu complexes (Scheme 1).

Originally, we started working on the bis-[12]aneN<sub>3</sub> ligand L3 that has its two [12]aneN<sub>3</sub> rings joined together by a propylene linker.<sup>24</sup> When comparing the two X-ray structures of the copper complexes L3Cu<sub>2</sub>(NO<sub>3</sub>)<sub>2</sub>(TfO)<sub>2</sub> (Fig. 1) and 9,<sup>24</sup> they revealed the high flexibility of the chosen ligand system. Both complexes were synthesized starting from ligand L3, in the latter case with copper(II) chloride to give the µ-hydroxo,µ-chlorido bridged complex 9 with a copper-copper distance of 3.2350(9) Å. In the former case, reaction of L3 with two equivalents of copper(II) nitrate gave the dinuclear complex



I	$\mathbf{L}, \mathbf{R} = \mathbf{H}$	$\mathbf{LEt}$ , $\mathbf{R} = \mathbf{Et}$	
II	<b>L2</b> , $n = 1$	<b>L3</b> , $n = 2$	<b>L4</b> , $n = 3$
III, R = H	$1, MX_2 = Ni(NO_3)_2$	2, $MX_2 = Cu(NO_3)_2$	$3, MX_2 =$
			$Zn(NO_3)_2$
III, R = Et	$4$ , $MX_2 = Ni(OAc)_2$	5, $MX_2 = Cu(OAc)_2$	
IV, $n = 1$	$6, MX_2 = Ni(OAc)_2$	7, $MX_2 = Cu(OAc)_2$	
IV, $n = 2$	$8, MX_2 =$	$9, M_2X_4 =$	$10, M_2X_4 =$
	Ni(OAc)(OTf)	$Cu_2(OH)(Cl)(OTf)_2$	$Zn_2(OH)(TfO)_3$
IV, $n = 3$	11, $MX_2 =$	12, $MX_2 = Cu(OTf)_2$	
	Ni(OTf) <sub>2</sub>		

Scheme 1 Studied compounds.




Fig. 1 ORTEP representation of  $L3Cu_2(NO_3)_2(TfO)_2 \cdot (CH_3OH)_{0.7}(H_2O)_{0.3}$  at 50% probability. Non-acidic hydrogen atoms and solvent molecules were omitted for clarity.

 $L3Cu_2(NO_3)_2(TfO)_2$  that does not show any copper-copper interaction (Cu(1)-Cu(2) distance: 8.5803(10) Å, Fig. 1). These findings were the reason why the linker length in the dinucleating ligand was varied in order to study the effect of increasing and reducing the flexibility of the system on the B- to Z-DNA transition.

In order to synthesize the known 1,4-bis(triazacyclododecyl)-butane (**L4**), we did not follow the route described in the literature<sup>25</sup> but chose an alternative way that we already had employed for the synthesis of 1,4-bis(triazacyclododecyl)-propane.<sup>24</sup> Our method led to an improvement of the overall yield of 1,4-bis(triazacyclododecyl)butane **L4** from 40%<sup>25</sup> to 74%. The synthesis of the corresponding precursor 1,4-butylenebistriflate could also be improved from 63%<sup>26</sup> to 98%. 1,4-butylenebistriflate was added to two equivalents of 1,5,9-triazatricyclo[7.3.1.0]tridecane at room temperature to yield the desired product **13** in 86% yield. Compound **13** was then

Scheme 2 Synthesis of 1,4-bis(triazacyclododecyl)butane (L4).

refluxed in triflic acid for 18 hours in order to obtain the hexaprotonated ligand L4 in 86% yield (Scheme 2). Deprotonation by sodium hydroxide and finally ion exchange column quantitatively yielded the free ligand L4.

Crystals of 1,4-bis(triazacyclododecyl)butane·4TfOH could be obtained from a partially neutralised ethanolic solution of the hexaprotonated ligand **L4**, simply by concentrating the sample and storing it at 4 °C (Fig. S1†). In the crystal structure of **L4·4TfOH**, the asymmetric unit consists of half of a molecule – the other half being generated by a centre of inversion. All four NH protons of one [12]aneN<sub>3</sub> ring were localized and, not surprisingly, all are involved in hydrogen bonding: The proton of N1 forms an internal bridge across the ring to N2 (2.693(2) Å), while the remaining 3 protons all bind to triflate anions with N–O distances between 2.732(3) Å (N3–O31\_x+0.5, y+0.5, z-0.5) to 2.934(3) Å (N2–O30). The synthesis of the ligand 1,2-bis(triazacyclododecyl)ethane (**L2**) has already been described by us before. <sup>27</sup>

The synthesis and isolation of the nickel (1), copper (2) and zinc (3) complexes has been achieved from ligand L in methanol and the corresponding nitrate salts. The synthesis of 1 with hydroxide and perchlorate as the anions has been reported together with the stability constant of  $\mathbf{L}\cdot\mathbf{Cu}(\Pi)$ . A  $\mu$ -chloro- $\mu$ -hydroxo dimer of  $\mathbf{Cu}(\Pi)$ [12]aneN $_3$  with perchlorates as remaining anions has been characterised. The *in situ* generation of 3, but not its characterisation, was reported.

Complex  $L4Ni_2(AcO)_4$  was obtained when ligand L4 was mixed with two equivalents of  $Ni(CH_3CO_2)_2$ . Since crystals could not be grown of the reaction product, an excess of NaB- $(C_6H_5)_4$  was added to a solution of complex  $L4Ni_2(AcO)_4$  in water. The resulting precipitate was filtered off and dissolved in acetonitrile. Crystals could finally be grown by vapour diffusion of tetrahydropyrane (THP) to the acetonitrile solu-




Fig. 2 ORTEP representation of [L4Ni<sub>2</sub>(AcO)<sub>2</sub>(BPh<sub>4</sub>)<sub>2</sub>(NCCH<sub>3</sub>)<sub>2</sub>]-THP at 50% probability. Non-acidic hydrogen atoms, non-coordinating anions and non-coordinating solvent molecules were omitted for clarity.




Fig. 3 ORTEP representation of  $6.2H_2O$  at 50% probability. Non-acidic hydrogen atoms were omitted for clarity.

tion.<sup>31</sup> The nickel centre of  $L4Ni_2(OAc)_2(B(C_6H_5)_4)_2(NCCH_3)_2$  presents a distorted octahedral coordination centre (Fig. 2). The triazamacrocycle binds to the nickel cation in a facial tridentate fashion; the remaining coordination sites are occupied by a bidentate acetate and an acetonitrile. The intramolecular nickel–nickel distance is 9.6723(7) Å. The asymmetric unit is composed of half a molecule, the other half of the complex is generated by a centre of inversion. A disordered THP molecule was found in the crystal lattice as well.

Ligand L2 was reacted with two equivalents of nickel acetate in a mixture of methanol–acetonitrile to yield a blue solution. The resulting nickel complex (6) was crystallised by vapour diffusion of pentane into the corresponding complex solution in dichloromethane. The nickel centre presents a slightly distorted octahedral coordination. The nickel–nickel distance is with 6.542(1) Å 3.13 Å shorter than in the butylene bridged case. The water molecule coordinated to the nickel centre is in hydrogen bonding with the non-coordinating oxygen atoms of both acetate groups (Fig. 3).

Ligand L3 was used to synthesise dinuclear nickel (8), copper (9) and zinc (10) complexes. All three metal complexes could induce Z-DNA (Fig. 4). These results showed for the very first time an opposite behaviour between mononuclear and dinuclear metal complexes. Thereby, our original hypothesis






Fig. 4 CD spectra of poly d(GC) and with added L3, 8, 9 or 10. The spectra are shown after the possible transition has been completed.

Table 1 Studied ligands and metal complexes and their effect upon poly d(GC): Z: induction of Z-DNA, denat.: denaturation. condens.: condensation; n.d.: not determined. Numbers after the compound number stand for the midpoint of the observed transition expressed as metal equivalents versus DNA phosphates (if observed)

Ligand	No metal	Ni	Cu
L	L·3TfOH, n.d.	1, none	2, none
LEt	LEt·3TfOH, none <sup>24</sup>	4, none <sup>24</sup>	5, denat. 0.53 <sup>24</sup>
L2	L2·6TfOH, n.d.	6, Z 0.20	7, none
L3	L3·6TfOH, none <sup>24</sup>	8, Z 0.21 <sup>24</sup>	9, Z 0.16 <sup>24</sup>
L4	L4·6TfOH, condens.	11, Z 0.25	12, Z 0.17

was confirmed that dinuclear metal complexes with the appropriate distance between the two metal centres would be more efficient Z-DNA inducers than mononuclear ones.<sup>32</sup> However, our experimental studies showed a much more dramatic result, since the mononuclear complexes 1, 2 and 3 (Fig. S2, pink; Fig. S3, blue; Fig. S3,† pink) did not induce Z-DNA at all, or even induced denaturation (complex 5), whereas their equivalent dinuclear complexes are one of the best Z-DNA inducers described so far (Table 1).

The nickel complex 8 induced the midpoint of the B- to Z-DNA transition at 0.21 equivalents of metal (0.11 equivalents of metal complex). The zinc complex 10 was the worst Z-DNA inducer among the three successful complexes having a midpoint at 0.30 equivalents of zinc (0.15 equivalents of complex, Fig. 5). The best Z-DNA inducer was the copper complex 9, even though it contained chloride which is retarding the B- to Z-DNA transition.<sup>33</sup> The midpoint of this transition happened at 0.16 equivalents of copper (0.08 equivalents of complex, Fig. 5). As the zinc complex 10 was the worst Z-DNA inducer (0.30 equivalent metals needed versus DNA phosphate) among the metal series of nickel, copper and zinc, we decided not to further study the other members of the homologous series of the zinc complexes.




Fig. 5 CD titration at 255 nm of poly d(GC) with added 1, 2, 3, 8, 9, or 10. The metal content of the complexes and not the equivalents of complexes is given in the x-axis (in order to compensate for the fact that dinuclear complexes have twice as much metal ions as the mononuclear ones).

The linker between the two units in ligand L3 was first increased from three to four carbon atoms to give ligand L4, which is expected to have a higher flexibility and to generate a longer distance between the two metal centres. Nickel (11) and copper (12) complexes induced Z-DNA at a slightly higher concentration than their equivalent complexes with ligand L3. Thus increasing the flexibility of the system had only a slightly negative if any effect on the ability of the complexes to induce Z-DNA (Fig. S4 and S5†).

Since increasing the flexibility of the system did not help to improve the efficiency of Z-DNA induction, a more rigid ligand was synthesised by reducing the length of the linker to two carbon atoms in ligand L2. Nickel (6) and copper (7) complexes were synthesised and tested for their ability to promote the B- to Z-DNA transition. The nickel complex (6) was slightly more efficient in inducing Z-DNA (Fig. S4†) than the other dinickel complexes. However, the copper complex (7) was not able to induce Z-DNA (Fig. S5†), demonstrating a dramatic difference when the flexibility of the ligand was reduced.

The mononucleating ligand LEt was synthesised in order to generate a mononuclear system that resembles a halved dinuclear complex, in which the ethyl substituent mimics the steric and electronic influences of the linker in the dinuclear systems. A similar behaviour as for nickel complex 1 was observed when the nickel complex (4, Fig. S6,† blue) with ligand LEt was tested in the B- to Z-DNA transition. However a different result was obtained with the copper complex 5 that contained ligand LEt. Denaturation of the DNA was induced after the addition of 0.3 equivalents of the copper complex 5 (Fig. S6,† pink). The denaturation is more easily seen by either looking at the whole CD or UV/vis spectra at different equivalents of the copper complex (Fig. S7†). In summary it can be said that neither the nickel 4 nor the copper 5 complexes induced Z-DNA.

**Dalton Transactions Paper** 

As a control, the mononuclear ligand LEt-3TfOH and dinuclear butylene bridged ligand L4.6TfOH were also tested for their ability to induce Z-DNA (Fig. S8†). Ligand LEt-3TfOH did not have any effect on DNA, however the dinucleating ligand L4.6TfOH induced condensation of DNA.34 This condensation started at around 0.5 equivalents of ligand versus DNA phosphates and lead to a strong increase of the absorption at 255 nm. This aggregation of DNA did not have an intermediate Z-DNA transition as observed in Zacharias' work, instead of that the condensation occurred directly from B-DNA as in Chaires studies<sup>34,35</sup> (Fig. S9†). Zacharias suggested that the hydrophobic part of the acetate group could be decisive for the condensation.<sup>34</sup> In our system, 6 triflates per ligand are present; the CF<sub>3</sub> group of the triflates could be the hydrophobic tale needed for the condensation. However, LEt-3TfOH and L3-6TfOH with the same ratio of nitrogen atoms versus triflates did not induce condensation of DNA.

EDTA was used to study the reversibility of the transition metal induced B- to Z-DNA transformation. 32,36 The action of EDTA alone upon poly d(GC) was studied first. As expected, EDTA did not induce Z-DNA but rather surprisingly, at high concentration caused condensation of DNA (Fig. S10†). Further addition of EDTA lowered the positive CD band at 275 nm. Then, the effect of EDTA upon the complexes 6 (Ni)/9 (Cu) and poly d(GC) were investigated (Fig. S11†). In both cases, the DNA essentially converted back to the B-form after the addition of two equivalents of EDTA, and a full conversion back to B-DNA was found after 3 equivalents of EDTA versus metal ions.

#### Conclusions

This study aimed to determine the factors that favour the induction of Z-DNA by metal complexes. Our original hypothesis was that sterically demanding dinuclear metal complexes with a metal to metal distance of about 5-7 Å would be more efficient inducers of the left-handed form than other dinuclear and mononuclear complexes. This idea was inspired by looking at the geometry of Z-DNA with its solvent exposed atom N7 of guanine and the interstrand N7-N7 distance between two guanine of a GC base pair step. We soon realized that for the 1,5,9-triazadodecane system, the reality was even better than our hypothesis: Only the dinuclear copper and nickel complexes of 1,3-bis-triazadodecyl-propane were inducing the left-handed form of poly d(GC) but not the corresponding N-ethyl-triazadodecane mononuclear metal complexes.<sup>24</sup> In the current publication, we have substantially expanded the set of studied compounds by including dinucleating ligands whose rings are either linked by an ethylene or butylene group. As a first conclusion, we can say that none of the studied ligands were inducing the Z-form. L4·6TfOH was causing condensation of the DNA, but apart from this result no effects were observed. This is interesting because various linear polyamines were found to be powerful Z-DNA

inducers.<sup>37</sup> To the best of our knowledge, no macrocyclic polyamines were ever reported to induce Z-DNA.<sup>38</sup> Summarising the results of the extended set of mononuclear metal complexes, none of them induce Z-DNA. On the other hand, almost all of the dinuclear complexes do induce the lefthanded form with the notable exception of the di-copper complex of 1,2-bis-triazadodecyl-ethane. At the moment, we still do not know, what causes this exception. Comparing the relative ability of the various metal ions, copper seems to be always better than nickel and zinc. This trend essentially follows the Irving-Williams series<sup>39</sup> as discussed by Barone et al. 40 Concerning the influence of the linker length, we can note that the differences between ethylene, propylene and butylene linker are very subtle, with the propylene linker being the most efficient among them. This indifference of the system towards the linker length might be caused by the nature of the ligands that lack an anionic central attachment point for the metal ions<sup>41</sup> and the flexibility of the chosen aliphatic linker.

As an outlook, one could imagine similar complexes having high affinity for these specific forms of DNA that could be functionalised with fluorescence markers or damaging moieties for future applications in a chemical detection of the transcription or in an antitumour therapy.

### Experimental section


#### Instrumentation and materials

All chemicals were purchased from Sigma-Aldrich (Buchs, CH) and used without further purification. The synthesis of compounds LEt, L3, 4, 5, 8, 9<sup>24</sup> and L2<sup>27</sup> has already been described by us previously. The synthesis of 1,4-butylenebistriflate was done slightly differently than reported in the literature<sup>26</sup> (see ESI† for details). All the reactions were performed under a nitrogen atmosphere. The reactions were monitored by HPLC or thin layer chromatography (TLC). TLC was carried out on 0.25 mm Merck silica gel aluminium plates (60 F254) or aluminium oxide pre-coated plastic sheets (alox N/UV254) using UV light or Schlittler's reagent42 as a visualising agent. Column chromatography was performed on silica gel (particle size 0.040-0.063 mm) or aluminium oxide (basic, 0.05-0.15 mm, pH  $9.5 \pm 0.5$ ). Ion exchange chromatography was performed on DOWEX 2 X8 20-50 (Fluka) or AMBERLYST A26-OH, macromolecular ion-exchange resin (ABCR). Elemental analyses were performed on a Leco CHNS-932 elemental analyser. Electrospray ionisation (ESI) mass spectra were recorded on a Merck Hitachi M-8000 spectrometer or a Bruker Daltonics HCT 6000 mass spectrometer. UV/Vis spectra were recorded on a Varian Cary 50 spectrometer and IR spectra on a Perkin Elmer, Spectrum Two spectrometer equipped with a Golden Gate ATR. NMR spectra were acquired on a Varian Mercury 200 MHz or Gemini 300 MHz, and on a Bruker DRX 400 or 500 MHz spectrometer. The chemical shifts are relative to residual solvent protons as reference. Circular dichroism measurements were performed using a Jasco J-810 spectro**Dalton Transactions** Paper

polarimeter equipped with a Jasco PFD-4255 Peltier temperature controller. A 0.1 mM poly d(GC) solution in 1 mM sodium cacodylate buffer (pH = 7) was titrated with various aliquots of 1 mM solutions of the ligands or metal complexes in the same buffer. The samples were warmed to 60 °C for 5 minutes and then cooled down to 25 °C for the CD measurements. The CD spectra were smoothed by adjacent averaging. Crystallographic data were collected at 183(2) K on a Stoe IPDS (Mo K<sub>r1</sub> radiation,  $\lambda = 0.71073$  Å) diffractometer using a graphitemonochromated radiation. Suitable crystals were covered with Paratone N oil, mounted on top of a glass fibre and immediately transferred to the diffractometer. A maximum of eight thousand reflections distributed over the whole limiting sphere were selected by the program SELECT and used for unit cell parameter refinement with the program CELL. 43 Data were collected for Lorentz and polarisation effects as well as for absorption (numerical). Structures were solved with direct methods using SHELXS-9744 or SIR9745 and were refined by full-matrix least-squares methods on  $F^2$  with SHELXL-97. Crystals with low symmetry space groups measured on the one-circle Stoe IPDS diffractometer have an intrinsically low completeness of their data sets. CCDC 1021588-1021591 contain the supplementary crystallographic data for this paper.

Synthesis of compound 13. 1,4-Butylenebistriflate (4.7 g, 13.3 mmol) dissolved in dry DCM (90 ml) was slowly added to a solution of 1,5,9-triazatricyclo[7.3.1.0]tridecane (3.93 g, 21.7 mmol) in dry DCM (25 ml) using Schlenk line techniques. After 18 hours of stirring, abundant precipitate was obtained. The precipitate was filtered off and the solution was dried under reduced pressure, but no further solid was obtained (13, 6.68 g, 86%). Elemental analysis:  $C_{26}H_{46}F_6N_6O_6S_2$ : calc. C (43.57%), H (6.47%), N (11.72%); exp. C (43.54%), H (6.11%), N (11.82%). MS (ESI): m/z (%) = 567 [M - F<sub>3</sub>CSO<sub>3</sub>]<sup>+</sup> (100%). <sup>1</sup>H-NMR (500 MHz, CD<sub>3</sub>OD):  $\delta$  = 1.31 (broad d, J = 15.1 Hz, 4 H,  $\mathbf{H}(d)$ ), 1.42 (t, J = 3.7 Hz,  $\mathbf{H}(a)$ ), 2.11–2.19 (m, 4 H, H(d') + H(g), 2.27-2.30 (m, 2 H, H(g')), 2.62 (t, J = 13.1 Hz, 4 H, H(e)), 2.77 (broad s, 4 H, H(b)), 3.08 (broad d, J = 13.2 Hz, 4 H,  $\mathbf{H}(e')$ ), 3.28 (broad s, 4 H,  $\mathbf{H}(e)$ ), 3.43 (broad t, J = 10.5 Hz, 4 H, H(c'), 3.49 (d, J = 14.2 Hz, H(f)), 3.79 (t, J = 13.1 Hz, 4 H, H(f')) and 8.22 (s, 2 H, H(h)). <sup>13</sup>C-NMR (125 MHz, CD<sub>3</sub>OD):  $\delta =$ 20.72 (C(g)), 22.15 (C(a)), 24.21 (C(d)), 43.65 (C(c)), 51.81 (C(b)), 55.94 (C(e)), 56.52 (C(f)), 123.43  $(q, J = 318.6 \text{ Hz}, CF_3)$ and 158.22 (C(h)). See Scheme 3 for numbering.

Synthesis of 1,4-bis(1,5,9-triazacyclododecyl)butane⋅6(trifluoromethanesulfonic acid) (L4·6 TfOH). Compound 13 (6.47 g, 9.05 mmol) was dissolved in 200 ml CF<sub>3</sub>SO<sub>3</sub>H (0.7 M) and refluxed for 18 hours. The resulting colourless solution was concentrated under reduced pressure observing crystal formation. When ethanol was added to the mixture, the hexatriflate salt of 1,4-bis(1,5,9-triazacyclododecyl)-butane (L4) precipitated as a white powder. Ligand (L4·6 TfOH) was filtered off and rinsed several times with a mixture of ethanoldiethylether (1:6) (10.11 g, 86%). Elemental analysis:  $C_{28}H_{54}F_{18}N_6O_{18}S_6$ : calc. C (25.93%), H (4.20%), N (6.48%); exp. C (25.81%), H (4.23%), N (6.43%). MS (ESI): m/z (%) = 547



Atom numbering for 13 (left) and L4 (right).

 $[M + TfOH + H]^{+}$  (100), 697  $[M + 2(TfOH) + H]^{+}$  (7), 847  $[M + 3(TfOH) + H]^{+}$  (4). IR (Golden Gate ATR):  $\tilde{\nu} = 3045 (NH_{2}^{+}, w)$ , 2830 (CH<sub>2</sub>, w), 1212 (C-F, s), 1165 (SO<sub>3</sub>, s), 1021 (S=O, s) cm<sup>-1</sup>. <sup>1</sup>H-NMR (400 MHz, D<sub>2</sub>O):  $\delta = 1.93$  (broad s, 4 H, H(a)), 2.30-2.33 (m, 12 H, H(d) + H(g)), 3.37 (broad s, 4 H, H(b)), 3.41-3.46 (m, 16 H, H(e) + H(f)) and 3.53 ppm (broad s, 8 H, **H**(c)). <sup>13</sup>C-NMR (100 MHz, D<sub>2</sub>O):  $\delta$  = 17.88 (C(d)), 20.96 (C(g)), 21.47 (Ca(a)), 41.39 (C(f)), 42.46 (C(e)), 47.57 (C(c)), 54.60 (C(b)) and 120.21 (q, J = 317.0 Hz, CF<sub>3</sub>). Crystals could be obtained from a partially neutralised ethanolic solution of the originally hexaprotonated ligand, simply by concentrating the sample and storing at 4 °C.

Synthesis of 1,4-bis(1,5,9-triazacyclododecyl)butane (L4). Ligand (L4·6 TfOH) (2.02 g, 1.6 mmol) was dissolved in 15 ml of NaOH (1 M) and NaCl (3.5 M). The resulting solution was extracted with dichloromethane. The combined organic phases were concentrated under reduced pressure. The resulting oil was then dissolved in water and run through an activated ion exchange column of the type amberlist A-26, macromolecular ion-exchange resin.47 All the fractions containing the product were concentrated together, yielding a colourless oil of L4 (0.688 g, quantitative). Elemental analysis:  $C_{22}H_{48}N_6$ : calc. C (66.62%), H (12.20%), N (21.19%); exp. C (66.61%), H (11.95%), N (21.00%).

Synthesis of the mono-nickel complex of ligand L (1). Ni(NO<sub>3</sub>)<sub>2</sub> (96.9 mg, 0.34 mmol) was dissolved in methanol and added to a solution of ligand L (57.0 mg, 0.33 mmol) in the same solvent. The resulting green solution was concentrated and dried to yield the green complex 1 in quantitative yield. Elemental analysis: C<sub>9</sub>H<sub>21</sub>N<sub>5</sub>NiO<sub>6</sub>: calc. C (30.54%), H (5.98%), N (19.78%); exp. C (30.38%), H (5.74%), N (19.62%). MS (ES): m/z (%) = 291 [M - NO<sub>3</sub>]<sup>+</sup>, (100), 644  $[2M - NO_3]^+$  (50). IR (Golden Gate ATR):  $\tilde{\nu} = 3197$  (NH, w), 2935  $(CH_2, w)$ , 1487  $(CH_2, m)$ , 1385  $(NO_3^-, s)$  cm<sup>-1</sup>.

Synthesis of the mono-copper complex of ligand L (2). Cu(NO<sub>3</sub>)<sub>2</sub> (208 mg, 0.861 mmol) was dissolved in methanol (8 ml) and added to a solution of ligand L (145 mg,

**Paper** 

0.848 mmol) also in methanol (6 ml). A precipitate was filtered off and the resulting dark green solution was concentrated under reduced pressure to give 212 mg (61% yield). Blue crystals of complex 2 were obtained after a week in the fridge. Elemental analysis:  $C_9H_{21}CuN_5O_6$ : calc. C (30.12%), H (5.86%), N (19.52%); exp. C (30.45%), H (5.78%), N (19.67%). MS (ESI): m/z (%) = 296 [M - NO<sub>3</sub>]<sup>+</sup> (100), 654 [2M - NO<sub>3</sub>]<sup>+</sup> (40). IR (Golden Gate ATR):  $\tilde{\nu}$  = 3233 (NH, w), 1421 (NO<sub>3</sub><sup>-</sup>, m), 1340

 $(NO_3^-, s)$ , 1007 (Cu-N, w) cm<sup>-1</sup>.

Synthesis of the mono-zinc complex of ligand L (3). Zn(NO<sub>3</sub>)<sub>2</sub> (87.5 mg, 0.29 mmol) was dissolved in methanol and added to a solution of ligand L (50.5 mg, 0.30 mmol) also in methanol. An abundant white precipitate appeared after a few minutes of stirring. The precipitate was filtered off and rinsed several times with cold methanol to give complex 3 in 56% yield. Elemental analysis: C<sub>9</sub>H<sub>21</sub>N<sub>5</sub>O<sub>6</sub>Zn: calc. C (29.97%), H (5.87%), N (19.42%); exp. C (30.18%), H (5.61%), N (19.42%). MS (ESI): m/z (%) = 172  $[M - Zn(NO_3)_2 + H]^+$  (100), 406  $[2M - INO_3]_2$  $Zn - 4NO_3^{\dagger}$  (50), 531  $[2M - Zn(NO_3)_2 + H]^{\dagger}$  (10). <sup>1</sup>H-NMR (200 MHz,  $D_2O$ ):  $\delta = 1.52$  (ttd, J = 2, 9, 16 Hz, 3H, CCHC), 1.92 (ttd, J = 2, 9, 16 Hz, 3H, CCH'C), 2.76 (ddd, J = 2, 9, 13 Hz, 6H, CHN) and 3.08 ppm (ddd, J = 2, 9, 13 Hz, 6H, CH'N). The <sup>1</sup>H-NMR of the SCN<sup>-</sup> salt was reported by Kimura. <sup>48</sup> IR (Golden Gate ATR):  $\tilde{\nu} = 3192$  (NH, m), 2934 (CH<sub>2</sub>, w), 1490 (CH<sub>2</sub>, m), 1457 (NO<sub>3</sub><sup>-</sup>, m), 1324 (NO<sub>3</sub><sup>-</sup>, s), 1009 (Zn-N, m)  $cm^{-1}$ .

Synthesis of the di-nickel complex of ligand L2 (6). Ligand L2 (50 mg, 0.14 mmol) in MeOH was slowly added to a stirred solution of nickel acetate (69 mg, 0.28 mmol) in MeOH, obtaining a blue solution. The solution was dried under reduced pressure. The resulting oil was dissolved in dichloromethane. Complex 6 crystallised by slow diffusion of pentane into the previous solution to give 76.2 mg (68% yield). Elemental analysis:  $C_{20}H_{44}N_6Ni_2(CH_3CO_2)_4(CH_2Cl_2)(H_2O)$ : calc. C (42.21%), H (7.33%), N (10.18%); exp. C (42.22%), H (7.43%), N (10.07%). MS (ESI): m/z (%) = 661 [M – OAc]<sup>+</sup> (100), 302 [M – 2(OAc)]<sup>2+</sup> (90). IR (Golden Gate ATR):  $\tilde{\nu}$  = 2912 (CH<sub>2</sub>, w), 1568 (AcO<sup>-</sup>, s), 1407 (AcO<sup>-</sup>, s), 877 (Ni–N, m) cm<sup>-1</sup>.

Synthesis of the di-copper complex of ligand L2 (7). To a stirred mixture of copper acetate (54 mg, 0.27 mmol) in MeOH (4 ml), ligand L2 (50 mg, 0.14 mmol) in MeOH (4 ml) was slowly added, resulting in a dark blue solution. The solution was dried under reduced pressure yielding a blue oil (7). Elemental analysis:  $C_{20}H_{44}Cu_2N_6(CH_3CO_2)_4(CH_2Cl_2)_{1.5}(H_2O)$ : calc.: C (40.40%), H (7.01%), N (9.58%); exp.: C (40.52%), H (6.81%), N (9.56%). MS (ESI): m/z (%) = 429 [M - Cu - 4(OAc) - 2H]<sup>+</sup> (50), 491 [M - Cu - 3(OAc) + H]<sup>+</sup> (100), 509 [M - Cu - 3(OAc) +  $H_2O + H$ ]<sup>+</sup> (30), 551 [M - Cu - 2(OAc) + 2H]<sup>+</sup> (43). IR (Golden Gate ATR):  $\tilde{\nu}$  = 2962 (CH<sub>2</sub>, w), 1557 (AcO<sup>-</sup>, m), 1393 (AcO<sup>-</sup>, m), 1015 (Cu-N, m) cm<sup>-1</sup>.

Synthesis of the di-zinc complex of ligand L3 (10). Ligand L3 (150.7 mg, 0.12 mmol) and KOH (39.5 mg, 0.70 mmol) was refluxed in ethanol for 10 minutes. Once it had cooled down, the resulting solution was added drop-wise to a solution of  $Zn(CF_3SO_3)_2$  (85.7 mg, 0.24 mmol) also in ethanol. The result-

ing solution was concentrated under reduced pressure, and needles grew directly from the reaction mixture after two days. After keeping in the fridge for one further night, the reaction mixture was filtered off and the white needles were rinsed several times with cold ethanol. NMR and single crystal X-ray studies confirmed the formation of the Zn complex 10 (66.4 mg, 55%). <sup>49</sup> <sup>1</sup>H-NMR (500 MHz, D<sub>2</sub>O):  $\delta = 1.74-1.78$  (m, 2 H, H(g), 1.97-2.01 (m, 6 H, H(d) and H(a)), 2.08-2.12 (m, 6 H, H(d') and H(g')), 2.91-2.98 (m, 12 H, H(b), H(f) and H(e)), 3.03-3.10 (m, 8 H, H(c)) and 3.31-3.34 ppm (m, 8 H, H(f') and H(e')). <sup>13</sup>C-NMR (125 MHz, D<sub>2</sub>O):  $\delta = 25.47$  (C(d)), 26.35 (C(g)), 30.17 (C(a)), 51.77 (C(e)), 52.01 (C(f)), 58.66 (C(b)), 60.02 (C(c)) and 121.88 ppm (q, J = 318.4 Hz, CF<sub>3</sub>). Elemental analysis:  $C_{21}H_{46}N_6Zn_2(CF_3SO_3)_2(OH)_2\cdot(CH_3CH_2OH)_{0.5}$  calc. C (33.19%), H (5.92%), N (9.67%); exp. C (33.01%), H (6.17%), N (9.88%). MS (ESI): m/z (%) = 533 [LH + CF<sub>3</sub>SO<sub>3</sub>H]<sup>+</sup> (100%), 745 [LH +  $Zn(CF_3SO_3)_2$  (85%).

Synthesis of the di-nickel complex of ligand L4 (11). Ligand L4 (51 mg, 0.13 mmol) and nickel triflate (136.6 mg, 0.26 mmol) were refluxed for 60 h in an acetonitrile–methanol–water mixture (10:10:1), yielding a bright blue solution with some solid material. The suspension was filtered while still hot and dried under reduced pressure obtaining a green oil in 90% yield (11). Elemental analysis:  $C_{26}H_{48}F_{12}N_6Ni_2O_{12}S_4\cdot(CH_3CN)$ : calc. C (29.21%), H (4.46%) N (8.52%); exp. C (29.67%), H (4.80%), N (8.58%). IR (Golden Gate ATR):  $\tilde{\nu}=3446$  (NH $_2^+$ , w), 2879 (CH $_2$ , w), 1224 (C–F, s), 1160 (SO $_3$ , s), 1026 (S=O, s) cm $^{-1}$ .

Ligand L4 (61 mg, 0.15 mmol) in dichloromethane was added to a stirred solution of nickel acetate (76.6 mg, 0.31 mmol) in EtOH–MeOH, obtaining a pale green solution. Part of this solution was treated with a saturated solution of NaBPh<sub>4</sub> in water in order to precipitate the dinickel complex of L4. Suitable crystals for X-ray analysis could be grown by slow vapour diffusion of tetrahydropyrane into a solution of the nickel complex in acetonitrile.<sup>31</sup>

Synthesis of the di-copper complex of ligand L4 (12). Copper triflate (92 mg, 0.25 mmol) in ethanol (2 ml), was slowly added to a stirred solution of ligand L4 (50 mg, 0.13 mmol) in ethanol (2 ml). The solution immediately changed from blue to green. The solution was dried under reduced pressure, yielding a green solid in 87% yield (12). Elemental analysis:  $C_{26}H_{48}Cu_2F_{12}N_6O_{12}S_4$ : calc. C (27.88%), H (4.31%), N (7.50%); exp. C (27.91%), H (4.38%) N (7.42%). IR (Golden Gate ATR):  $\tilde{\nu}=3251$  (NH<sub>2</sub><sup>+</sup>, w), 2883 (CH<sub>2</sub>, w), 1220 (C–F, s), 1158 (SO<sub>3</sub>, s), 1022 (S=O, s) cm<sup>-1</sup>.

## Acknowledgements

We thank the Swiss National Science Foundation, the Eidgenössische Stipendien Kommission für ausländische Studierende and the Department of Chemistry, University of Zurich for financial support.

#### Notes and references

- 1 L. Yang, S. Wang, T. Tian and X. Zhou, *Curr. Med. Chem.*, 2012, **19**, 557–568.
- 2 C. K. Singleton, J. Klysik, S. M. Stirdivant and R. D. Wells, *Nature*, 1982, 299, 312–316.
- 3 A. Herbert and A. Rich, *J. Biol. Chem.*, 1996, 271, 11595–11598
- 4 L. F. Liu and J. C. Wang, *Proc. Natl. Acad. Sci. U. S. A.*, 1987, **84**, 7024–7027.
- 5 S. Rothenburg, F. Koch-Nolte, A. Rich and F. Haag, *Proc. Natl. Acad. Sci. U. S. A.*, 2001, **98**, 8985–8990; D.-B. Oh, Y.-G. Kim and A. Rich, *Proc. Natl. Acad. Sci. U. S. A.*, 2002, **99**, 16666–16671.
- 6 B. Wong, S. Chen, J.-A. Kwon and A. Rich, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 2229–2234.
- 7 H. Liu, N. Mulholland, H. Fu and K. Zhao, *Mol. Cell. Biol.*, 2006, 26, 2550–2559.
- M. Steinmetz, D. Stephan and K. F. Lindahl, *Cell*, 1986, 44, 895–904; A. Weinreb, D. R. Katzenberg, G. L. Gilmore and B. K. Birshtein, *Proc. Natl. Acad. Sci. U. S. A.*, 1988, 85, 529–533.
- 9 G. Wang, S. Carbajal, J. Vijg, J. DiGiovanni and K. M. Vasquez, *J. Natl. Cancer Inst.*, 2008, **100**, 1815– 1817.
- 10 A. M. Freund, M. Bichara and R. P. P. Fuchs, *Proc. Natl. Acad. Sci. U. S. A.*, 1989, **86**, 7465–7469.
- D. T. Kha, G. Wang, N. Natrajan, L. Harrison and K. M. Vasquez, J. Mol. Biol., 2010, 398, 471–480.
- 12 G. Wang, L. A. Christensen and K. M. Vasquez, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 2677–2682.
- 13 H. Li, J. Xiao, J. Li, L. Lu, S. Feng and P. Dröge, *Nucleic Acids Res.*, 2009, 37, 2737–2746.
- 14 A. Cerna, A. Cuadrado, N. Jouve, S. M. Diaz de la Espina and C. De la Torre, *Eur. J. Histochem.*, 2004, 48, 49–55.
- H. J. Lipps, A. Nordheim, E. M. Lafer, D. Ammermann,
   B. D. Stollar and A. Rich, *Cell*, 1983, 32, 435–441.
- 16 B. Wittig, T. Dorbic and A. Rich, *Proc. Natl. Acad. Sci. U. S. A.*, 1991, 88, 2259–2263.
- 17 S. Rothenburg, F. Koch-Nolte and F. Haag, *Immunol. Rev.*, 2001, **184**, 286–298.
- 18 R. R. Sinden, *DNA: Structure and Function*, Academic Press, San Diego, 1994.
- 19 B. Spingler, in *Metal Ions and Nucleic Acids*, ed. A. Sigel, H. Sigel and R. K. O. Sigel, Springer Science+Business Media, 2012, vol. 10, pp. 103–118.
- 20 B. Spingler and C. Da Pieve, *Dalton Trans.*, 2005, 1637–1643; M. G. Santangelo, A. Medina-Molner, A. Schweiger, G. Mitrikas and B. Spingler, *J. Biol. Inorg. Chem.*, 2007, 12, 767–775; B. Spingler, C. Da Pieve, A. Medina-Molner, P. M. Antoni and M. G. Santangelo, *Chimia*, 2009, 63, 153–156; M. G. Santangelo, P. M. Antoni, B. Spingler and G. Jeschke, *ChemPhysChem*, 2010, 11, 599–606.
- 21 B. Spingler, F. Zobi, P. M. Antoni, A. Medina-Molner and R. Alberto, *Chimia*, 2005, **59**, 826–831.

- 22 R. Haidar, M. Ipek, B. DasGupta, M. Yousaf and L. J. Zompa, *Inorg. Chem.*, 1997, 36, 3125–3132; B. Graham, L. Spiccia, A. M. Bond, M. T. W. Hearn and C. M. Kepert, *J. Chem. Soc., Dalton Trans.*, 2001, 2232–2238; T. Moufarrej, K. Bui and L. J. Zompa, *Inorg. Chim. Acta*, 2002, 340, 56–64.
- 23 Y. Song, J. Zan, H. Yan, Z.-L. Lu and R. Wang, *Org. Biomol. Chem.*, 2012, **10**, 7714–7720.
- 24 A. Medina-Molner and B. Spingler, *Chem. Commun.*, 2012, 48, 1961–1963.
- 25 G. R. Weisman, D. J. Vachon, V. B. Johnson and D. A. Gronbeck, J. Chem. Soc., Chem. Commun., 1987, 886– 887
- 26 R. W. Alder, D. D. Ellis, R. Gleiter, C. J. Harris, H. Lange, A. G. Orpen, D. Read and P. N. Taylor, J. Chem. Soc., Perkin Trans. 1, 1998, 1657–1668.
- 27 A. Medina-Molner, O. Blacque and B. Spingler, *Org. Lett.*, 2007, **9**, 4829–4831.
- 28 L. J. Zompa, Inorg. Chem., 1978, 17, 2531-2536.
- 29 P. G. Graham, D. C. Weatherburn, F. C. March and W. T. Robinson, *Inorg. Chim. Acta*, 1990, **178**, 227–232.
- 30 V. M. Shelton and J. R. Morrow, *Inorg. Chem.*, 1991, 30, 4295–4299.
- 31 B. Spingler, S. Schnidrig, T. Todorova and F. Wild, *CrystEng-Comm*, 2012, **14**, 751–757.
- 32 B. Spingler and P. M. Antoni, *Chem. Eur. J.*, 2007, **13**, 6617–6622.
- 33 B. Spingler, Inorg. Chem., 2005, 44, 831-833.
- 34 W. Zacharias, J. C. Martin and R. D. Wells, *Biochemistry*, 1983, 22, 2398–2405.
- 35 J. B. Chaires and J. M. Sturtevant, *Biopolymers*, 1988, 27, 1375–1387.
- 36 J. H. van de Sande and T. M. Jovin, *EMBO J.*, 1982, 1, 115–120; W. C. Russell, B. Precious, S. R. Martin and P. M. Bayley, *EMBO J.*, 1983, 2, 1647–1653; M. Govindaraju, F. S. Monica, R. Berrocal, K. R. S. Sambasiva Rao and K. S. Rao, *Curr. Trends Biotechnol. Pharm.*, 2012, 6, 204–209.
- 37 M. Behe and G. Felsenfeld, *Proc. Natl. Acad. Sci. U. S. A.*, 1981, **78**, 1619–1623.
- 38 A. Parkinson, M. Hawken, M. Hall, K. J. Sanders and A. Rodger, *Phys. Chem. Chem. Phys.*, 2000, 2, 5469–5478.
- 39 H. Irving and R. J. P. Williams, Nature, 1948, 162, 746-747.
- 40 G. Barone, A. Terenzi, A. Lauria, A. M. Almerico, J. M. Leal, N. Busto and B. Garcia, *Coord. Chem. Rev.*, 2013, 257, 2848– 2862.
- 41 A. L. Gavrilova and B. Bosnich, *Chem. Rev.*, 2004, **104**, 349–383.
- 42 E. Schlittler and J. Hohl, Helv. Chim. Acta, 1952, 35, 29-45.
- 43 STOE-IPDS Software package, STOE & Cie, GmbH, Darmstadt, Germany, 1998.
- 44 G. M. Sheldrick, Acta Crystallogr., Sect. A: Fundam. Crystallogr., 1990, 46, 467–473.
- 45 A. Altomare, M. C. Burla, M. Camalli, G. L. Cascarano, C. Giacovazzo, A. Guagliardi, A. G. G. Moliterni, G. Polidori and R. Spagna, *J. Appl. Crystallogr.*, 1999, 32, 115– 119.

Paper

- 46 G. M. Sheldrick, Acta Crystallogr., Sect. A: Fundam. Crystallogr., 2008, 64, 112–122.
- 47 C. Da Pieve, A. Medina-Molner and B. Spingler, *Synthesis*, 2007, 679–682.
- 48 E. Kimura, T. Koike, M. Shionoya and M. Shiro, *Chem. Lett.*, 1992, 21, 787–790.
- 49 A. A. Neverov, Z.-L. Lu, C. I. Maxwell, M. F. Mohamed, C. J. White, J. S. W. Tsang and R. S. Brown, *J. Am. Chem. Soc.*, 2006, 128, 16398–16405; Z.-L. Lu, C. T. Liu, A. A. Neverov and R. S. Brown, *J. Am. Chem. Soc.*, 2007, 129, 11642–11652; S. E. Bunn, C. T. Liu, Z.-L. Lu, A. A. Neverov and R. S. Brown, *J. Am. Chem. Soc.*, 2007, 129, 16238–16248.