

Cite this: *Dalton Trans.*, 2015, **44**, 46Received 20th October 2014,  
Accepted 22nd October 2014

DOI: 10.1039/c4dt03239j

www.rsc.org/dalton

Redox-active multinuclear Pd(II) complexes with  
bis- and tris-mesoionic carbenes†

Ramananda Maity, Margarethe van der Meer and Biprajit Sarkar\*

**Synthesis of a ligand platform to generate di- and tri-mesoionic carbenes is reported together with their multinuclear Pd(II) complexes. Complete structural characterization and preliminary electrochemical data are presented.**

N-heterocyclic carbenes (NHCs) are privileged ligands in organometallic chemistry.<sup>1</sup> While homogeneous catalysis has traditionally dominated NHC research,<sup>1</sup> these ligands are useful in a variety of other fields, as has been seen in recent years for supramolecular chemistry.<sup>2</sup> Their mesoionic carbene (MIC) counterparts are currently gaining immense popularity.<sup>3</sup> We have recently used metal complexes of these ligands as catalysts for the “click” reactions,<sup>4a-c</sup> Suzuki–Miyaura cross-coupling reactions,<sup>4d</sup> reduction of aromatic nitro compounds<sup>4e</sup> and for oxidation catalysis.<sup>4f,g</sup>

Most work on triazolylidene ligands has been on their monodentate forms. Metal complexes of these ligands have been in general used for a variety of homogeneous catalytic processes.<sup>3-5</sup>

Only a few bis-MIC ligands possessing a metal center at each MIC donor have been reported in the literature.<sup>6,7</sup> There have been no reports on tris-MIC ligands and their complexes till now. Bis-carbene ligands can have certain advantages over their mono-counterparts.<sup>8</sup> We present here the synthesis of a phenylene based ligand platform for generating MIC ligands of various denticity and report on the mono-, di- and trinuclear Pd(II)-complexes with these ligands. Apart from synthetic and structural aspects, we also present preliminary results on the redox properties of these complexes, a field that has been largely ignored to date for metal complexes of MIC ligands.<sup>3b</sup>

Reaction of **1**<sup>3a</sup> with PdCl<sub>2</sub> in the presence of a base resulted in the formation of the mononuclear complex **[2]**, which was

characterized by <sup>1</sup>H and <sup>13</sup>C NMR spectroscopy. The related mononuclear PEPPSI<sup>1e</sup> type complexes with MIC ligands have been reported earlier in the literature.<sup>5a,7c</sup> **[2]** was synthesized here to be able to compare a mononuclear case with the di- and trinuclear complexes where all substituents on ligands are the same. The reaction of the bis-triazolium salt **3**<sup>6</sup> with PdCl<sub>2</sub> in pyridine yielded the dinuclear complex **[4]**.

The complex **[4]** is well soluble in dichloromethane and sparingly soluble in chloroform. The formation of **[4]** was easily monitored by <sup>1</sup>H NMR spectroscopy which showed the absence of the triazolium C5–H signals ( $\delta = 9.25$  ppm)<sup>6</sup> observed for the original bis-triazolium salt. A singlet was observed for the aryl C–H protons at  $\delta = 8.21$  ppm. The resonances for the C–H protons of pyridine rings were detected as multiplets at  $\delta = 8.92$ – $8.93$ ,  $7.60$ – $7.63$  and  $7.21$ – $7.24$  ppm. Upon complex formation the resonance for  $\alpha$ -hydrogen atoms of the pyridine ring ( $\delta = 8.92$ – $8.93$ ) was more downfield shifted compared to their corresponding resonance in free pyridine ( $\delta = 8.62$ ).<sup>9</sup> The <sup>13</sup>C{<sup>1</sup>H} NMR spectrum of **[4]** showed the characteristic resonance for the C<sub>trz</sub>–Pd carbon atoms at  $\delta = 128.3$  ppm, which is not significantly shifted from the C5 resonance of the precursor triazolium salt **3** ( $\delta = 129.6$  ppm).<sup>6</sup> The ESI mass spectrum (positive ions) of **[4]** showed a peak for the cationic complex ion **[[4] – I + CH<sub>3</sub>CN]<sup>+</sup>** which supports the formation of the dipalladium complex **[4]** depicted in Scheme 1.


Single crystals suitable for X-ray diffraction studies were obtained by slow evaporation of the saturated dichloromethane–chloroform solution of **[4]** at ambient temperature. The molecular structure analysis of these crystals confirmed the formation of the dinuclear palladium complex **[4]** where each palladium atom is coordinated by a carbene C-donor and one pyridine donor in a *trans*-fashion. The remaining coordination sites at the Pd<sup>II</sup> center are occupied by two iodido donors (Fig. 1).

The planes of the NHC donors are rotated out of the central aryl ring plane to incorporate the palladium–pyridine moieties. The dihedral angle between the NHC planes and the central aryl ring plane is measured to be 41.97°. Two Pd–Py


Institut für Chemie und Biochemie, Anorganische Chemie, Freie Universität Berlin,  
Fabeckstraße 34–36, D-14195 Berlin, Germany. E-mail: biprajit.sarkar@fu-berlin.de;  
Fax: +49-30-838-53310

† Electronic supplementary information (ESI) available: Preparation and spectroscopic properties of complexes. CV table. CCDC 1019285 (**[4]**·2CHCl<sub>3</sub>), 1019286 (**[7]**·CHCl<sub>3</sub>) and 1021276 (**[9]**). For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c4dt03239j





Scheme 1 Synthesis of complexes [2] and [4].



Scheme 2 Preparation of complexes [7] and [9].




Fig. 1 ORTEP plot of [4]. Ellipsoids are drawn at 50% probability. Hydrogen atoms and solvent molecules have been omitted for clarity.

moieties are oriented in an opposite direction, thus featuring an *anti*-geometry of the complex. The asymmetric unit contains a half formula unit of the complex [4] and a chloroform molecule. The half unit is related to the other half by a crystallographic inversion center. The coordination geometry around the Pd atom is distorted square planar. The C1–Pd1–N4 bond angle is measured to be 178.8(3)°. This value, like the Pd1–C1 bond length (1.967(10) Å), falls in the range previously described for palladium MIC complexes.<sup>5a,7c</sup>

The tris-triazole **5** was prepared by an *in situ* Cu(I) catalyzed click reaction. Subsequent trimethylation of **5** delivered the new tris-triazolium salt **6** (see ESI†). For comparison purposes we also used the tris-imidazolium salt **8**<sup>2d</sup> to prepare the corresponding trinuclear Pd(II) complex. The reaction of **1** equiv. of the azolium salts **6** or **8** with 3 equiv. of PdCl<sub>2</sub> in the presence of K<sub>2</sub>CO<sub>3</sub> in pyridine led to the formation of trinuclear complexes [7] and [9], respectively (Scheme 2). Complexes [7] and [9] were obtained in good yields of 60.5 and 72%, respectively. Both complexes are stable towards air and moist-

ure in the solid state. These trinuclear complexes are well soluble in dichloromethane. Changing the amount of the base did not result in the formation of orthometalated complexes.<sup>6</sup>

The formation of complexes [7] and [9] was confirmed by <sup>1</sup>H, <sup>13</sup>C, 2D correlation NMR spectroscopy and mass spectrometry (ESI†). The <sup>1</sup>H NMR spectra of both complexes show the absence of azolium C–H protons which were observed at δ = 9.27 (**6**) and 10.07 (**8**) ppm for their parent azolium salts. The resonance for the aryl C–H protons (δ = 9.68 ppm) in complex [9] is significantly downfield shifted compared to their corresponding resonance (δ = 8.95 ppm) in complex [7]. The resonances for the characteristic carbene carbon atoms were observed at δ = 133.4 and 147.9 ppm in the <sup>13</sup>C{<sup>1</sup>H} NMR spectra of complexes [7] and [9], respectively. The resonance for the imidazol-2-ylidene carbon atoms (147.9 ppm for [9]) appeared more downfield shifted compared to 1,2,3-triazol-5-ylidene carbon atoms (δ = 133.4 ppm for [7]). These resonances fall in the range reported for mono- and dinuclear palladium(II) complexes of similar type.<sup>5a,10</sup> The resonance for the α-carbon atoms of the pyridine rings appeared downfield shifted ([7]: δ = 154.5 ppm; [9]: 154.4 ppm) compared to their corresponding resonances in free pyridine.<sup>9</sup>

Single crystals of both complexes [7] and [9] were obtained by slow evaporation of the solvents from a concentrated dichloromethane–chloroform solution at room temperature. X-ray diffraction analysis of these crystals confirmed the formation of trinuclear palladium complexes [7] (Fig. 2) and [9] (Fig. 3).

The metric parameters for complex [7] are almost identical to the previously described dinuclear complex [4]. To coordinate with palladium atoms the NHC donor planes are rotated out of the central aryl ring plane by 34.21 to 44.04° in complex [7] and by 35.56 to 48.26° in complex [9]. The Pd–C<sub>MIC</sub> bond distances (range: 1.950(5)–1.952(5) Å) are slightly shorter in






Fig. 2 ORTEP plot of [7]. Ellipsoids are drawn at 50% probability. Hydrogen atoms and solvent molecules have been omitted for clarity.




Fig. 3 ORTEP plot of [9]. Ellipsoids are drawn at 50% probability. Hydrogen atoms and solvent molecules have been omitted for clarity. One of the iodine atoms is disordered over two positions and only one position is shown.

complex [7] compared to their corresponding bond distances in complex [9] (range: 1.962(3)–1.970(3) Å). The Pd–C<sub>NHC</sub> in [9] bond distances are in similar range described for previously reported palladium NHC complexes.<sup>11</sup> The N–C<sub>NHC</sub>–N bond angles (range: 105.2(3)–105.5(5)°) in [9] are slightly larger than C–C<sub>MIC</sub>–N bond angles (range: 102.6(5)–102.9(4)°) in [7].

Mononuclear Pd(II)-MIC complexes of the PEPPSI type have been used as pre-catalysts for cross-coupling reactions.<sup>5a,7c</sup> The initial step in these catalytic cycles is believed to be reductive elimination which results in the *in situ* generation of the catalytically active Pd(0) species. In general, electrochemistry can help in addressing questions related to such elementary steps, as has been shown for studies with Pd-NHC and Pd-phosphine complexes.<sup>12</sup> To the best of our knowledge no electrochemical investigation has been carried out on Pd-MIC complexes so far.<sup>3b–d,5a,7c</sup> We were interested in comparing the electrochemical responses of the complexes as a function of their nuclearity and we wanted to compare the results of the Pd-MIC complexes with that of the tris-palladium NHC counterpart [9]. The mononuclear complex [2] displays an irreversible reduction peak at –1.91 V (Fig. 4). The corresponding re-oxidation peak is observed at –0.11 V. This oxidation peak shows up only after the reductive side has been scanned proving this to be a follow-up of the reduction peak at –1.91 V. This




Fig. 4 Cyclic voltammograms of the complexes in DMF–0.1 M Bu<sub>4</sub>NPF<sub>6</sub> at 298 K. Scan rate: 250 mV s<sup>-1</sup>. Fc/Fc<sup>+</sup> was used as an internal standard.

process is assigned to the reduction of Pd(II) to Pd(0) combined with the ligand loss, making this step irreversible.<sup>12</sup> The complexes [4] and [7] also display an irreversible reduction step at potentials comparable to that of [2] (Fig. 4 and Table S4†). These processes are also associated with re-oxidation steps at more positive potentials just like for [2]. However, for the dinuclear complex [4] one further reduction step is observed and for the trinuclear complex [7] two further reduction steps are observed. These data thus likely point to a stepwise reduction of the Pd(II) centers in these multinuclear complexes suggesting strong electrochemical coupling among them. For the trinuclear complex [9] containing a tris-imidazol-2-ylidene ligand, only one irreversible reduction step is observed at –1.80 V indicating a different electronic situation in [9] as compared to [4] and [7].

In conclusion, we have presented here a synthetic strategy for building multinuclear complexes with MIC ligands based on a phenylene platform. [7] is the first example of a trinuclear complex containing MIC donors. Electrochemical data clearly show that the mononuclear complex [2] undergoes a reductive elimination step, and that electrochemical communication exists between the palladium centers in the multinuclear complexes [4] and [7]. These results differ from the electrochemical response of the imidazole-2-ylidene containing complex [9] where no such communication is observed. In view of the modular synthesis for introducing additional triazolylidene units into the phenylene rings, and the lability of the pyridine ligands in these PEPPSI complexes, the systems reported here are likely to be appropriate synthons for generating supramolecular assemblies. Additionally, the electrochemical data presented here should be helpful in deciphering mechanisms (including cooperativity) of cross-coupling reactions with these



and other Pd-MIC complexes. Current research in our laboratory is focused in these directions.

The authors thank the Fonds der Chemischen Industrie (FCI) and the Freie Universität Berlin for financial support.

## Notes and references

- For selected reviews see: (a) P. de Frémont, N. Marion and S. P. Nolan, *Coord. Chem. Rev.*, 2009, **253**, 862; (b) M. Melaimi, M. Soleilhavoup and G. Bertrand, *Angew. Chem., Int. Ed.*, 2010, **49**, 8810; (c) F. E. Hahn and M. C. Jahnke, *Angew. Chem., Int. Ed.*, 2008, **47**, 3122; (d) W. A. Herrmann, *Angew. Chem., Int. Ed.*, 2002, **41**, 1290; (e) E. A. B. Kantchev, C. J. O'Brien and M. G. Organ, *Angew. Chem., Int. Ed.*, 2007, **46**, 2768; (f) S. Díez-González, N. Marion and S. P. Nolan, *Chem. Rev.*, 2009, **109**, 3612; (g) C. C. Loh and D. Enders, *Chem. – Eur. J.*, 2012, **18**, 10212; (h) J. D. Egbert, C. S. J. Cazin and S. P. Nolan, *Catal. Sci. Technol.*, 2013, **3**, 912.
- For selected examples see: (a) C. Radloff, F. E. Hahn, T. Pape and R. Fröhlich, *Dalton Trans.*, 2009, 7215; (b) C. Radloff, J. J. Weigand and F. E. Hahn, *Dalton Trans.*, 2009, 9392; (c) Y. Han, L. J. Lee and H. V. Huynh, *Chem. – Eur. J.*, 2010, **16**, 771; (d) A. Rit, T. Pape and F. E. Hahn, *J. Am. Chem. Soc.*, 2010, **132**, 4572; (e) R. Maity, A. Rit, C. Schulte to Brinke, C. G. Daniliuc and F. E. Hahn, *Chem. Commun.*, 2013, **49**, 1011; (f) C. Segarra, G. Guisado-Barrios, F. E. Hahn and E. Peris, *Organometallics*, 2014, **30**, 5077.
- (a) P. Mathew, A. Neels and M. Albrecht, *J. Am. Chem. Soc.*, 2008, **130**, 13534; (b) K. F. Donnelly, A. Petronilho and M. Albrecht, *Chem. Commun.*, 2013, **49**, 1145; (c) R. H. Crabtree, *Coord. Chem. Rev.*, 2013, **257**, 755; (d) J. M. Aizpurua, R. M. Fratila, Z. Monasterio, N. Perez-Esnaola, E. Andreieff, A. Irastorza and M. Sagartzazu-Aizpurua, *New J. Chem.*, 2014, **38**, 474; (e) D. Schweinfurth, N. Deibel, F. Weisser and B. Sarkar, *Nat. Chem.*, 2011, **59**, 937; (f) G. Guisado-Barrios, J. Bouffard, B. Donnadiou and G. Bertrand, *Angew. Chem., Int. Ed.*, 2010, **49**, 4759; (g) J. D. Crowley, A. Lee and K. J. Kilpin, *Aust. J. Chem.*, 2011, **64**, 1118.
- (a) S. Hohloch, C.-Y. Su and B. Sarkar, *Eur. J. Inorg. Chem.*, 2011, 3067; (b) S. Hohloch, D. Scheiffele and B. Sarkar, *Eur. J. Inorg. Chem.*, 2013, 3956; (c) S. Hohloch, B. Sarkar, L. Nauton, F. Cisnetti and A. Gautier, *Tetrahedron Lett.*, 2013, **54**, 1808; (d) S. Hohloch, W. Frey, C.-Y. Su and B. Sarkar, *Dalton Trans.*, 2013, **42**, 11355; (e) S. Hohloch, L. Suntrup and B. Sarkar, *Organometallics*, 2013, **32**, 7376; (f) S. Hohloch, L. Hettmanczyk and B. Sarkar, *Eur. J. Inorg. Chem.*, 2014, 3164; (g) A. Bolje, S. Hohloch, D. Urankar, A. Pevec, M. Gazvoda, B. Sarkar and J. Kosmrlj, *Organometallics*, 2014, **33**, 2588.
- For selected examples see: (a) D. Canseco-Gonzalez, A. Gniewek, M. Szulmanowicz, H. Müller-Bunz, A. M. Trzeciak and M. Albrecht, *Chem. – Eur. J.*, 2012, **18**, 6055–6062; (b) T. Nakamura, T. Terashima, K. Ogata and S. Fukuzawa, *Org. Lett.*, 2011, **13**, 620; (c) J. R. Wright, P. C. Young, N. T. Lucas, A.-L. Lee and J. D. Crowley, *Organometallics*, 2013, **32**, 7065; (d) J. Huang, J.-T. Hong and S. H. Hong, *Eur. J. Org. Chem.*, 2012, 6630; (e) E. C. Keske, O. V. Zenkina, R. Wang and C. M. Crudden, *Organometallics*, 2012, **31**, 6215.
- R. Maity, S. Hohloch, C.-Y. Su, M. van der Meer and B. Sarkar, *Chem. – Eur. J.*, 2014, **20**, 9952–9961.
- (a) M. T. Zamora, M. J. Farguson and M. Cowie, *Organometallics*, 2012, **31**, 5384; (b) J. Cai, X. Yang, K. Arumugam, C. W. Bielawski and J. L. Sessler, *Organometallics*, 2011, **30**, 5033; (c) E. C. Keske, O. V. Zenkina, R. Wang and C. M. Crudden, *Organometallics*, 2012, **31**, 456; (d) K. J. Kilpin, U. S. D. Paul, A.-L. Lee and J. D. Crowley, *Chem. Commun.*, 2011, **47**, 328.
- (a) G. Guisado-Barrios, J. Hiler and E. Peris, *Chem. – Eur. J.*, 2013, **19**, 10405; (b) S. Sabater, J. A. Mata and E. Peris, *Eur. J. Inorg. Chem.*, 2013, 4764; (c) S. Gonell, M. Poyatos and E. Peris, *Angew. Chem., Int. Ed.*, 2013, **125**, 7147.
- G. R. Fulmer, A. J. M. Miller, N. H. Sherden, H. E. Gottlieb, A. Nudelman, B. M. Stoltz, J. E. Bercaw and K. I. Goldberg, *Organometallics*, 2010, **29**, 2176.
- (a) G. Wang, G. Liu, Y. Du, W. Li, S. Yin, S. Wang, Y. Shi and C. Cao, *Transition Met. Chem.*, 2014, **39**, 691; (b) S. Ruiz-Botella and E. Peris, *Organometallics*, 2014, **33**, 5509.
- (a) R. Maity, H. Koppetz, A. Hepp and F. E. Hahn, *J. Am. Chem. Soc.*, 2013, **135**, 4966; (b) R. Maity, A. Rit, C. Schulte to Brinke, J. Kösters and F. E. Hahn, *Organometallics*, 2013, **32**, 6174.
- For a detailed review see: A. Jutand, *Chem. Rev.*, 2008, **108**, 2300.

