

Showcasing collaborative research from the Youichi Ishii lab at Chuo University and Noriko Tsuchida lab at Saitama Medical University.

Title: Competition between vinylidene rearrangement and 1,2-insertion of carbon-disubstituted internal alkynes at a Cp*Ir(III) complex

Even carbon-disubstituted internal alkynes can select the "vinylidene rearrangement route". Vinylidene rearrangement of diarylacetylenes in an aryl-Ir(III) complex has been shown to become a kinetically (and thermodynamically) more favorable path than common 1,2-insertion.

As featured in:



See Youichi Ishii et al. Dalton Trans., 2015, **44**, 17448.



www.rsc.org/dalton

Dalton Transactions



View Article Online

COMMUNICATION

CrossMark

Cite this: Dalton Trans., 2015, 44, 17448

Received 10th June 2015, Accepted 20th August 2015 DOI: 10.1039/c5dt02207j

www.rsc.org/dalton

Competition between vinylidene rearrangement and 1,2-insertion of carbon-disubstituted internal alkynes at a Cp*Ir(III) complex*

Yousuke Ikeda,^a Shintaro Kodama,^a Noriko Tsuchida^b and Youichi Ishii*^a

Competition between vinylidene rearrangement/1,1-insertion and 1,2-alkyne insertion into the Ir-Ar bond of $[Cp*Ir(ppy-F_4)]^+$ was observed on reaction with diarylacetylenes. The former process afforded the iridacycle 2 via the subsequent 1,4-Ir migration, whereas the latter led to the pyridoisoquinolinium complex 4. Detailed analysis revealed that 4 isomerizes to 2 by heating at 50 °C.

Vinylidene rearrangement is now recognized as one of the most valuable methods for the transformation of terminal alkynes.¹ Details of this tautomerization involving mechanistic studies^{1*f*,2} and applications to organic synthesis¹ have been well discussed. The rich reactivities of vinylidenes as well as the accumulated mechanistic information concerning their formation brought about interest in the vinylidene rearrangement of more general alkynes.¹¹ However, the tautomerization of internal alkynes is still not recognized as a common process,^{1b} although a few examples of the internal alkyne-disubstituted vinylidene rearrangement have been studied by us and other groups during the last decade.^{3,4} This is partly because the vinylidene rearrangement of internal alkynes is a slower process than that of terminal alkynes, hence cannot compete favorably with other metal promoted reactions. For example, internal alkynes readily insert into the metal-carbon bond of an alkyl- or aryl-metal complex in a 1,2 fashion, and therefore the vinylidene rearrangement of the alkynes does not take place. A representative reaction is found in Cp*Rh(III)-mediated annulation of 2-phenylpyridine with internal alkynes, which is first developed by Jones in 2008,⁵ and the mechanism of this

interesting reaction has been proposed to involve the first ortho-metallation of phenylpyridine to form a five-membered metallacycle [Cp*RhCl(ppy)] (ppy = 2-(2-pyridyl)phenyl) followed by the 1,2-insertion of the alkyne into the Rh-aryl bond and the reductive elimination of a C-N bond. This strategy has been applied to a variety of substrates other than 2-phenylpyridine and also expanded to catalytic versions.⁶ Needless to say, vinylidene rearrangement of internal alkynes and subsequent 1,1-insertion of the disubstituted vinylidene ligand has never been observed throughout these studies.

Obviously, the general kinetic trend that the 1,2-insertion of internal alkynes is faster than their vinylidene rearrangement/ 1,1-insertion limits the synthetic utility of the latter process. To overcome this drawback and broaden applicability of the vinylidene rearrangement, we have explored how we can control the preference between these processes. As a model system, we have adopted $[Cp*IrCl(ppy-F_4)]$ (1) (ppy-F₄ = 2,3,4,5tetrafluoro-6-(2-pyridyl)phenyl), in which the C_6F_4 group is expected to bind to the metal more strongly than the C₆H₄ group in ppy and hence to slow down the 1,2-insertion. To our surprise, the products derived from the vinylidene rearrangement of diarylacetylenes were obtained in preference to those from the 1,2-insertion as the thermodynamic product, but not necessarily the kinetic product. This also provides the first example of vinylidene rearrangement of internal alkynes at metal complexes other than group 8 metals.

The iridium precursor 1 was readily synthesized by the reaction of [Cp*IrCl₂]₂ with 2-(2,3,4,5-tetrafluorophenyl)pyridine in the presence of NaOAc·3H2O and fully characterized by spectroscopic as well as crystallographic analysis (see ESI[†]). When 1 was allowed to react with diphenylacetylene and NaBAr^F₄ in $C_2H_4Cl_2$ (1,2-dichloroethane) at 50 °C for 4 h, the colour of the reaction mixture turned from yellow to dark purple (Scheme 1). Recrystallization of this mixture afforded the ninemembered iridacycle complex 2a with an Ir-(vinyl CH) agostic interaction as dark purple crystals in 87% isolated yield, and 2a was fully characterized by means of NMR analysis as well as a single-crystal X-ray diffraction study (Fig. 1, left). In the ¹H NMR spectrum, the vinyl CH signal of **2a** appears in a

Dpen Access Article. Published on 20 August 2015. Downloaded on 8/22/2025 12:07:42 AM.

^aDepartment of Applied Chemistry, Faculty of Science and Engineering, Chuo University, 1-13-27 Kasuga, Bunkyo-ku, Tokyo, Japan.

E-mail: yo-ishii@kc.chuo-u.ac.jp; Fax: +81-3-3817-1895; Tel: +81-3-3817-1901 ^bDepartment of Liberal Arts, Faculty of Medicine, Saitama Medical University, 38 Morohongo, Moroyama-machi, Iruma-gun, Saitama 350-0495, Japan. E-mail: n_tsuchi@saitama-med.ac.jp; Fax: +81-49-276-1901; Tel: +81-49-276-1901 †Electronic supplementary information (ESI) available: Text, figures and CIF files giving experimental procedures and crystallographic data. CCDC 1052465-1052470. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c5dt02207j



Scheme 1 Reaction of 1 with diphenylacetylene derivatives and $NaBAr^{F}_{4}$ and deprotonation of 2a to form 3.



Fig. 1 ORTEP drawings of 2a (left) and 3 (right). Anionic part and hydrogen atoms except for H1 of 2a are omitted for clarity. Selected bond lengths (Å) and angles (°): 2a, Ir1–N1, 2.121(5); Ir1–C1, 2.213(5); Ir1–C4, 2.026(6); Ir1–H1, 1.893; C1–C2, 1.461(9); C1–C2–C3, 114.0(5); C1–C2– C9, 123.0(6); C3–C2–C9, 122.5(6); C2–C1–C15, 119.3(5). 3, Ir1–N1, 2.085(3); Ir1–C1, 2.033(5); Ir1–C4, 2.056(4); C1–C2, 1.346(6); Ir1–C1– C2, 120.3(3); Ir1–C1–C15, 112.8(3); C2–C1–C15, 126.0(4); C1–C2–C3, 112.5(4); C1–C2–C9, 125.4(4); C3–C2–C9, 121.7(4).

significantly low-field region (δ 4.16) as an agostic CH, whereas its ¹³C¹H NMR signal exhibited notable high-field shift (δ 49.3). It is interesting to note that these spectroscopy data are in marked difference to those of the related (o-vinyl)aryliricomplex $[Cp*Ir{o-C_6H_4C(Ph)=CHPh}(PMe_3)][BAr_4]$ dium which showed its vinyl CH signal at δ –0.30 in the ¹H NMR and δ 86.4 in the ¹³C{¹H} NMR spectra.⁷ The molecular structure of 2a shows that the Ir1-C1 distance at 2.214(5) Å is explicitly shorter than common agostic iridium-CH distances,^{7,8} suggesting that the iridium center of 2a interacts more strongly with the C1 atom than the common Ir-CH agostic interaction. In fact, complex 2a easily dissociates the agostic vinyl proton by stirring in MeOH at room temperature for 30 min to form the neutral iridium(m) complex 3 in 73% isolated yield (Scheme 1). Complex 3 was fully characterized by spectroscopic as well as crystallographic analysis (Fig. 1, right). Importantly, the two Ph groups of 2a are bound to the same carbon atom (C2), indicating that one of the Ph groups of diphenylacetylene

has migrated across the C=C bond during the reaction. In addition, one Ph group is ortho-metallated by the iridium center, and the H atom is transferred to the vinyl carbon (C1). These results clearly demonstrate that the formation of 2a occurred through the initial vinylidene rearrangement to generate the diphenylvinylidene intermediate followed by the 1,1insertion of the vinylidene ligand into the Ir-C₆F₄ bond and 1,4-Ir(\mathbf{m}) migration^{7,9} to the *ortho* position of the Ph group originated from the alkyne (vide infra). Similarly, para-substituted diphenylacetylene derivatives p-XC₆H₄C \equiv CC₆H₄X-p (X = Me, Cl) were applied to this transformation, and the corresponding cyclometallated complexes 2b and 2c were formed in good yield under appropriate conditions (Scheme 1). Apparently, in this system, vinylidene rearrangement of internal alkynes was more favored than 1,2-insertion into the Ir-Carvl bond. It should also be mentioned that the present reaction provides the first example of the vinylidene rearrangement of carbondisubstituted internal alkynes at an Ir complex;¹⁰ related reactions have so far been observed only at group 8 metal complexes.3,4

To gain deeper insight into this reaction, we monitored the progress of the formation of 2c at 50 °C in CDCl₃ by means of ¹H NMR.¹¹ After 30 min, **1** was consumed completely, and two Cp* signals were observed at δ 1.58 and 1.63 in the intensity ratio of 1:2.1. The former signal is assigned to 2c, whereas the latter species was isolated in 50% yield as yellow crystals by column chromatography on silica gel and characterized unambiguously by a X-ray diffraction study (see ESI†) as the Ir(1) η^4 -pyridoisoquinolinium complex **4c** (Scheme 2). A structurally related rhodium complex was already reported by Huang.^{6a} Obviously complex **4c** is formed from the 1,2-insertion of the alkyne into the Ir-C₆F₄ bond to form the seven-membered metallacycle instead of the vinylidene rearrangement, and this species was further isomerized to **4c** through the reductive elimination (*vide infra*).

Surprisingly, as the reaction proceeded, isomerization of **4c** to **2c** was observed, and it was completed after 10 h (Scheme 3). Thus, complex **4c** is kinetically formed at the early stage of this reaction, and on maintaining at 50 °C, **4c** slowly isomerizes to **2c**, the thermodynamic product of this reaction. Similarly, Cp*Ir(1) complex **4a** was observed at the early stage of the reaction of **1** with PhC=CPh as a minor species and isomerized to **2a** over 4 h (Scheme 3), although it could not be isolated in a pure form (Scheme 2).¹² On the other hand, the



Scheme 2 Formation of complexes 4.



Scheme 3 Isomerization of 4 to 2 at 50 °C.

formation of **2b** was so fast that the corresponding **4b** could not be observed.

The above observation indicates that the rate of the formation of 2 is notably enhanced by introducing electron donating groups into the para positions of the diphenylacetylene. Recently we have shown both experimentally and theoretically that the internal alkyne-disubstituted vinylidene rearrangement at [CpRu(dppe)]⁺ is facilitated by an electrondonating substituent,^{4b,e} and this tendency is in good agreement with the present observations, suggesting that the present vinylidene rearrangement at the Ir(m) center involves nucleophilic aryl migration in the rate determining step. In addition, the ppy- F_4 ligand is essential for the formation of 2; the reaction of [Cp*IrCl(ppy)] (5) with PhC=CPh in the presence of NaBArF4 at 50 °C for 30 min resulted in selective formation of 6 as the sole product in 67% yield, and the corresponding vinylaryliridium species was not formed any more (Scheme 4). For a better understanding of the above observations, preliminary density functional theory (DFT) calculations on the cationic part of 2a and 4a were performed with the B3PW91 functional. As expected, it was confirmed that 2a is more stable in energy than 4a by 8.52 kcal mol⁻¹, which gives good explanation for the selective formation of 2 as the thermodynamic product (Scheme 5). Although we must await a more detailed theoretical study, several mechanisms are considered to be plausible for the conversion of 4 to 2. One is the C-N bond oxidative addition followed by the β -carbon elimination (back-reaction from 4) to regenerate the η^2 -alkyne complex,¹³ which then undergo vinylidene rearrangement. β-Carbon elimination from a vinyl complex to form the corresponding η^2 -alkyne complex is a rare process, but some examples are known in the literature.^{4f,14} Alternatively, direct isomerization of the seven-membered iridacycle A may be operative. In this case, concerted migration of the iridium center and an aryl group (Fig. 2(a)) or an aryl group migration to eliminate C₅H₄N-C₆F₄⁻ anion followed by its nucleophilic attack at the vinylidene α -carbon (Fig. 2(b)) is assumed to be involved.15



Scheme 4 Reactions of 5 with PhC=CPh



Scheme 5 Plausible mechanism of formation of 2 and 4, and Gibbs energy differences between 2a and 4a.



Fig. 2 Possible structures for (a) the transition state in a concerted migration of A into B (Scheme 5) and (b) the intermediate derived from aryl migration of A to eliminate $C_5H_4N-C_6F_4^-$ anion.

Finally, we have investigated the reactivity of **1** with acyl alkyne, which is known to be a reactive substrate towards a vinylidene rearrangement.^{3,4c} When a mixture of **1** with PhC=CCOPh and NaBAr^F₄ in $C_2H_4Cl_2$ was allowed to react at 50 °C for 30 min, a mixture of yellow and red crystals was obtained after recrystallization. X-ray diffraction studies disclosed that the yellow product is the ten-membered iridacycle complex **7**, while the red one is attributed to the vinyliridium complex **8** (see ESI†).¹⁶ These complexes could be separated by column chromatography on silica gel and isolated in 36% and 38% yields, respectively.

Judging from these structures, the 1,2-insertion and vinylidene rearrangement of PhC=CCOPh competitively occurred to generate the seven-membered vinyliridium species **9** and the iridium vinylidene species **10**. Complex **7** was formed from **9** by the 1,4-Ir migration from the vinyl to the *ortho* position of the COPh group, whereas **8** was produced by the 1,1-insertion of the vinylidene ligand in **10** into the Ir-C₆F₄ bond and the



final coordination of the oxygen atom of the carbonyl group (Scheme 6). Unlike the reaction with diarylalkynes, the ratio of 7 and 8 was not changed by further heating at 50 °C.¹⁷

In conclusion, we have revealed that the reaction of $[Cp*Ir-(ppy-F_4)]^+$ with diphenylacetylene derivatives afforded the ninemembered metallacycle complex 2 by way of the sequential vinylidene rearrangement, 1,1-insertion of the vinylidene ligand, and the 1,4-Ir migration to the Ar group. Detailed analysis of this reaction disclosed that 2 and the Ir(I) pyridoisoquinolinium complex 4, the latter of which is a normal 1,2insertion-reductive elimination product, are competitively generated at the early stage of the reaction, but 4 is gradually isomerized to 2, indicating that 2 is the thermodynamically favoured product. The present reaction provides not only the first example of the internal alkyne-disubstituted vinylidene rearrangement at an iridium complex but also a rare example of actual observation of the competition between vinylidene rearrangement and 1,2-insertion of internal alkynes.

We thank the Research Center for Computational Science in Okazaki, Japan, for the use of the computer facilities. This research was financially supported by JST ACT-C. Y. Ikeda thanks the Japan Society for the Promotion of Science (JSPS) Research Fellowships for Young Scientists.

Notes and references

 For reviews, see: (a) Metal Vinylidenes and Allenylidenes in Catalysis: From Reactivity to Applications in Synthesis, ed. C. Bruneau and P. H. Dixneuf, Wiley-VCH, Weinheim, Germany, 2008; (b) G. Marinelli, W. E. Streib, J. C. Huffma, K. G. Caulton, M. R. Gagné, J. Takats, M. Dartiguenave, C. Chardon, S. A. Jackson and O. Eisenstein, Polyhedron, 1990, 9, 1867–1881; (c) M. I. Bruce, Chem. Rev., 1991, 91, 197–257; (d) C. Bruneau and P. H. Dixneuf, Acc. Chem. Res., 1999, 32, 311-323; (e) P. Valerga and M. C. Puerta, *Coord. Chem. Rev.*, 1999, **193-195**, 977-1025; (f) Y. Wakatsuki, *J. Organomet. Chem.*, 2004, **689**, 4092-4109; (g) V. Cadierno, M. P. Gamasa and J. Gimeno, *Coord. Chem. Rev.*, 2004, **248**, 1627-1657; (h) B. M. Trost, M. U. Frederiksen and M. T. Rudd, *Angew. Chem., Int. Ed.*, 2005, **44**, 6630-6666; (*i*) C. Bruneau and P. H. Dixneuf, *Angew. Chem., Int. Ed.*, 2006, **45**, 2176-2203; (*j*) J. A. Varela and C. Saá, *Chem. – Eur. J.*, 2006, **12**, 6450-6456; (*k*) B. M. Trost and A. McClory, *Chem. – Asian J.*, 2008, **3**, 164-194; (*l*) J. M. Lynam, *Chem. – Eur. J.*, 2010, **16**, 8238-8247.

- 2 O. J. S. Pickup, I. Khazal, E. J. Smith, A. C. Whitwood and J. M. Lynam, *Organometallics*, 2014, 33, 1751–1761.
- 3 (a) P. J. King, S. A. R. Knox, M. S. Legge, A. G. Orpen, J. N. Wilkinson and E. A. Hill, *J. Chem. Soc., Dalton Trans.*, 2000, 1547–1548; (b) M. J. Shaw, S. W. Bryant and N. Rath, *Eur. J. Inorg. Chem.*, 2007, 3943–3946; (c) E. Bustelo, I. de los Rios, M. C. Puerta and P. Valerga, *Organometallics*, 2010, 29, 1740–1749; (d) V. K. Singh, E. Bustelo, I. de los Ríos, I. Macías-Arce, M. C. Puerta, P. Valerga, M. A. Ortuño, G. Ujaque and A. Lledós, *Organometallics*, 2011, 30, 4014–4031; (e) F. E. Fernández, M. C. Puerta and P. Valerga, *Inorg. Chem.*, 2013, 52, 6502–6509.
- 4 (a) Y. Ikeda, T. Yamaguchi, K. Kanao, K. Kimura, S. Kamimura, Y. Mutoh, Y. Tanabe and Y. Ishii, J. Am. Chem. Soc., 2008, 130, 16856–16857; (b) Y. Mutoh, Y. Ikeda, Y. Kimura and Y. Ishii, Chem. Lett., 2009, 38, 534–535; (c) Y. Mutoh, K. Imai, Y. Kimura, Y. Ikeda and Y. Ishii, Organometallics, 2011, 30, 204–207; (d) Y. Mutoh, Y. Kimura, Y. Ikeda, N. Tsuchida, K. Takano and Y. Ishii, Organometallics, 2012, 31, 5150–5158; (e) M. Otsuka, N. Tsuchida, Y. Ikeda, Y. Kimura, Y. Mutoh, Y. Ishii and K. Takano, J. Am. Chem. Soc., 2012, 134, 17746–17756; (f) Y. Ikeda, Y. Mutoh, K. Imai, N. Tsuchida, K. Takano and Y. Ishii, Organometallics, 2013, 32, 4353–4358; (g) M. Otsuka, N. Tsuchida, Y. Ishii and K. Takano, J. Ishii and K. Takano, Organometallics, 2015, 34, 3934–3943.
- 5 L. Li, W. W. Brennessel and W. D. Jones, *J. Am. Chem. Soc.*, 2008, **130**, 12414–12419.
- 6 For the recent examples, see: (a) G. Zhang, L. Yang, Y. Wang, Y. Xie and H. Huang, J. Am. Chem. Soc., 2013, 135, 8850–8853; (b) X.-C. Huang, X.-H. Yang, R.-J. Song and J.-H. Li, J. Org. Chem., 2014, 79, 1025–1031; (c) W. Han, G. Zhang, G. Li and H. Huang, Org. Lett., 2014, 16, 3532–3535; (d) C. Ma, C. Ai, Z. Li, B. Li, H. Song, S. Xu and B. Wang, Organometallics, 2014, 33, 5164–5172; (e) D. Ghorai and J. Choudhury, Chem. Commun., 2014, 50, 15159–15162.
- 7 Y. Ikeda, K. Takano, S. Kodama and Y. Ishii, *Organometallics*, 2014, **33**, 3998–4004.
- 8 (a) A. C. Albéniz, G. Schulte and R. G. Crabtree, Organometallics, 1992, 11, 242–249; (b) C. S. Chin, M. Kim, H. Lee, S. Noh and K. M. Ok, Organometallics, 2002, 21, 4785–4793.
- 9 Recent reviews of the 1,4-metal migration, see: (a) S. Ma and Z. Gu, Angew. Chem., Int. Ed., 2005, 44, 7512–7517;

(b) F. Shi and R. C. Larock, Top. Curr. Chem., 2010, 292, 123–164.

- 10 For recent examples concerning Ir-based vinylidene rearrangement of terminal alkynes, see: W. Iali, F. L. Paglia, X.-F. L. Goff, D. Sredojević, M. Pfeffer and J.-P. Djukic, *Chem. Commun.*, 2012, 48, 10310–10312.
- 11 Instead of $CDCl_3$, $C_2H_4Cl_2$ was used as the solvent for initial 30 min of this reaction because of the insolubility of $NaBAr^F_4$ for $CDCl_3$.
- 12 The structure of **4a** was determined by a preliminary singlecrystal X-ray diffraction study.
- 13 Preliminary DFT calculation indicated that the free energy barrier of activation for the β -carbon elimination is *ca*.

19 kcal mol⁻¹ (see ESI[†]), and therefore we consider that the formation of the η^2 -alkyne complex from **A** is acceptable in the present reaction conditions, while we have not obtained satisfactory calculation results to explain the C–N bond oxidative addition yet.

- 14 M. Etienne, R. Mathieu and B. Donnadieu, J. Am. Chem. Soc., 1997, **119**, 3218–3228.
- 15 We appreciate one of the reviewers' helpful suggestions.
- 16 The ¹H NMR analysis of the crude reaction mixture indicated that complexes 7 and 8 were formed in the ratio of 1.4 : 1.
- 17 When isolated 7 and 8 are dissolved in $C_2H_4Cl_2$ and kept at 50 °C overnight, no isomerization was observed for both complexes.