## Dalton Transactions

## PAPER



**Cite this:** *Dalton Trans.*, 2015, **44**, 20898

Received 8th October 2015, Accepted 6th November 2015 DOI: 10.1039/c5dt03941j

www.rsc.org/dalton

### Introduction

Elemental silicon and germanium and their compounds with oxygen and chalcogens are key technological materials, with applications in electronics, glasses, ceramics and optics.<sup>1-3</sup> We are currently developing routes to electrochemically deposit the elements and their binary and ternary alloys from both organic solvents and supercritical fluids.<sup>4-6</sup> In the search for silicon and germanium reagents with appropriate chemical stabilities and solubilities in these media for electrochemical studies, we have explored a variety of coordination complexes, to establish how the properties may be tuned by incorporating various ligands. Most coordination chemistry of germanium(II), germanium(IV) and silicon(IV) involves neutral adducts of the di- or tetra-halides,<sup>7</sup> and cationic complexes are rather rare, while Si(II) halide complexes are limited to

# Cationic aza-macrocyclic complexes of germanium(II) and silicon(IV)†

Matthew Everett, Andrew Jolleys, William Levason, Mark E. Light, David Pugh and Gillian Reid\*

[GeCl<sub>2</sub>(dioxane)] reacts with the neutral aza-macrocyclic ligands L, L = Me<sub>3</sub>tacn (1,4,7-trimethyl-1,4,7-triazacyclononane), Me<sub>4</sub>cyclen (1,4,7,10-tetramethyl-1,4,7,10-tetraazacyclododecane) or Me<sub>4</sub>cyclam (1,4,8,11-tetramethyl-1,4,8,11-tetraazacyclotetradecane) and two mol. equiv. of Me<sub>3</sub>SiO<sub>3</sub>SCF<sub>3</sub> in thf solution to yield the unusual and hydrolytically very sensitive [Ge(L)][O<sub>3</sub>SCF<sub>3</sub>]<sub>2</sub> as white solids in moderate yield. Using shorter reaction times [Ge(Me<sub>3</sub>tacn)]Cl<sub>2</sub> and [Ge(Me<sub>3</sub>tacn)]Cl[O<sub>3</sub>SCF<sub>3</sub>] were also isolated; the preparation of [Ge(Me<sub>4</sub>cyclen)][GeCl<sub>3</sub>]<sub>2</sub> is also described. The structures of the Me<sub>3</sub>tacn complexes show  $\kappa^3$ -coordination of the macrocycle, with the anions interacting only weakly to produce very distorted fiveor six-coordination at germanium. In contrast, the structure of [Ge(Me<sub>4</sub>cyclen)][O<sub>3</sub>SCF<sub>3</sub>]<sub>2</sub> shows no anion interactions, and a distorted square planar geometry at germanium from coordination to the tetra-aza macrocycle. Crystal structures of the Si(w) complexes, [SiCl<sub>3</sub>(Me<sub>3</sub>tacn)]Y (Y = O<sub>3</sub>SCF<sub>3</sub>, BAr<sup>F</sup>; [B{3,5-(CF<sub>3</sub>)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>)<sub>4</sub>]) and [SiHCl<sub>2</sub>(Me<sub>3</sub>tacn)][BAr<sup>F</sup>], obtained from reaction of SiCl<sub>4</sub> or SiHCl<sub>3</sub> with Me<sub>3</sub>tacn, followed by addition of either Me<sub>3</sub>SiO<sub>3</sub>SCF<sub>3</sub> or Na[BAr<sup>F</sup>], contain distorted octahedral cations, with *facial*  $\kappa^3$ coordinated Me<sub>3</sub>tacn. The open-chain triamine, Me<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>N(Me)CH<sub>2</sub>CH<sub>2</sub>NMe<sub>2</sub> (pmdta), forms [SiCl<sub>3</sub>(pmdta)][BAr<sup>F</sup>] and [SiBr<sub>3</sub>(pmdta)][BAr<sup>F</sup>]</sup> under similar conditions, containing *mer*-octahedral cations.

N-heterocyclic carbenes.<sup>8</sup> Aza-macrocyclic complexes reported include  $[GeF_3(Me_3tacn)]_2[GeF_6]$  (Me<sub>3</sub>tacn = 1,4,7-trimethyl-1,4,7-triazacyclononane),<sup>9</sup> [(GeF\_4)\_2( $\kappa^2\kappa'^2$ -Me<sub>4</sub>cyclam)] (Me<sub>4</sub>cyclam = 1,4,8,11-tetramethyl-1,4,8,11-tetraazacyclotetradecane),<sup>9</sup> [GeCl<sub>3</sub>-(Me\_3tacn)]\_2(H\_3O)\_2Cl\_3,<sup>10</sup> [SiF\_3(Me\_3tacn)][SiF\_5],<sup>11</sup> [Ge(Me\_3tacn)]Br-[GeBr\_3],<sup>12</sup> and [Ge(Me\_4cyclam)][GeCl\_3]\_2.<sup>12</sup>

Here we report the synthesis of several new Ge(II) monoand dications and Si(IV) monocations based upon neutral triaza- and tetra-aza macrocyclic ligands with a variety of anions. Single crystal X-ray structural studies on representative examples are described and compared.

## Experimental

SiCl<sub>4</sub>, SiHCl<sub>3</sub>, Me<sub>3</sub>SiO<sub>3</sub>SCF<sub>3</sub>, [GeCl<sub>2</sub>(dioxane)] and the N-donor ligands were obtained from Sigma Aldrich, except for Me<sub>3</sub>tacn which was prepared using the literature route.<sup>13</sup> Na[BAr<sup>F</sup>] ([BAr<sup>F</sup>]<sup>-</sup> = [B{3,5-(CF<sub>3</sub>)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>}]<sup>-</sup>) was synthesised by a modification of Brookhart's procedure.<sup>14</sup> Me<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>N(Me)-CH<sub>2</sub>CH<sub>2</sub>NMe<sub>2</sub> (pmdta) was distilled from CaH<sub>2</sub>. All experiments were performed under strictly anhydrous conditions using glove-boxes and Schlenk techniques. CH<sub>2</sub>Cl<sub>2</sub> was dried by distillation from CaH<sub>2</sub>, toluene was distilled from sodium, hexane was distilled from Na/K alloy and thf was distilled from

View Article Online

School of Chemistry, University of Southampton, Southampton, SO17 1BJ, UK. E-mail: G.Reid@soton.ac.uk: Tel: +44 (0)23 8059 3609

 $<sup>\</sup>dagger$  Electronic supplementary information (ESI) available: CCDC 1430051 ([Ge(Me\_{3}tacn)][O\_{3}SCF\_{3}]\_{2}\cdotMeCN), 1430052 ([Ge(Me\_{4}cyclen)][O\_{3}SCF\_{3}]\_{2}\cdot0.8MeCN), 1430053 ([Ge(Me\_{3}tacn)]Cl[O\_{3}SCF\_{3}]\_{2}), 1430054 ([Ge(Me\_{3}tacn)]Cl\_{2}\cdotMeCN), 14300555 ([SiCl\_{3}(Me\_{3}tacn)][O\_{3}SCF\_{3}]\_{2}), 1430056 ([SiCl\_{3}(Me\_{3}tacn)]Cl\_{2}\cdotMeCN), 16300555 ([SiCl\_{3}(Me\_{3}tacn)]Cl\_{2}\cdotMeCN), 1630056 ([SiCl\_{3}(Me\_{3}tacn)]Cl\_{2}\cdotMeCN), 1630057 ([SiCl\_{3}(Me\_{3}tacn)]Cl\_{2}\cdotMeCN), 1630056 ([SiCl\_{3}(Me\_{3}tacn)]Cl\_{2}\cdotMeCN), 1630056 ([SiCl\_{3}(Me\_{3}tacn)]Cl\_{2}\cdotMeCN), 1630056 ([SiCl\_{3}(Me\_{3}tacn)]Cl\_{2}\cdotMeCN), 1630056 ([SiCl\_{3}(Me\_{3}tacn)]Cl\_{2}\cdotMeCN), 1630057 ([SiCl\_{3}(Me\_{3}tacn)]Cl\_{2}\cdotMeCN), 1630056 ([SiCl\_{3}(Me\_{3}tacn)]Cl\_{2}\cdotMeCN), 1630057 ([SiCl\_{3}(Me\_{3}tacn)]Cl\_{2}\cdotMeCN), 1630056 ([SiCl\_{3}(Me\_{3}tacn)]Cl\_{2}\cdotMeCN), 1630057 ([SiCl\_{3}(Me\_{3

Na/benzophenone ketyl. IR spectra were recorded as Nujol mulls between CsI plates using a Perkin Elmer Spectrum 100 spectrometer over the range 4000–200 cm<sup>-1</sup>. <sup>1</sup>H and <sup>19</sup>F NMR spectra were recorded using a Bruker DPX-400 spectrometer and referenced to the residual solvent resonance and external CFCl<sub>3</sub>, respectively. Microanalytical measurements were performed by London Metropolitan University.

#### [Ge(Me<sub>4</sub>cyclen)][O<sub>3</sub>SCF<sub>3</sub>]<sub>2</sub>

[GeCl<sub>2</sub>(dioxane)] (0.059 g, 0.26 mmol) was dissolved in thf (10 mL) and a solution of  $Me_3SiO_3SCF_3$  (0.112 g, 0.50 mmol) in thf (5 mL) was added with stirring, giving a clear, colourless solution. After 10 min, a solution of  $Me_4$ cyclen (0.057 g, 0.25 mmol) in thf (5 mL) was added, causing the formation of a white precipitate. After stirring for a further 1 h, the product was collected by filtration and dried *in vacuo*. Yield: 0.120 g (80%). Anal. calc. for  $C_{14}H_{28}F_6GeN_4O_6S_2$  (599.1): C, 28.08; H, 4.71; N, 9.35. Found: C, 28.19; H, 4.61; N, 9.26%. <sup>1</sup>H NMR (CD<sub>3</sub>CN, 298 K): 2.78 (s, [12H], NCH<sub>3</sub>), 3.20–3.40 (m, [16H], NCH<sub>2</sub>). <sup>19</sup>F{<sup>1</sup>H} NMR (CD<sub>3</sub>CN, 298 K): -79.4 (O<sub>3</sub>SCF<sub>3</sub>). IR (Nujol/cm<sup>-1</sup>): 474w, 517m, 573m, 638s, 749m, 793w, 919w, 960w, 1030s, 1068w, 1155s, 1225s, 1260s.

Crystals of  $[Ge(Me_4cyclen)][O_3SCF_3]_2$ ·CH<sub>3</sub>CN suitable for X-ray diffraction were obtained by layering an acetonitrile solution with diethyl ether.

#### [Ge(Me<sub>4</sub>cyclam)][O<sub>3</sub>SCF<sub>3</sub>]<sub>2</sub>

[GeCl<sub>2</sub>(dioxane)] (0.058 g, 0.25 mmol) was dissolved in thf (10 mL), and a solution of Me<sub>3</sub>SiO<sub>3</sub>SCF<sub>3</sub> (0.115 g, 0.52 mmol) in thf (5 mL) was added with stirring, giving a colourless solution. After 10 min a solution of Me<sub>4</sub>cyclam (0.065 g, 0.25 mmol) in thf (5 mL) was added, causing the formation of a white, microcrystalline precipitate. After 1 h, the product was collected by filtration, washed with diethyl ether and dried in vacuo. Yield: 0.092 g (59%). Anal. calc. for C<sub>16</sub>H<sub>32</sub>F<sub>6</sub>GeN<sub>4</sub>O<sub>6</sub>S<sub>2</sub> (627.2): C, 30.65; H, 5.14; N, 8.93. Found: C, 30.81; H, 5.17; N, 8.86%. <sup>1</sup>H NMR (CD<sub>3</sub>CN, 298 K): 1.96 (s, [4H], NCH<sub>2</sub>CH<sub>2</sub>), 2.56 (br s, [12H], NCH<sub>3</sub>), 2.94-2.99 (v br, [16H], overlapping NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub> and NCH<sub>2</sub>CH<sub>2</sub>N).  ${}^{19}F{}^{1}H{}^{1}$ NMR (CD<sub>3</sub>CN, 298 K): -79.4 (O<sub>3</sub>SCF<sub>3</sub>). <sup>1</sup>H NMR (DMF-d<sub>7</sub>, 298 K): 2.09 (quintet, [4H],  ${}^{3}J_{HH} = 6.3$  Hz, NCH<sub>2</sub>CH<sub>2</sub>), 2.68 (br s, [12H], NCH<sub>3</sub>), 3.08 (v br, [8H], NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>) 3.21 (br s, [8H], NCH<sub>2</sub>CH<sub>2</sub>N).  ${}^{19}$ F{ ${}^{1}$ H} NMR (DMF-d<sub>7</sub>, 298 K): -79.38 (O<sub>3</sub>SCF<sub>3</sub>). IR (Nujol/cm<sup>-1</sup>): 517m, 574m, 592w, 639s, 743w, 757w, 802w, 1011m, 1034s, 1160s, 1226m, 1256s, 1278s, 1354m.

#### [Ge(Me<sub>4</sub>cyclen)][GeCl<sub>3</sub>]<sub>2</sub>

[GeCl<sub>2</sub>(dioxane)] (0.104 g, 0.45 mmol) was suspended in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) and a solution of Me<sub>4</sub>cyclen (0.035 g, 0.15 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) was added with stirring, giving a dense white suspension. After stirring at room temperature for 2 h, the product was collected by filtration, washed with Et<sub>2</sub>O and dried *in vacuo*. Yield: 0.048 g (48%). Anal. calc. for:  $C_{12}H_{28}Cl_6Ge_3N_4$  (658.9): C, 21.87; H, 4.28; N, 8.50. Found: C, 22.02; H, 4.19; N, 8.41%. <sup>1</sup>H NMR (CD<sub>3</sub>CN, 298 K): 2.78 (s,

[12H], NCH<sub>3</sub>), 3.20–3.38 (m, [16H], NCH<sub>2</sub>). IR (Nujol/cm<sup>-1</sup>): 277s, vbr [GeCl<sub>3</sub>]<sup>-</sup>, 317s, 413w, 472m, 547m, 737s, 747s, 791m, 917s, 945m, 958s, 1014s, 1024s, 1051s, 1064s, 1151m, 1262s, 1275s, 1298s.

#### [Ge(Me<sub>3</sub>tacn)][O<sub>3</sub>SCF<sub>3</sub>]<sub>2</sub>

[GeCl<sub>2</sub>(dioxane)] (0.059 g, 0.26 mmol) was suspended in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) and a solution of Me<sub>3</sub>SiO<sub>3</sub>SCF<sub>3</sub> (0.112 g, 0.50 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) was added with stirring, giving a colourless solution. After 10 min Me<sub>3</sub>tacn (0.044 g, 0.26 mmol) was added, causing the formation of a white precipitate. After stirring for approximately 1 h, the product was collected by filtration and dried *in vacuo*. Yield: 0.085 g (62%). Anal. calc. for C<sub>11</sub>H<sub>21</sub>F<sub>6</sub>GeN<sub>3</sub>O<sub>6</sub>S<sub>2</sub> (542.0): C, 24.38; H, 3.91; N, 7.75. Found: C, 24.51; H, 3.74; N, 7.84%. <sup>1</sup>H NMR (CD<sub>3</sub>CN, 298 K): 3.02 (s, [9H], NCH<sub>3</sub>), 3.34–3.54 (m, [12H], NCH<sub>2</sub>). <sup>19</sup>F{<sup>1</sup>H} NMR (CD<sub>3</sub>CN, 298 K): -79.4 (O<sub>3</sub>SCF<sub>3</sub>). IR (Nujol/cm<sup>-1</sup>): 420m, 451w, 516m, 573m, 639s, 739m, 785m, 898m, 982w, 994m, 1030s, 1049m, 1138m, 1167s, 1226s, 1259s.

Crystals of [Ge(Me<sub>3</sub>tacn)][O<sub>3</sub>SCF<sub>3</sub>]<sub>2</sub>·CH<sub>3</sub>CN suitable for X-ray diffraction were obtained by layering an acetonitrile solution with diethyl ether.

#### [Ge(Me<sub>3</sub>tacn)]Cl[O<sub>3</sub>SCF<sub>3</sub>]

In a similar reaction using a 1:1 molar ratio of  $[GeCl_2(diox$ ane)] and Me<sub>3</sub>tacn, conducted in MeCN solution, the mixture was stirred at room temperature for *ca.* 3 h following addition of Me<sub>3</sub>SiO<sub>3</sub>SCF<sub>3</sub>, then concentrated *in vacuo*. This did not cause any precipitation, and the solution was layered with Et<sub>2</sub>O and stored in the freezer. Small rod-shaped crystals formed, which were found to be  $[Ge(Me_3tacn)]Cl[O_3SCF_3]$  as identified by an X-ray crystal structure determination. Anal. calc. for C<sub>10</sub>H<sub>21</sub>ClF<sub>3</sub>GeN<sub>3</sub>O<sub>3</sub>S (428.4): C, 28.04; H, 4.94; N, 9.81. Found: C, 28.76; H, 4.85; N, 9.77%. <sup>1</sup>H NMR (298 K, CD<sub>2</sub>Cl<sub>2</sub>): 2.94 (s, [9H], NCH<sub>3</sub>), 3.21–3.47 (m, [12H], CH<sub>2</sub>). (A second minor species is also evident in the NMR spectrum). <sup>19</sup>F{<sup>1</sup>H} NMR (298 K, CD<sub>2</sub>Cl<sub>2</sub>): –79.4 (O<sub>3</sub>SCF<sub>3</sub>).

#### [Ge(Me<sub>3</sub>tacn)]Cl<sub>2</sub>

In a further reaction conducted in MeCN solution, the mixture was stirred at room temperature for *ca.* 15 min following addition of  $Me_3SiO_3SCF_3$ , then concentrated *in vacuo*. This caused the rapid precipitation of a white solid, which redissolved on warming. Storage of this solution in the freezer furnished colourless rod-shaped crystals which were found by X-ray crystallographic analysis to be [Ge(Me\_3tacn)]-Cl<sub>2</sub>·MeCN.

#### [SiCl<sub>3</sub>(Me<sub>3</sub>tacn)][O<sub>3</sub>SCF<sub>3</sub>]

SiCl<sub>4</sub> (0.170 g, 1.0 mmol) and Me<sub>3</sub>SiO<sub>3</sub>SCF<sub>3</sub> (0.222 g, 1.0 mmol) were dissolved in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) and stirred for 5 min. A solution of Me<sub>3</sub>tacn (0.171 g, 1.0 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was added and the reaction was stirred for 16 h. After this time, a white solid had formed. This was collected by filtration and dried *in vacuo*. Yield: 0.204 g (45%). Anal. calc.

for  $C_{10}H_{21}Cl_3F_3N_3O_3SSi$  (454.8): C, 26.40; H, 4.66; N, 9.24. Found: C, 26.61; H, 4.73; N, 9.25%. <sup>1</sup>H NMR (295 K, CD<sub>3</sub>CN): 3.54–3.66 (m, [6H], NCH<sub>2</sub>), 3.31–3.41 (m, [6H], NCH<sub>2</sub>), 3.17 (s, [9H], NCH<sub>3</sub>). <sup>13</sup>C{<sup>1</sup>H} NMR (295 K, CD<sub>3</sub>CN): 122.20 (q, <sup>1</sup> $J_{C-F} =$ 320 Hz, CF<sub>3</sub>), 54.86 (CH<sub>2</sub>), 53.49 (CH<sub>3</sub>). <sup>19</sup>F{<sup>1</sup>H} NMR (295 K, CD<sub>3</sub>CN): -78.7 (O<sub>3</sub>SCF<sub>3</sub>). IR (Nujol, cm<sup>-1</sup>): 231s, 428s (SiCl), 460s (SiCl), 497m, 517m, 574m, 601m, 637s, 754m, 899m, 967m, 998m, 1029s, 1053m, 1155s, 1225m.

Crystals were obtained by layering a concentrated  $\mathrm{CH}_2\mathrm{Cl}_2$  solution with hexane.

#### [SiHCl<sub>2</sub>(Me<sub>3</sub>tacn)][BAr<sup>F</sup>]

SiHCl<sub>3</sub> (0.068 g, 0.50 mmol) and NaBAr<sup>F</sup> (0.443 g, 0.50 mmol) were dissolved in toluene (10 mL) and stirred for 5 min. A solution of Me<sub>3</sub>tacn (0.085 g, 0.50 mmol) in toluene (5 mL) was added then the reaction was stirred for 4 h. After this time, volatiles were removed in vacuo and the solid extracted into CH<sub>2</sub>Cl<sub>2</sub> (5 mL), filtered and hexane (30 mL) added to precipitate a white solid. Yield: 0.421 g (74%). Anal. calc. for C41H34BCl2F24N3Si (1134.12): C, 43.38; H, 3.02; N, 3.70. Found: C, 43.45; H, 3.11; N, 3.80%. <sup>1</sup>H NMR (295 K, CD<sub>2</sub>Cl<sub>2</sub>): 7.74 (s, [8H], BAr<sup>F</sup> H2/6), 7.59 (s, [4H], BAr<sup>F</sup> H4), 4.78 (s, [1H], Si-H), 3.54-3.70 (m, [4H], NCH<sub>2</sub>), 3.26 (s, [3H], NCH<sub>3</sub>), 3.21 (s, [4H], NCH<sub>2</sub>), 3.12-3.17 (m, [4H], NCH<sub>2</sub>), 2.99 (s, [6H], NCH<sub>3</sub>). <sup>13</sup>C{<sup>1</sup>H} NMR (295 K, CD<sub>2</sub>Cl<sub>2</sub>): 162.33 (q,  ${}^{1}J_{C-B}$  = 49.8 Hz, BAr<sup>1</sup> C1), 135.42 (CH, BAr<sup>F</sup> C2/6), 129.48 (C, q,  ${}^{1}J_{C-F} = 3.3$  Hz, BAr<sup>F</sup> C3/5), 125.22 (C, q,  ${}^{1}J_{C-F}$  = 272 Hz, CF<sub>3</sub>), 118.11 (CH, BAr<sup>F</sup> C4), 54.72, 53.77, 51.41, 50.45 (NCH<sub>2</sub> and NCH<sub>3</sub>). IR (Nujol, cm<sup>-1</sup>): 449 m (SiCl), 480 m (SiCl), 683s, 744m, 838m, 888m, 899m, 1005m, 1056m, 1086m, 1114s, 1163s, 1280s, 1289s, 1356s, 1612w, 2137w (SiH).

#### [SiCl<sub>3</sub>(Me<sub>3</sub>tacn)][BAr<sup>F</sup>]

Made similarly using SiCl<sub>4</sub> (0.085 g, 0.50 mmol) in place of SiHCl<sub>3</sub>. Yield: 0.466 g (82%). Anal. calc. for  $C_{41}H_{33}BCl_3F_{24}N_3Si$  (1168.9): C, 42.10; H, 2.85; N, 3.59. Found: C, 41.85; H, 3.08; N, 3.74%.

A concentrated  $CH_2Cl_2$  solution of this complex layered with hexane deposited a few colourless crystals which were identified from an X-ray crystallographic study as [SiCl<sub>3</sub>(Me<sub>3</sub>tacn)]Cl. Attempts to prepare the latter directly from SiCl<sub>4</sub> and Me<sub>3</sub>tacn in  $CH_2Cl_2$  gave a pale yellow solid which was very poorly soluble in chlorocarbons and which was not analytically pure.

#### [SiCl<sub>3</sub>(pmdta)][BAr<sup>F</sup>]

SiCl<sub>4</sub> (0.085 g, 0.50 mmol) and Na[BAr<sup>F</sup>] (0.443 g, 0.50 mmol) were dissolved in toluene (10 mL) and stirred for 5 min. A solution of pmdta (0.087 g, 0.50 mmol) in toluene (5 mL) was added, then the reaction was stirred for 4 h. After this time, volatiles were removed *in vacuo* and the solid extracted into CH<sub>2</sub>Cl<sub>2</sub> (5 mL), filtered and hexane (30 mL) added to precipitate a white solid. Yield: 0.447 g (76%). Anal. calc. for C<sub>41</sub>H<sub>35</sub>BCl<sub>3</sub>F<sub>24</sub>N<sub>3</sub>Si (1170.58): C, 42.03; H, 3.01; N, 3.59. Found: C, 41.88; H, 3.12; N, 3.71%. <sup>1</sup>H NMR (295 K, CD<sub>2</sub>Cl<sub>2</sub>): 7.72 (s, [8H], BAr<sup>F</sup> H2/6), 7.57 (s, [4H], BAr<sup>F</sup> H4), 2.76–2.98 (br s, [4H],

NCH<sub>2</sub>), 2.65 (v br s, [16H], N(CH<sub>3</sub>)<sub>2</sub> and NCH<sub>2</sub>), 2.35 (s, [3H], NCH<sub>3</sub>) ppm. <sup>13</sup>C{<sup>1</sup>H} NMR (295 K, CD<sub>2</sub>Cl<sub>2</sub>): 162.37 (q, <sup>1</sup> $J_{C-B}$  = 49.7 Hz, BAr<sup>F</sup> C1), 135.41 (CH, BAr<sup>F</sup> C2/6), 129.45 (C, q, <sup>1</sup> $J_{C-F}$  = 3.3 Hz, BAr<sup>F</sup> C3/5), 125.22 (C, q, <sup>1</sup> $J_{C-F}$  = 272 Hz, CF<sub>3</sub>), 118.10 (CH, BAr<sup>F</sup> C4), 57.91, 57.16 (CH<sub>2</sub>), 45.08 (N(CH<sub>3</sub>)<sub>2</sub>), 44.51 (NCH<sub>3</sub>). IR (Nujol cm<sup>-1</sup>): 448w, 506w (SiCl), 522w (SiCl), 682m, 713m, 889w, 1112s, 1143s, 1358s, 1367s.

Crystals were obtained by layering a concentrated  $\mathrm{CH}_2\mathrm{Cl}_2$  solution with hexane.

#### [SiBr<sub>3</sub>(pmdta)][BAr<sup>F</sup>]

SiBr<sub>4</sub> (0.174 g, 0.50 mmol) and Na[BAr<sup>F</sup>] (0.443 g, 0.50 mmol) were dissolved in toluene (10 mL) and stirred for 5 min. A solution of pmdta (0.087 g, 0.50 mmol) in toluene (5 mL) was added then the reaction was stirred for 4 h. After this time, volatiles were removed *in vacuo* and the solid extracted into CH<sub>2</sub>Cl<sub>2</sub> (5 mL), filtered and hexane (30 mL) added to precipitate a white solid. Yield: 0.291 g (45%). Anal. calc. for C<sub>41</sub>H<sub>35</sub>BBr<sub>3</sub>F<sub>24</sub>N<sub>3</sub>Si (1303.93): C, 37.73; H, 2.71; N, 3.22. Found: C, 37.57; H, 2.80; N, 3.26%. <sup>1</sup>H NMR (CD<sub>2</sub>Cl<sub>2</sub>): 7.72 (s, [8H], BAr<sup>F</sup> H2/6), 7.57 (s, [4H], BAr<sup>F</sup> H4), 2.76–2.98 (br s, [4H], NCH<sub>2</sub>), 2.65 (br s, [16H], N(CH<sub>3</sub>)<sub>2</sub> and NCH<sub>2</sub>), 2.35 (s, [3H], NCH<sub>3</sub>). IR (Nujol, cm<sup>-1</sup>): 363 m (SiBr), 471w, 521w, 584w, 682s, 712s, 838m, 887m, 898m, 925w, 955w, 1112s, 1144s, 1279s, 1358s.

#### X-ray crystallography

Crystals were obtained as described above. Details of the crystallographic data collection and refinement are in Table 1. Diffractometer: Rigaku AFC12 goniometer equipped with an enhanced sensitivity (HG) Saturn724+ detector mounted at the window of an FR-E+ SuperBright molybdenum rotating anode generator ( $\lambda_1 = 0.71073$  Å) with VHF Varimax optics (70 or 100 μm focus). Cell determination, data collection, data reduction, cell refinement and absorption correction: CrystalClear-SM Expert 2.0 r7.4.<sup>15a</sup> Structure solution and refinement were carried out using WinGX or Olex2 and software packages within.<sup>15b-d</sup> No positional disorder was observed in complexes of [BAr<sup>F</sup>]<sup>-</sup>, despite this being a common issue with weaklycoordinating anions containing CF<sub>3</sub> groups, especially [BAr<sup>F</sup>]<sup>-</sup>.<sup>16</sup> [Ge(Me<sub>3</sub>tacn)]Cl<sub>2</sub>·MeCN crystallised as an inversion twin with a BASF of 0.21. Unusually large Z values were observed for [Ge(Me4cyclen)][O3SCF3]2.0.8MeCN and [SiCl3-(Me<sub>3</sub>tacn)]Cl (10 and 24, respectively) which is explained by the presence of multiple cation units with very similar, but not identical, metrical parameters in the asymmetric unit. H atoms attached to C atoms were placed in geometrically assigned positions, with C–H distances of 0.95 Å (CH), 0.98 Å  $(CH_3)$  or 0.99 Å  $(CH_2)$  and refined using a riding model, with  $U_{\rm iso}({\rm H}) = 1.2U_{\rm eq}({\rm C})$  (CH, CH<sub>2</sub>) or  $1.5U_{\rm eq}({\rm C})$  (CH<sub>3</sub>). Si-H and N-H protons were located in the Fourier difference map and allowed to refine freely. enCIFer was used to prepare CIFs for publication.15e

[Ge(Me3tacn)]-

C11H24Cl2GeN4

Orthorhombic

Cl<sub>2</sub>·MeCN

 $Pna2_{1}(33)$ 

14.399(5)

10.139(5)

21.491(5)

90

355.83

Table 1         Selected X-ray crystallographic data <sup>a</sup>			
Compound	[Ge(Me₃tacn)]- [O₃SCF₃]₂·MeCN		
Formula	C13H24F6GeN4O6S2		
$M/g \text{ mol}^{-1}$	583.07		
Crystal system	Monoclinic		
Space group (No.)	$P2_{1}/n$ (14)		
a/Å	13.541(5)		
b/Å	8.543(5)		
c/Å	19.620(5)		
$\alpha/^{\circ}$	90		
$\beta  ^{\circ}$	97.495(5)		
γ/°	90		
$U/Å^3$	2250.3(17)		
Z	4		
$\mu$ (Mo-K $\alpha$ )/mm <sup>-1</sup>	1.634		
F(000)	1184		
Total reflections	10 515		
Unique reflections	5123		
	0.041		
Goodness-of-fit on $F^2$	1 036		
$R_{i}^{b} \left[ I > 2\sigma(I) \right]$	0.037		
$R_1$ [ $R_2$ (all data)	0.050		
$wR_{o}^{b}[I \geq 2\sigma(I)]$	0.082		
$wR_{2}$ (all data)	0.088		
Compound	[SiCl <sub>3</sub> (Me <sub>3</sub> ta		
Formula	C10H21Cl2F2		
$M/g \text{ mol}^{-1}$	454.80		
Crystal system	Triclinic		
Space group (no.)	$P\bar{1}(2)$		
a/Å	6.997(2)		
b/Å	10.820(4)		
c/Å	11.974(4)		
$\alpha/^{\circ}$	99.110(7)		
$\beta/\circ$	93.245(5)		
v/°	92.108(8)		
$I I / Å^3$	892.7(5)		
7.	092.7(0)		
$u(Mo-K\alpha)/mm^{-1}$	0 743		
F(000)	468		
Total reflections	7788		
Unique reflections	//00		
	4019		
$\Lambda_{\text{int}}$	0.019		
$D^{b}[I > 2\sigma(I)]$	1.042		
$\pi_1  I_0 \ge 20(I_0) $	0.032		

β/°	97.495(5)	90	101.705(7)	90
$\gamma/^{\circ}$	90	90	90	90
$U/Å^3$	2250.3(17)	6486.1(4)	1593.2(3)	3138(2)
Ζ	4	10(Z' = 1.25)	3	8
$\mu$ (Mo-K $\alpha$ )/mm <sup>-1</sup>	1.634	1.425	2.267	2.284
F(000)	1184	3236	872	1472
Total reflections	10 515	34 358	8862	11 548
Unique reflections	5123	7099	3647	5491
R <sub>int</sub>	0.041	0.042	0.074	0.0823
Goodness-of-fit on $F^2$	1.036	1.030	0.987	1.025
$R_1^{b} \left[ I_0 > 2\sigma(I_0) \right]$	0.037	0.054	0.047	0.061
$R_1$ (all data)	0.050	0.071	0.060	0.108
$wR_2^{b} [I_o > 2\sigma(I_o)]$	0.082	0.131	0.108	0.103
$wR_2$ (all data)	0.088	0.142	0.116	0.119
Compound	[SiCl <sub>3</sub> (Me <sub>3</sub> tacr	h)][O <sub>3</sub> SCF <sub>3</sub> ]	[SiCl <sub>3</sub> (Me <sub>3</sub> tacn)]Cl	[SiCl <sub>3</sub> (pmdta)][BAr <sup>F</sup> ]
Formula	$C_{10}H_{21}Cl_3F_3N_3$	<sub>3</sub> O <sub>3</sub> SSi	$C_9H_{21}Cl_4N_3Si^2/_3(CH_2Cl_2)$	C41H35BCl3F24N3Si
$M/g \text{ mol}^{-1}$	454.80		397.80	1170.97
Crystal system	Triclinic		Monoclinic	Orthorhombic
Space group (no.)	$P\overline{1}(2)$		C2/c (15)	<i>Pbca</i> (61)
a/Å	6.997(2)		34.475(8)	17.728(5)
b/Å	$10.820(4) \\ 11.974(4) \\ 99.110(7) \\ 93.245(5) \\ 92.108(8)$		$26.707(6) \\11.820(3) \\90 \\106.684(5) \\90$	19.613(6)
c/Å				26.662(8)
$\alpha / ^{\circ}$				90
$\beta / ^{\circ}$				90 90
$\gamma/^{\circ}$				
$U/\text{\AA}^3$	892.7(5)		10 425(4)	9270(5)
Ζ	2		24(Z'=3)	8
$\mu$ (Mo-K $\alpha$ )/mm <sup>-1</sup>	0.743		0.946	0.357
F(000)	468		4944	4704
Total reflections	7788		46 773	17 463
Unique reflections	4019		10659	8124
R <sub>int</sub>	0.019		0.114	0.070
Goodness-of-fit on $F^2$	1.042		1.107	1.176
$R_1^{b} \left[ I_0 > 2\sigma(I_0) \right]$	0.032		0.094	0.097
$R_1$ (all data)	0.039		0.121	0.147
$wR_2^{b} [I_o > 2\sigma(I_o)]$	0.069		0.227	0.137
$wR_2$ (all data)	0.072		0.247	0.155
<sup><i>a</i></sup> Common items: $T = 100$ H	K; wavelength (Mo-Kα) = 0.7	71073 Å; $\theta(\max) = 27.5^{\circ}$ .	$R_1 = \sum   F_{\sigma}  -  F_{c}   / \sum  F_{o} ; wR_2 = \sum w(A_{\sigma})$	$F_{\rm o}^{2} - F_{\rm c}^{2})^{2} / \sum w F_{\rm o}^{2} ]^{1/2}.$

[Ge(Me4cyclen)]-

631.96

Tetragonal

P4cc (103)

17.1253(3)

17.1253(3)

90

22.1160(10)

[O<sub>3</sub>SCF<sub>3</sub>]<sub>2</sub>·0.8MeCN

 $C_{15.6}H_{30.4}F_6GeN_{4.8}O_6S_2$ 

[Ge(Me3tacn)]-

C10H21ClF3GeN3O3S

Cl[O<sub>3</sub>SCF<sub>3</sub>]

Monoclinic

13.9764(15)

14.2804(16)

 $P2_1/c$  (14)

8.1521(9)

90

438.40

#### **Results and discussion**

#### Germanium(II) complexes

We have previously reported<sup>12</sup> that reaction of GeBr<sub>2</sub> with Me3tacn in anhydrous MeCN solution gives colourless [Ge-(Me<sub>3</sub>tacn)]Br[GeBr<sub>3</sub>], containing unusual discrete three-coordinate  $[Ge(Me_3tacn)]^{2+}$  dications (Ge-N = 2.124(3)-2.156(3) Å), with Br<sup>-</sup> and [GeBr<sub>3</sub>]<sup>-</sup> anions providing charge balance. We find that reaction of  $[GeCl_2(dioxane)]$  with Me<sub>3</sub>tacn in anhydrous CH2Cl2 followed by addition of Me3SiO3SCF3 colourless powder,  $[Ge(Me_3tacn)][O_3SCF_3]_2,$ gave а subsequently obtained as colourless crystals of [Ge(Me<sub>3</sub>tacn)]-

[O<sub>3</sub>SCF<sub>3</sub>]<sub>2</sub>·MeCN by recrystallisation from MeCN/Et<sub>2</sub>O. The structure (Fig. 1) also reveals a pyramidal GeN<sub>3</sub> unit, however, in this species there are also weak directional Ge---O interactions from one oxygen in each triflate anion at 2.850(2) and 3.179(2) Å, well within the sum of vdW radii for O and Ge (3.79 Å).17

While there is no evidence for 2:1 Me3tacn: Ge species (presumably due to steric clashing of the Me groups on the relatively small Ge(II) centre), the anions present in the products proved to be very sensitive to the reaction conditions; if the reaction of [GeCl<sub>2</sub>(dioxane)], Me<sub>3</sub>tacn and Me<sub>3</sub>SiO<sub>3</sub>SCF<sub>3</sub> was conducted in MeCN solution as described in the



Fig. 1 The X-ray structure of  $[Ge(Me_3tacn)][O_3SCF_3]_2$ ·MeCN. Thermal ellipsoids are drawn at 50% probability, and hydrogen atoms omitted for clarity. Selected bond lengths (Å) and angles (°): N1-Ge1 = 2.084(2), N2-Ge1 = 2.106(2), N3-Ge1 = 2.089(2), O5…Ge1 = 2.850(2), O3…Ge1 = 3.179(2), N1-Ge1-N3 = 82.73(8), N1-Ge1-N2 = 82.84(8), N3-Ge1-N2 = 83.23(9).

Experimental section, concentrated *in vacuo*, and the solution layered with Et<sub>2</sub>O, the product was isolated as rod-like crystals of [Ge(Me<sub>3</sub>tacn)]Cl[O<sub>3</sub>SCF<sub>3</sub>]. The X-ray structure of this species (Fig. 2) shows the complex forms a weakly associated dimer in the solid state *via* the chloride bridges (Ge…Cl = 3.0254(9), 3.214(1) Å) and with a similar pyramidal 'Ge<sup>II</sup>(Me<sub>3</sub>tacn)' core. These Ge…Cl interactions are substantially within the sum of vdW radii for Ge and Cl (4.11 Å). One  $\kappa^1$ -coordinated triflate completes a distorted six-coordinate environment at each germanium centre.

The same reaction conducted in MeCN solution, but worked up after 15 min and placed in a freezer, deposited a few rod-like crystals that were identified as  $[Ge(Me_3tacn)]$ - $Cl_2$ ·MeCN by X-ray crystallography (Fig. 3). In this case the 'Ge<sup>II</sup>(Me\_3tacn)' core has two long contacts to the chlorides at



**Fig. 2** The X-ray structure of  $[Ge(Me_3tacn)]Cl[O_3SCF_3]$  showing the dimeric unit. Thermal ellipsoids are drawn at 50% probability, and hydrogen atoms omitted for clarity. Selected bond lengths (Å) and angles (°): N1-Ge1 = 2.119(3), N2-Ge1 = 2.139(3), N3-Ge1 = 2.130(2), Cl1...Ge1 = 3.0254(9), O1...Ge1 = 3.381(2), Cl1<sup>i</sup>...Ge1 3.214(1), N1-Ge1-N3 = 82.48(10), N1-Ge1-N2 = 81.63(10), N3-Ge1-N2 = 80.98(10).



Fig. 3 The X-ray structure of  $[Ge(Me_3tacn)]Cl_2$ ·MeCN showing the Ge1centred molecule. The crystallographically independent Ge2-centred molecule is very similar. Thermal ellipsoids are drawn at 50% probability, and hydrogen atoms omitted for clarity Selected bond lengths (Å) and angles (°): N1–Ge1 = 2.124(10), N2–Ge1 = 2.147(10), N3–Ge1 = 2.154(9), Cl1…Ge1 = 3.028(4), Cl2…Ge1 = 3.028(3), N1–Ge1–N3 = 80.9(3), N1– Ge1–N2 = 81.2(4), N3–Ge1–N2 = 80.8(4).

3.028(4) and 3.028(3) Å. The closest intermolecular contact is to a chloride of an adjacent molecule, but the Ge…Cl distance is the same as the sum of vdW radii for Ge and Cl (within experimental error), most likely a consequence of crystal packing.

Comparison of the Ge–N distances in this series of complexes shows only small differences as a function of the anion(s) present, suggesting that the structures are dominated by the 'Ge<sup>II</sup>(Me<sub>3</sub>tacn)' core. All of the Ge–N distances are considerably longer than the sum of the covalent radii (1.85 Å),<sup>18</sup> but well within the sum of the van der Waals radii (3.66 Å).<sup>17</sup>

The spectroscopic data provide very limited information; the IR spectra show the Me<sub>3</sub>tacn and the  $[O_3SCF_3]^-$  (when present), whilst the <sup>1</sup>H NMR spectra show small high frequency shifts corresponding to coordinated Me<sub>3</sub>tacn. The complexes are extremely sensitive to hydrolysis by trace water, readily forming protonated Me<sub>3</sub>tacn.

The reaction of  $[GeCl_2(dioxane)]$  and  $Me_3SiO_3SCF_3$  in thf followed by addition of  $Me_4cyclam$  gave  $[Ge(Me_4cyclam)]$  $[O_3SCF_3]_2$ . Attempts to obtain crystals of this complex were unsuccessful with recrystallisation from MeCN/Et<sub>2</sub>O or CH<sub>2</sub>Cl<sub>2</sub>/ Et<sub>2</sub>O giving  $[Me_4cyclamH_2][O_3SCF_3]_2$  (identified crystallographically). We reported the structure of  $[Ge(Me_4cyclam)]$  $[GeCl_3]_2$  in our preliminary communication,<sup>12</sup> which revealed an essentially coplanar N<sub>4</sub> with the Ge out of the plane by 0.83 Å, and with no significant interaction with the anions. The data also revealed a spread of Ge–N distances (2.151(2)– 2.349(2) Å), suggesting germanium( $\pi$ ) is not a good fit to the relatively large 14-membered macrocycle cavity; this may correlate with the very ready hydrolysis in solution.

Replacing the 14-membered  $Me_4$ cyclam ring by the 12membered  $Me_4$ cyclen (1,4,7,10-tetramethyl-1,4,7,10-tetraazacyclotetradecane) gave colourless  $[Ge(Me_4cyclen)][O_3SCF_3]_2$ . Crystals of  $[Ge(Me_4cyclen)][O_3SCF_3]_2 \cdot 0.8CH_3CN$  suitable for



**Fig. 4** The X-ray structure of the dication present in  $[Ge(Me_4cyclen)]$ - $[O_3SCF_3]_2 \cdot 0.8CH_3CN$  showing the Ge1-containing molecule. Thermal ellipsoids are drawn at 50% probability, and hydrogen atoms omitted for clarity. Selected bond lengths (Å) and angles (°): Ge1-N1 = 2.244(5), Ge1-N2 = 2.168(9), Ge1-N3 = 2.206(6), Ge1-N4 = 2.165(6), N1-Ge1-N3 = 131.8(2), N2-Ge1-N4 = 118.5(3), N2-Ge1-N1 = 78.2(2), N2-Ge1-N3 = 77.6(3), N4-Ge1-N1 = 77.8(2), N4-Ge1-N3 = 78.2(2). Symmetry codes: (i) x, y, 0.5 + z; (ii) -x, 1 - y, 0.5 + z.

X-ray diffraction were obtained by layering an acetonitrile solution of the complex with diethyl ether. The structure (Fig. 4) also shows the Ge( $\pi$ ) centre coordinated to a tetradentate Me<sub>4</sub>cyclen macrocycle with the Ge–N bonds alternating shortlong-short-long around the ring, Ge–N = 2.165(6)–2.244(5) Å, with the Ge lying 1.009(3) Å above the mean N<sub>4</sub> plane. In both complexes the methyl substituents on nitrogen are directed to the same side of the plane as the germanium centre. There are no significant interactions to the triflate anions, therefore, the germanium is in a highly distorted square planar, or tetragonal pyramidal, environment.

The reaction of  $[GeCl_2(dioxane)]$  in  $CH_2Cl_2$  with  $Me_4cyclen$  in a 3:1 molar ratio gave the corresponding  $[Ge(Me_4cyclen)]$ - $[GeCl_3]_2$  which was less stable in solution.

#### Silicon(IV) complexes

The reaction of SiCl<sub>4</sub>, Me<sub>3</sub>tacn and Me<sub>3</sub>SiO<sub>3</sub>SCF<sub>3</sub> in anhydrous CH<sub>2</sub>Cl<sub>2</sub> solution produced [SiCl<sub>3</sub>(Me<sub>3</sub>tacn)][O<sub>3</sub>SCF<sub>3</sub>], which is *much* more hydrolytically sensitive than the [SiF<sub>3</sub>(Me<sub>3</sub>tacn)][SiF<sub>5</sub>] salt.<sup>11</sup> Crystals of this chloro-complex were obtained by layering a concentrated CH<sub>2</sub>Cl<sub>2</sub> solution with hexane. The structure of the cation shows the expected *fac*-octahedral coordination (Fig. 5). Comparison with the [SiF<sub>3</sub>(Me<sub>3</sub>tacn)]<sup>+</sup> cation shows negligible differences between the Si–N distances.

The corresponding  $[SiCl_3(Me_3tacn)][BAr^F]$  was obtained by reacting  $SiCl_4$  and  $Na[BAr^F]$  in toluene, followed by addition of a solution of  $Me_3tacn$ . As we have described elsewhere,<sup>19</sup> if the reagents are added simultaneously, the product is the sodium complex of the ligand, rather than the silicon cation. Crystals of  $[SiCl_3(Me_3tacn)][BAr^F]$  were not obtained, however, a concentrated  $CH_2Cl_2$  solution of the  $[BAr^F]^-$  salt, layered with hexane, produced a few colourless crystals identified by an X-ray structure determination as  $[SiCl_3(Me_3tacn)]Cl$  (see Fig. S1 in ESI<sup>†</sup>) which separate adventitiously due to the lower solubility of



Fig. 5 View of the cation present in  $[SiCl_3(Me_3tacn)][O_3SCF_3]$ . Thermal ellipsoids are drawn at 50% probability, hydrogen atoms omitted for clarity. Selected bond lengths (Å) and angles (°): Si1-Cl1 = 2.1867(9), Si1-Cl2 = 2.1688(8), Si1-Cl3 = 2.1416(9), Si1-N1 = 2.010(2), Si1-N2 = 2.013(2), Si1-N3 = 2.026(2); N1-Si1-N2 = 85.61(7), N2-Si1-N3 = 84.70(6), N1-Si1-N3 = 85.37(6), Cl1-Si1-Cl2 = 91.79(4), Cl1-Si1-Cl3 = 92.18(3), Cl2-Si1-Cl3 = 92.76(3).

this salt. Direct reaction of  $SiCl_4$  and  $Me_3tacn$  in  $CH_2Cl_2$  resulted in precipitation of a pale yellow solid containing the same [SiCl\_3(Me\_3tacn)]Cl, however we have been unable to obtain it in analytically pure form by this route.

The <sup>1</sup>H NMR spectrum of [SiCl<sub>3</sub>(Me<sub>3</sub>tacn)]<sup>+</sup> in CD<sub>3</sub>CN at 298 K shows sharp second order multiplets characteristic of symmetrical *fac*-coordinated Me<sub>3</sub>tacn in solution.<sup>20</sup>

Similar reaction of SiHCl<sub>3</sub> with Na[BAr<sup>F</sup>] in toluene, followed by addition of Me<sub>3</sub>tacn, produced the corresponding dichlorosilane complex, [SiHCl<sub>2</sub>(Me<sub>3</sub>tacn)][BAr<sup>F</sup>]. The presence of the Si-H group is shown by a singlet in the <sup>1</sup>H NMR spectrum at  $\delta = 4.78$  and by  $\nu$ (SiH) in the IR spectrum at 2137 cm<sup>-1</sup>. The presence of the *fac*-SiHCl<sub>2</sub> removes the threefold symmetry of the Me<sub>3</sub>tacn found in [SiCl<sub>3</sub>(Me<sub>3</sub>tacn)]<sup>+</sup>, and this is reflected in both the <sup>1</sup>H and <sup>13</sup>C $_{1}^{1}H$  NMR spectra, which show two  $\delta(Me)$  resonances and corresponding splitting of the NCH<sub>2</sub> resonances. Attempts to determine the structure of this complex have been unsuccessful. Poor quality crystals were obtained, but structure solution showed disordered [SiCl<sub>3</sub>(Me<sub>3</sub>tacn)]<sup>+</sup> and [SiHCl<sub>2</sub>(Me<sub>3</sub>tacn)]<sup>+</sup> were both present (note that  $[SiCl_3(Me_3tacn)]^+$  is not present in the NMR spectra of the bulk product). Disproportionation of silane complexes of amine ligands has been noted in other systems, and presumably occurs here slowly over the time taken to grow crystals from the [SiHCl<sub>2</sub>(Me<sub>3</sub>tacn)]<sup>+</sup> solution.<sup>21</sup>

Finally, two complexes of the linear triamine,  $Me_2NCH_2CH_2N(Me)CH_2CH_2NMe_2$  (pmdta), [SiX<sub>3</sub>(pmdta)]-[BAr<sup>F</sup>] (X = Cl or Br) were isolated by reacting the appropriate SiX<sub>4</sub> with Na[BAr<sup>F</sup>] in toluene, followed by addition of pmdta. The X-ray crystal structure of the chloride reveals a *mer*-geometry (Fig. 6) which contrasts with the *fac* arrangement found in the Me<sub>3</sub>tacn complexes. The geometry of the [SiCl<sub>3</sub>(pmdta)]<sup>+</sup> cation is close to octahedral, and the Si–N and Si–Cl distances are little different from those found in [SiCl<sub>3</sub>(Me<sub>3</sub>tacn)]<sup>+</sup>



**Fig. 6** View of the cation in  $[SiCl_3(pmdta)][BAr^F]$ . Thermal ellipsoids are drawn at 50% probability and hydrogen atoms are omitted for clarity. Selected bond lengths (Å) and angles (°): Si1–Cl1 = 2.147(2), Si1–Cl2 = 2.124(2), Si1–Cl3 = 2.184(2), Si1–N1 = 2.059(5), Si1–N2 = 2.028(5), Si1–N3 = 2.075(5); N1–Si1–N2 = 85.6(2), N2–Si1–N3 = 85.0(2) Cl1–Si1–Cl2 = 91.12(9), Cl2–Si1–Cl3 = 89.82(9).

despite the different donor arrangements. The geometry is also similar to that found in *mer*-[SiHCl<sub>2</sub>(pmdta)]<sup>+</sup> (which has H *trans* to Cl).<sup>22</sup>

The spectroscopic data are unexceptional, but consistent with the *mer* geometries, although it is notable that the terminal –NMe<sub>2</sub> groups appear as a broad singlet rather than the two resonances expected due to the inequivalence produced by the central –NMe group lying out of the N<sub>3</sub>Cl plane. The ease of formation of the [SiX<sub>3</sub>(pmdta)]<sup>+</sup> cations (X<sub>3</sub> = Cl<sub>3</sub>, Br<sub>3</sub>, HCl<sub>2</sub>) is in contrast with the  $\kappa^2$ -coordinated pmdta adduct formed with SiF<sub>4</sub>, reflecting the much higher Si–F bond strength.<sup>11</sup>

#### Conclusions

Two series of complexes with neutral aza-macrocyclic coordination to  $Ge(\pi)$  and Si(w) are reported and structurally characterised. The hydrolytically sensitive  $Ge(\pi)$  triaza macrocyclic complexes show pyramidal  $GeN_3$  coordination with very weak, but variable, interactions from the chloride and triflate anions. This contrasts with the 'naked' dication identified in the previously reported  $[Ge(Me_3tacn)]Br[GeBr_3]$ .<sup>12</sup> The tetra-aza macrocyclic complexes of  $Ge(\pi)$  give  $GeN_4$  dications in highly distorted square planar geometries, with no significant anion interactions towards  $Ge(\pi)$ .

The triaza ligand complexes of Si(IV), formed by halide abstraction using either  $Me_3SiO_3SCF_3$  or  $Na[BAr^F]$ , adopt distorted octahedral coordination geometries, and the trichloro-, tribromo- and hydridodichloro-species reported here are significantly more readily hydrolysed than  $[SiF_3(Me_3tacn)]^+$ .<sup>11</sup>

#### Acknowledgements

We thank EPSRC for support (EP/I010890/1 and EP/I033394/1). The SCFED Project (http://www.scfed.net) is a multidisciplin-

ary collaboration of British universities investigating the fundamental and applied aspects of supercritical fluids.

#### References

- 1 D. Vaughn II and R. E. Schaak, *Chem. Soc. Rev.*, 2013, 42, 2861.
- 2 S. Raoux, W. Welnic and D. Ielmini, *Chem. Rev.*, 2010, **110**, 240.
- 3 D. V. Talapin, J. S. Lee, M. V. Kovalenko and E. V. Shevchenko, *Chem. Rev.*, 2010, **110**, 389.
- 4 (a) J. Ke, W. Su, S. M. Howdle, M. George, D. Cook, M. Perdjon-Abel, P. N. Bartlett, W. Zhang, F. Cheng, W. Levason, G. Reid, J. Hyde, J. Wilson, D. C. Smith, K. Mallik and P. Sazio, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 14768; (b) J. Ke, P. N. Bartlett, D. Cook, T. L. Easun, M. George, W. Levason, G. Reid, D. C. Smith, W. Su and W. Zhang, *Phys. Chem. Chem. Phys.*, 2014, **16**, 9202.
- 5 P. N. Bartlett, S. L. Benjamin, C. H. de Groot,
  A. L. Hector, A. Jolleys, R. Huang, G. P. Kissling,
  W. Levason, S. J. Pearce, G. Reid and Y. Wang, *Mater. Horizons*, 2015, 2, 420.
- 6 (a) P. N. Bartlett, C. Y. Cummings, W. Levason, D. Pugh and G. Reid, *Chem. Eur. J.*, 2014, 20, 5019;
  (b) J. Ke, P. N. Bartlett, D. Cook, T. L. Easun, M. W. George, W. Levason, G. Reid, D. Smith, W. Su and W. Zhang, *Phys. Chem. Chem. Phys.*, 2012, 14, 1517.
- 7 W. Levason, G. Reid and W. Zhang, *Coord. Chem. Rev.*, 2011, 255, 1319.
- 8 (*a*) R. S. Ghadwal, H. W. Roesky, S. Merkel, J. Henn and D. Stalke, *Angew. Chem., Int. Ed.*, 2009, 48, 5683;
  (*b*) A. C. Filippou, O. Chernov and G. Schnakenburg, *Angew. Chem., Int. Ed.*, 2009, 48, 5687.
- 9 F. Cheng, M. F. Davis, A. L. Hector, W. Levason, G. Reid, M. Webster and W. Zhang, *Eur. J. Inorg. Chem.*, 2007, 4897.
- 10 G. Willey, T. J. Woodman, U. Somasundaram, D. R. Aris and W. Errington, *J. Chem. Soc., Dalton Trans.*, 1998, 2573.
- 11 F. Cheng, A. L. Hector, W. Levason, G. Reid, M. Webster and W. Zhang, *Chem. Commun.*, 2009, 1334.
- F. Cheng, A. L. Hector, W. Levason, G. Reid, M. Webster and W. Zhang, *Angew. Chem., Int Ed.*, 2009, 48, 5152.
- 13 K. Wieghardt, P. Chaudhuri, B. Nuber and J. Weiss, *Inorg. Chem.*, 1982, **21**, 3086.
- 14 M. Brookhart, B. Grant and A. F. Volpe Jr., *Organometallics*, 1992, **11**, 3920.
- (a) CrystalClear-SM Expert 2.0 r7.4, Rigaku Corporation, Tokyo, Japan, 2011; (b) L. J. Farrugia, J. Appl. Crystallogr., 2012, 45, 849; (c) O. V. Dolomanov, L. J. Bourhis, R. J. Gildea, J. A. K. Howard and H. Puschmann, J. Appl. Crystallogr., 2009, 42, 339; (d) G. M. Sheldrick, Acta Crystal-

*logr., Sect. C: Cryst. Struct. Commun.*, 2015, **71**, 3; (*e*) F. H. Allen, O. Johnson, G. P. Shields, B. R. Smith and M. Towler, *J. Appl. Crystallogr.*, 2004, **37**, 335.

- See for example: J. Dyke, W. Levason, M. E. Light, D. Pugh, G. Reid, H. Bhakhoa, P. Ramasami and L. Rhyman, *Dalton Trans.*, 2015, 44, 13853.
- 17 M. Mantina, A. C. Chamberlin, R. Valero, C. J. Cramer and D. G. Truhlar, *J. Phys. Chem. A*, 2009, **113**, 5806.
- 18 B. Cordero, V. Gomez, A. E. Platero-Prats, M. Reves, J. Echeverria, E. Cremades, F. Barragan and S. Alvarez, *Dalton Trans.*, 2008, 2832.
- 19 M. Everett, A. Jolleys, W. Levason, D. Pugh and G. Reid, *Chem. Commun.*, 2014, **50**, 5843.
- 20 R. Bhalla, C. Darby, W. Levason, S. K. Luthra, G. McRobbie,
  G. Reid, G. W. Sanderson and W. Zhang, *Chem. Sci.*, 2014,
  5, 381.
- 21 (*a*) P. Boudjouk, S. D. Kloos, B.-K. Kim, M. Page and D. Thweatt, *J. Chem. Soc., Dalton Trans.*, 1998, 877;
  (*b*) H. J. Campbell-Ferguson and E. A. V. Ebsworth, *J. Chem. Soc. A*, 1966, 1508.
- 22 B.-K. Kim, S.-B. Choi, S. D. Kloos and P. Boudjouk, *Inorg. Chem.*, 2000, **39**, 728.