



Cite this: *Org. Biomol. Chem.*, 2015, **13**, 2092

Asymmetric syntheses of three-membered heterocycles using chiral amide-based ammonium ylides†

Mathias Pichler,^a Johanna Novacek,^a Raphaël Robiette,^b Vanessa Poscher,^a Markus Himmelsbach,^c Uwe Monkowius,^d Norbert Müller^a and Mario Waser*^a

The use of carbonyl-stabilised ammonium ylides to access chiral glycidic amides and the corresponding aziridines has so far been limited to racemic *trans*-selective protocols. We herein report the development of an asymmetric approach to access such compounds with high levels of stereoselectivity using easily accessible chiral auxiliary-based ammonium ylides. The use of phenylglycinol as the chiral auxiliary was found to be superior to Evans or pseudoephedrine-based auxiliaries resulting in good to excellent stereoselectivities in both, epoxidation and aziridination reactions.

Received 31st October 2014,
Accepted 1st December 2014


DOI: 10.1039/c4ob02318h

www.rsc.org/obc

Introduction

The use of onium ylides has emerged as a powerful strategy for (dia)-stereoselective epoxide, aziridine, and cyclopropane syntheses.^{1,2} Whereas sulfonium ylides have been very frequently used for a variety of often very selective three-ring forming reactions,^{3,4} easily available ammonium ylides have been less routinely employed in the past.^{5–9} Especially the use of chiral amines to render such reactions enantioselective has been mainly limited to cyclopropanations⁵ so far. Indeed, their use in asymmetric epoxidation and aziridination reactions was found to be rather difficult, mainly due to the weaker leaving group ability of the amine group compared to the use of sulfonium ylides.^{6,7,9}

We have recently introduced a highly *trans*-selective protocol for the synthesis of glycidic amides and the corresponding aziridines by reacting amide-stabilised achiral ammonium ylide precursors **1** with aldehydes **2** or imines **3** (Scheme 1).^{7,9b} Key to high yields was the use of trimethylamine as the amine leaving group, whereas the use of sterically more demanding



Scheme 1 Recently developed racemic *trans*-epoxidation and aziridination protocol and the targeted auxiliary-based stereoselective approach.

chiral amines like Cinchona alkaloids did not result in any product formation. Due to this limitation, alternative strategies to control the absolute stereochemistry in these epoxide and aziridine forming reactions are necessary. The use of chiral auxiliary containing amides to control the face selectivity in enolate reactions is a very commonly employed and powerful strategy for numerous applications.¹⁰ A few examples about their use in sulfonium ylide-mediated reactions have been reported in the past¹¹ but the use of chiral amide-based ammonium ylides has, to the best of our knowledge, not been reported yet. We have therefore undertaken a systematic study about the use of different classes of chiral auxiliaries to access chiral ammonium ylide precursors **6** and explore their potential in asymmetric three-membered ring heterocycle-forming reactions (Scheme 1). The main focus was on the development

^aInstitute of Organic Chemistry, Johannes Kepler University Linz, Altenbergerstraße 69, 4040 Linz, Austria. E-mail: Mario.waser@jku.at; Fax: +43 732 2468 8747; Tel: +43 732 2468 8748


^bInstitute of Condensed Matter and Nanosciences, Université catholique de Louvain, Place Louis Pasteur 1 box 14.01.02, 1348 Louvain-la-Neuve, Belgium

^cInstitute of Analytical Chemistry, Johannes Kepler University Linz, Altenbergerstraße 69, 4040 Linz, Austria

^dInstitute of Inorganic Chemistry, Johannes Kepler University Linz, Altenbergerstraße 69, 4040 Linz, Austria

† Electronic supplementary information (ESI) available: Experimental and computational details, characterization of new compounds, and copies of NMR spectra. CCDC 1023711 and 1030561. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c4ob02318h





Scheme 5 Use of phenylglycinol (**14**) to access chiral sulfonium^{11e} and ammonium salts for asymmetric epoxidation reactions.

Because of the limited applicability of the Evans oxazolidinones and pseudoephedrine as chiral auxiliaries for our ammonium ylide-mediated epoxidation we next turned our attention on the use of 1,3-oxazolidine-containing auxiliaries. These are less commonly employed in asymmetric transformations as compared to oxazolidinones or pseudoephedrine. However, we reasoned that the corresponding ammonium salts **6C** may work well for our target reaction as they should be stable under the basic reaction conditions and the addition step should hopefully proceed with satisfying face-selectivity too. In addition, during the initial phase of this project, Teran *et al.* reported the use of chiral 1,3-oxazolidines (derived from the reaction of phenylglycinol (**14**) with different aldehydes) in sulfur ylide-mediated epoxidation reactions.^{11e} Interestingly, they found that the absolute configuration of the product **16** depends exclusively on the configuration of the phenylglycinol moiety (C4) – and not on the configuration of the newly installed stereogenic center (C2) of the auxiliary. We opted for a slightly different approach by carrying out the oxazolidine formation by reacting (*R*)-**14** with acetone, which proceeds very quickly giving the auxiliary in high yield and purity. We anticipated that this ketone-based 1,3-oxazolidine should show a higher stability under basic conditions than aldehyde-based ones and should therefore be suitable for our approach. The ylide precursor **6C** could then easily be obtained in two more steps (Scheme 5).

In the initial epoxidation experiment of **6C** with benzaldehyde (**2a**) in dichloromethane (using 6 eq. of Cs₂CO₃ as a solid base) epoxide **17a** was formed as a single *trans*-diastereomer. Based on this promising result a systematic screening of different conditions was undertaken (Table 1 gives an overview about the most significant results obtained herein). In our attempts to determine the best-suited base and the optimum stoichiometric ratio we found that the strongly basic aqueous conditions that worked well in our racemic protocol⁷ did not give any product herein (entry 2). The use of K₂CO₃ (entry 3) or an equimolar amount of Cs₂CO₃ (entry 4) yielded also almost no product. However, an excess of Cs₂CO₃ and two equivalents of aldehyde allowed us to obtain *trans*-**17a** in 50% isolated

Table 1 Identification of the optimum reaction conditions for the epoxidation using the chiral ammonium ylide precursor **6C**



Entry	2a (eq.)	Solv.	Base (eq.)	<i>T</i> [°C]	<i>t</i> [h]	Yield ^a [%]	dr ^b [%] (<i>trans</i>) ^c
1	4	CH ₂ Cl ₂	Cs ₂ CO ₃ (6×)	25	24	35	>98
2	2	CH ₂ Cl ₂	NaOH (aq) (140×)	25	24	n.r.	n.d.
3	2	CH ₂ Cl ₂	K ₂ CO ₃ (20×)	25	24	n.r.	n.d.
4	1	CH ₂ Cl ₂	Cs ₂ CO ₃ (1×)	25	24	<10	>98
5	2	CH ₂ Cl ₂	Cs ₂ CO ₃ (20×)	25	24	50 ^d	>98
6	2	CH ₂ Cl ₂	Cs ₂ CO ₃ (20×)	25	72	74	>98
7	2	THF	Cs ₂ CO ₃ (20×)	25	24	n.r.	n.d.
8	2	Toluene	Cs ₂ CO ₃ (20×)	25	24	50 ^d	>98
9	2	Toluene	Cs ₂ CO ₃ (20×)	25	72	72	>98
10	2	Toluene	Cs ₂ CO ₃ (20×)	60	24	73	>98
11	2	Toluene	Cs ₂ CO ₃ (20×)	60 ^e	3	75	>98
12	2	i-PrOH	Cs ₂ CO ₃ (20×)	25	24	78	>98
13	2	i-PrOH	Cs ₂ CO ₃ (20×)	60 ^e	3	84	>98

^a Isolated yield. ^b Determined by ¹H NMR of the crude reaction mixture. ^c In neither experiment any *cis*-diastereomer could be detected. ^d Incomplete conversion of **6C**. ^e Carried out in an ultrasonic bath.


yield after 24 h and in 74% after 72 h reaction time (entries 5 and 6). As expected from our previous experience using amide-stabilised ammonium ylides⁷ no *cis*-epoxide was formed under either conditions. In addition, also no second diastereoisomeric *trans*-epoxide could be observed in NMR spectra of the crude reaction mixture, illustrating that this chiral auxiliary leads to a high face-differentiation in the ylide addition step. Further screening of different solvents showed that toluene and i-PrOH can also be used as solvents for this reaction. Whereas the reaction was found to be slow in toluene, requiring either a prolonged reaction time (entry 9) or a higher reaction temperature (entry 10) to obtain **17a** in good yield, reactions in i-PrOH were usually significantly faster (entry 12). However, hereby also, significant amounts of Cannizzaro disproportionation products of benzaldehyde could be detected whereas this side reaction is less pronounced in toluene. In addition, we also found that these reactions can be accelerated by ultrasonication, giving **17a** in comparable quality and yield after 3 h (entries 11 and 13). Accordingly, this screening allowed us to identify different conditions which all gave the target epoxide **17a** with very high diastereoselectivity and with isolated yields >70%.

The absolute configuration of **17a** was unambiguously proven by two different methods. First X-ray diffraction analysis of crystals of **17a** proved the (2*R*,3*S*)-configuration of the *trans*-epoxyamide moiety.¹⁵ In addition **17a** was also transferred into the known epoxyalcohol **18** upon treatment with LiBHET₃. Comparison of the specific optical rotation with literature values of **18**^{11a,e} also confirmed this absolute con-





Scheme 6 Molecular structure of 17a and conversion into the known epoxyalcohol 18.



Scheme 7 DFT calculations of the ylide geometry (relative energies given in brackets) and proposed rationale for observed diastereoselectivity.¹⁶

figuration of the epoxy moiety (Scheme 6). Unfortunately, this transformation did not allow us to recover the auxiliary.

In order to get an insight into the interpretation of the observed high diastereoselectivity in all these epoxidation reactions, we undertook DFT calculations investigating the geometry of the most stable conformer of the ylide.^{16,17} It turned out that (Z)-ylides are significantly more stable than (E)-ylides (Scheme 7). This can most probably be accounted for by de-

Table 2 Application scope of the asymmetric epoxidation using amide 6C

Entry	R	Ald.	Prod.	Cond. ^a	Yield ^b [%]	dr ^c [%]
1	Ph	2a	17a	A	78	>98
2				B	73	>98
3	4-MeC ₆ H ₄ -	2b	17b	A	80	>98
4	2-MeC ₆ H ₄ -	2c	17c	A	79	>98
5	4-ClC ₆ H ₄ -	2d	17d	A	75	>98
6	4-BrC ₆ H ₄ -	2e	17e	A	85	>98
7	4-PhC ₆ H ₄ -	2f	17f	A	89	>98
8	4-MeOC ₆ H ₄ -	2g	17g	A	73	>98
9	4-Me ₂ NC ₆ H ₄ -	2h	17h	B	(60) ^d	>98
10	4-CNC ₆ H ₄ -	2i	17i	C	62	>98
11	3-NO ₂ C ₆ H ₄ -	2j	17j	C	68	>98
12	<i>n</i> -Decyl-	2k	17k	D	42	>98
13	Cyclohexyl-	2l	17l	D	39	>98

^a A: *i*-PrOH, 25 °C, 24 h, Cs₂CO₃ (s, 20 eq.); B: toluene, 60 °C, 24 h, Cs₂CO₃ (s, 20 eq.); C: toluene, 25 °C, 72 h, Cs₂CO₃ (s, 20 eq.); D: toluene, 25 °C, 24 h, Cs₂CO₃ (s, 20 eq.). ^b Isolated yield. ^c Determined by ¹H NMR of the crude reaction mixture. ^d Full decomposition on silica gel, crude NMR yield given in brackets.


stabilizing steric interactions between the ammonium group and the auxiliary in the (E)-ylide and the presence of stabilizing electrostatic interactions between the ammonium group and the enolate oxygen in the (Z)-isomer.¹⁸ In addition, the (Z)-ylide in which the phenyl group blocks the *Re*-face of the α -carbon ((Z)-ylide^{Re}) is thermodynamically more stable than the one where the *Si*-face is shielded ((Z)-ylide^{Si}) (Scheme 7, upper part).

Based on these results, the observed stereoselectivity could be explained by addition of the most stable ylide ((Z)-ylide^{Re}) to benzaldehyde 2a via the *Si*-face (Scheme 7, lower part).¹⁷

With a set of high yielding reaction conditions available we next investigated the application scope of this epoxidation reaction. As shown in Table 2 a series of different aldehydes was tested. Notably, in neither case any *cis*-epoxide and no second *trans*-diastereomer could be detected. Due to the faster reaction rate of the parent benzaldehyde 2a in *i*-PrOH at room temperature the majority of the reactions were carried out under these conditions (cond. A). These worked reasonably fine for a series of aromatic aldehydes such as *para*- or *ortho*-methyl benzaldehydes (entries 3 and 4), halide-substituted ones (entries 5 and 6) and the biphenyl carbaldehyde 2f (entry 7). Also the more electron-rich *para*-methoxy benzaldehyde 2g could be reacted in good yield under these conditions, whereas the less active dimethylamino benzaldehyde 2h required harsher conditions (60 °C in toluene) to obtain the epoxide 17h in good crude yield. Remarkably, this was the first time that this aldehyde could be successfully used in any of our ammonium ylide-mediated epoxidation reactions.⁷



Table 3 Asymmetric aziridination using amide 6C




Entry	PG	Ar	3	19 (dr) ^{a,b} [%]	20 ^a [%]	Yield [%] (<i>trans</i> -19) ^c
1	Tos	Ph	3a	>98 <i>trans</i> ^d	n.d.	39 (70) ^e
2	Boc		3b	82 (87/3/10) ^b	18	62
3		2-MeC ₆ H ₄ -	3c	96 (85/3/12) ^b	4	56
4		4-MeOC ₆ H ₄ -	3d	95 (88/3/9) ^b	5	57 (75) ^f
5		Naph-2-yl	3e	77 (85/3/12) ^b	23	58
6		4-BrC ₆ H ₄ -	3f	39 (86/0/14) ^b	61	32
7		3-NO ₂ C ₆ H ₄ -	3g	n.d.	>99	n.d.

^a Determined by ¹H NMR of the crude reaction mixture. ^b Values in brackets give the diastereomeric ratios of aziridines (*trans*_{major}/*trans*_{minor}/*cis*) – only one *cis*-isomer could be detected. ^c Isolated yield of the major *trans*-aziridine. ^d Only one *trans*-aziridine detected. ^e Partial decomposition on silica gel, crude NMR yield given in brackets. ^f Full decomposition on silica gel and partial on alumina, crude NMR yield given in brackets.

Unfortunately, epoxide **17h** quickly decomposed during different purification methods. As expected, the more reactive cyano- and nitrobenzaldehydes **2i** and **2j** did not give any epoxide under the standard conditions (cond. A) but mainly the corresponding Cannizzaro disproportionation products. However, carrying out the reaction in toluene as a less polar solvent at room temperature allowed us to obtain the epoxides **17i** and **17j** in reasonable isolated yields. Notably, for the first time we have been able to use the aliphatic aldehyde **2k** (entry 12) to obtain the corresponding epoxide in moderate yield. Such enolisable aldehydes had primarily undergone self aldol condensation reactions under the strongly basic conditions developed previously.⁷ Now the use of Cs₂CO₃ in toluene allowed us to overcome this limitation to some extent. Similar yield was obtained when reacting cyclohexanecarbaldehyde (**2l**) under these conditions (entry 13).

Based on the detailed knowledge obtained on the use of the chiral amide **6C** for asymmetric ylide-mediated epoxidations at hand, we carried out a short screening to test the potential of this concept for the related aziridination reaction (Table 3). In analogy to our previous experience using achiral amides **6** for racemic aziridinations^{9b} the use of Cs₂CO₃ in CH₂Cl₂ was also found superior here compared to the use of *i*-PrOH or toluene as solvents. Interestingly, the use of *N*-tosyl imine **3a** allowed the synthesis of the *trans*-aziridine **19a** in high selectivity (entry 1). Unfortunately this compound was found to be relatively unstable and partially decomposed under a variety of purification conditions. This may be attributed to the rather strained nature of this *trans*-aziridine with the bulky tosyl group being *cis* to either the phenyl group or the chiral auxiliary.^{4c} The use of *N*-Boc imine **3b** gave an interesting result (entry 2). Formation of the expected major *trans*-aziridine

Fig. 1 Molecular structure of **19f**.

idine **19b** (isolated in 61% yield) was accompanied by minor amounts of the second *trans*-aziridine as well as around 10% of *cis*-**19b**. Besides this reduced stereoselectivity in the aziridine formation (compared to epoxidation) also notable amounts of the α,β -unsaturated β -amino amide **20b** were obtained. Formation of analogous olefins was already observed during our previous racemic studies but only when using electron-poor *N*-Boc benzaldehydes (*e.g.* NO₂-substituted) or hetero-aromatic *N*-Boc aldimines, whereas not even traces thereof could be detected using more electron-rich imines like **3b**.^{9b} Unfortunately we were not able to suppress formation of **20** by changing the reaction conditions (in contrast: using less base gave a larger amount of **20**). When using more electron-rich imines **3c** and **3d** the formation of olefins **20c** and **20d** was reduced, but still not totally suppressed (entries 3 and 4). Again, formation of the minor *trans*-aziridine and *cis*-diastereomer was observed too. While the isolated yield of **19c** may be a bit lower because of a reduced rate of addition due to the *ortho*-substituent, the more electron-rich *p*-methoxy containing **19d** was found to be totally unstable on silica gel and also partially decomposed during alumina column chromatography.¹⁹ The naphthyl-based imine **3e** performed similarly to imine **3b** (compare entries 5 and 2) giving the major *trans*-aziridine **19e** in a reasonable isolated yield of 59%. By contrast, the less electron-rich bromine-substituted **3f** gave the olefin **20f** as the main product (entry 6), while, as expected, the electron-poor **3g** did not allow us to obtain any aziridine at all (entry 7).

The absolute configuration of the major *trans*-aziridine was determined by anomalous X-ray diffraction analysis of crystals of bromine-containing **19f**. Again, in analogy to the epoxidation reaction, the (2*R*,3*S*)-configuration of the *trans*-heterocyclic moiety was found for the major isomer (Fig. 1).¹⁵

Experimental

General

¹H- and ¹³C-NMR spectra were recorded on a Bruker Avance DRX 500 MHz spectrometer, a Bruker Avance III 300 MHz spectrometer, and on a Bruker Avance III 700 MHz spectrometer with TCI cryoprobe. All NMR spectra were referenced using the residual ¹H solvent peak as secondary reference. High resolution mass spectra were obtained using a Thermo



Fisher Scientific LTQ Orbitrap XL with an Ion Max API Source. All analyses were made in the positive ionisation mode. IR spectra were recorded on a Bruker Tensor 27 FT-IR spectrometer with ATR unit. All chemicals were purchased from commercial suppliers and used without further purification unless otherwise stated. All reactions were performed under an Ar-atmosphere. CH₂Cl₂ was distilled over P₂O₅ and stored under Ar (it was not necessary to dry CH₂Cl₂ prior to every experiment and usually this quality could be used successfully in these reactions over the course of 3–4 weeks after distillation). Single-crystal structure analyses were carried out on a Bruker Smart X2S diffractometer operating with Mo-K_α radiation ($\lambda = 0.71073 \text{ \AA}$).

Geometry optimization has been performed using the Jaguar 8.0 pseudospectral program package using the well-established B3LYP hybrid density functional with the D3 dispersion correction and the standard split valence polarized 6-31G* basis as implemented in Jaguar. All the optimization calculations were carried out using the Poisson–Boltzmann polarizable continuum method as incorporated in Jaguar, and parameters for CH₂Cl₂. Energies were obtained by single point energy calculations at the B3LYP-D3/6-311+G**(CH₂Cl₂) level. The correct nature of each stationary point has been checked by performing frequency calculations at the B3LYP/6-31G*(CH₂Cl₂) level of theory. Thermal and entropic contributions to free energy (at 298.15 K) and zero-point energy have been obtained from these frequency calculations. We have made a systematic attempt to locate all possible local minima, with the data presented referring to the lowest energy form.

Ammonium amide 6C

(*R*)-**14** (3.00 g, 21.9 mmol) was dissolved in 50 mL acetone and 3.2 g anhydrous MgSO₄ were added and the reaction mixture was stirred for 3 h at 20 °C. After filtration and evaporation to dryness the product **21** was obtained in 95% (3.68 g, 20.7 mmol) and used without further purification. The ¹H-NMR-spectrum is in full accordance to literature.²⁰ ¹H-NMR (300 MHz, δ , CDCl₃, 298 K): 1.45 (s, 3H), 1.52 (s, 3H), 2.04 (b, 1H), 3.71 (t, 1H, $J = 7.8 \text{ Hz}$), 4.29 (t, 1H, $J = 7.8 \text{ Hz}$), 4.54 (t, 1H, $J = 7.8 \text{ Hz}$), 7.26–7.41 (m, 5H, Ar-H) ppm.

Compound **21** (3.68 g, 20.7 mmol) was dissolved in 25 mL CH₂Cl₂ and 83 mL aqueous saturated Na₂CO₃ solution were added. Then bromide **7** (2.9 mL, 21.7 mmol, 1.05 eq.) was added and the mixture was vigorously stirred for 4 h. After addition of aqueous saturated NaHCO₃ the aqueous layer was separated and washed three times with 20 mL CH₂Cl₂. The combined organic phases were dried over anhydrous MgSO₄, filtrated and the solvent removed under reduced pressure. The product was purified by column chromatography (silica gel, heptanes–EtOAc = 5 : 1) to give **8C** (2.57 g, 8.6 mmol, 41% yield) as a light-brown solid. The ¹H-NMR-spectrum was in accordance to literature.²¹ ¹H-NMR (300 MHz, δ , CDCl₃, 298 K): 1.64 (s, 3H), 1.87 (s, 3H), 3.45 (d, 1H, $J = 11.0 \text{ Hz}$), 3.52 (d, 1H, $J = 11.0 \text{ Hz}$), 3.94 (dd, 1H, $J_1 = 9.0 \text{ Hz}$, $J_2 = 2.7 \text{ Hz}$), 4.41 (dd, 1H, $J_1 = 9.0 \text{ Hz}$, $J_2 = 6.5 \text{ Hz}$), 5.07 (dd, 1H, $J_1 = 6.5 \text{ Hz}$, $J_2 = 2.7 \text{ Hz}$), 7.25–7.45 (m, 5H) ppm.

Compound **8C** (2.57 g, 8.6 mmol) was dissolved in THF (26 mL) and NMe₃ (2.50 mL, 20.4 mmol, 1.2 eq., 33% solution in EtOH) was added. After stirring for 24 h at r.t. the solvent was removed with vacuum distillation. The crude product (purity >90%) was purified by column chromatography (heptanes → CH₂Cl₂–MeOH = 5 : 1) to give the ammonium salt **6C** in 90% yield (2.90 g, 8.1 mmol) as a white foam. [α]_D²² ($c = 0.6$, CH₂Cl₂) = –92; ¹H-NMR (700 MHz, δ , CDCl₃, 298 K): 1.65 (s, 3H), 1.86 (s, 3H), 2.98 (d, 1H, $J = 16.3 \text{ Hz}$), 3.44 (s, 9H), 3.91 (dd, 1H, $J_1 = 9.2 \text{ Hz}$, $J_2 = 1.7 \text{ Hz}$), 4.46 (dd, 1H, $J_1 = 9.2 \text{ Hz}$, $J_2 = 6.5 \text{ Hz}$), 5.83 (dd, 1H, $J_1 = 6.5 \text{ Hz}$, $J_2 = 1.7 \text{ Hz}$), 6.07 (d, 1H, $J = 16.3 \text{ Hz}$), 7.28–7.51 (m, 5H) ppm; ¹³C NMR (176 MHz, δ , CDCl₃, 298 K): 23.5, 25.5, 54.7, 60.1, 64.8, 71.8, 97.5, 126.8, 128.5, 129.5, 140.5, 161.0 ppm; IR (film): $\bar{\nu} = 3011, 2987, 2937, 2882, 1654, 1434, 1412, 1378, 1351, 1237, 1204, 1133, 1064, 1048, 923, 896, 843, 703, 664, 604, 579, 563, 517, 501 \text{ cm}^{-1}$; HRMS (ESI): m/z calcd for C₁₆H₂₅N₂O₂⁺: 277.1910 [M]⁺; found: 277.1904.

General epoxidation procedure using ammonium amide 6C

Ammonium salt **6C** was dissolved in the appropriate solvent (20 mL mmol^{–1} ammonium salt) and Cs₂CO₃ (20 eq.) was added to the reaction mixture. After 5 min the aldehyde (2 eq.) was added and the suspension was stirred for the indicated time at the given temperature. The reaction was quenched with water and extracted with toluene. The organic phase was washed with brine and dried with anhydrous Na₂SO₄, filtrated and the solvent was removed under reduced pressure. The epoxide was purified by column chromatography (silica gel, heptanes–EtOAc = 7 : 3).

trans-Epoxyde 17a. Obtained in 78% (1 mmol scale) as a white solid after column chromatography (Cond. A). [α]_D²² ($c = 1.4$, CH₂Cl₂) = –178; ¹H NMR (700 MHz, δ , CDCl₃, 298 K): 1.72 (s, 3H), 1.88 (s, 3H), 3.22 (d, 1H, $J = 1.8 \text{ Hz}$), 3.68 (d, 1H, $J = 1.8 \text{ Hz}$), 3.90 (dd, 1H, $J_1 = 9.2 \text{ Hz}$, $J_2 = 4.2 \text{ Hz}$), 4.41 (dd, 1H, $J_1 = 9.2 \text{ Hz}$, $J_2 = 6.6 \text{ Hz}$), 5.10 (dd, 1H, $J_1 = 6.6 \text{ Hz}$, $J_2 = 4.2 \text{ Hz}$), 6.78 (m, 2H), 7.00–7.20 (m, 8H) ppm; ¹³C NMR (176 MHz, δ , CDCl₃, 298 K): 24.3, 25.1, 58.4, 58.9, 61.5, 72.1, 97.3, 125.9, 126.1, 128.4, 128.5, 128.7, 129.4, 135.3, 140.6, 164.2 ppm; IR (film): $\bar{\nu} = 3032, 2989, 2933, 2873, 1658, 1458, 1437, 1387, 1363, 1252, 1205, 1081, 1066, 894, 849, 772, 749, 695, 660, 596, 552, 513 \text{ cm}^{-1}$; HRMS (ESI): m/z calcd for C₂₀H₂₁NO₃: 324.1594 [M + H]⁺; found: 324.1595.

General procedure for the preparation of aziridines

A mixture of **6C** (0.4 mmol), aldimine **3** (2 eq.), and Cs₂CO₃ (20 eq.) in CH₂Cl₂ (8 mL) was vigorously stirred for 24 h at room temperature. CH₂Cl₂ and brine were added and the phases separated. The aqueous layer was extracted twice with CH₂Cl₂, the combined organic layers were extracted with brine and the aqueous layer was re-extracted twice with CH₂Cl₂. The combined organic layers were dried over Na₂SO₄, filtrated, evaporated, and dried *in vacuo*. Column chromatography (silica gel, heptanes–EtOAc = 20 : 1–2 : 1) gave the aziridines **19** in the reported yields. In most cases the minor *cis*-isomers and the olefins **20** could not be obtained in pure form.



trans-N-Boc aziridine 19b. Obtained in 62% as a colourless residue. $[\alpha]_{\text{D}}^{23}$ ($c = 1.6$, CH_2Cl_2) = -110 ; $^1\text{H NMR}$ (500 MHz, δ , CDCl_3 , 298 K): 1.48 (s, 9H), 1.73 (s, 3H), 1.89 (s, 3H), 2.87 (d, 1H, $J = 2.5$ Hz), 3.69 (d, 1H, $J = 2.5$ Hz), 3.96 (dd, 1H, $J = 9.1$, 3.5 Hz), 4.47 (dd, 1H, $J = 9.1$, 6.8 Hz), 5.28 (dd, 1H, $J = 6.8$, 3.5 Hz), 6.88 (d, 2H, $J = 6.9$ Hz), 7.19–7.28 (m, 8H) ppm; $^{13}\text{C NMR}$ (125 MHz, δ , CDCl_3 , 298 K): 23.8, 25.0, 28.0, 44.5, 44.8, 61.5, 71.6, 81.7, 96.8, 125.7, 126.5, 127.7, 128.0, 128.1, 129.1, 135.3, 140.9, 159.2, 163.7 ppm; IR (film): $\bar{\nu} = 3009$, 2984, 2935, 2868, 1715, 1652, 1433, 1395, 1364, 1333, 1253, 1223, 1204, 1074, 1053, 819, 755, 746, 705, 691, 635 cm^{-1} ; HRMS (ESI): m/z calcd for $\text{C}_{25}\text{H}_{30}\text{N}_2\text{O}_4$: 423.2278 $[\text{M} + \text{H}]^+$; found: 423.2287.

Conclusions

The use of easily available phenylglycinol as a chiral auxiliary in ammonium ylide-mediated reactions was found to be a promising strategy to obtain chiral three-membered ring heterocycles with good to excellent stereoselectivities and in good yields. In general it was found that the epoxidation reaction is rather broad in its application scope giving glycidic amides as single stereoisomers in all cases. The aziridination is slightly less selective giving the major *trans*-isomer with at least 85% diastereoselectivity but, depending on the electronic nature of the starting imine, with varying amounts of an α,β -unsaturated β -amino amide side-product.

Acknowledgements

This work was supported by the Austrian Science Funds (FWF): Project no. P26387-N28. Computational resources have been provided by the supercomputing facilities of the Université catholique de Louvain (CISM/UCL) and the Consortium des Équipements de Calcul Intensif en Fédération Wallonie Bruxelles (CÉCI) funded by the Fond de la Recherche Scientifique de Belgique (F.R.S.-FNRS) under convention 2.5020.11. The NMR spectrometers used were acquired in collaboration with the University of South Bohemia (CZ) with financial support from the European Union through the EFRE INTER-REG IV ETC-AT-CZ program (project M00146, "RERI-uasb"). RR is a *Chercheur qualifié* of the F.R.S.-FNRS.

Notes and references

- For reviews see: (a) V. K. Aggarwal, in *Comprehensive Asymmetric Catalysis*, ed. E. N. Jacobsen, A. Pfaltz and H. Yamamoto, Springer, New York, 1999, vol. 2, p. 679; (b) E. M. McGarrigle, E. L. Myers, O. Illa, M. A. Shaw, S. L. Riches and V. K. Aggarwal, *Chem. Rev.*, 2007, **107**, 5841–5883.
- For seminal contributions about sulfur ylides see: (a) A. W. Johnson and R. B. Lacount, *Chem. Ind.*, 1958, 1440–1441; (b) E. J. Corey and M. Chaykovsky, *J. Am. Chem. Soc.*, 1962, **84**, 867–868; (c) E. J. Corey and M. Chaykovsky, *J. Am. Chem. Soc.*, 1965, **87**, 1353–1364.
- For selected examples highlighting the use of sulfur ylides in epoxidation reactions see: (a) O. Illa, M. Arshad, A. Ros, E. M. McGarrigle and V. K. Aggarwal, *J. Am. Chem. Soc.*, 2010, **132**, 1828–1830; (b) J. Zanardi, C. Leriverend, D. Aubert, K. Julienne and P. Metzner, *J. Org. Chem.*, 2001, **66**, 5620–5623; (c) M. Davoust, J.-F. Briere, P.-A. Jaffres and P. Metzner, *J. Org. Chem.*, 2005, **70**, 4166–4169; (d) V. K. Aggarwal and J. Richardson, *Chem. Commun.*, 2003, 2644–2651; (e) Y.-G. Zhou, X.-L. Hou, L.-X. Dai, L.-J. Xia and M.-H. Tang, *J. Chem. Soc., Perkin Trans. 1*, 1999, 77–80; (f) V. K. Aggarwal, J. P. H. Charmant, D. Fuentes, J. N. Harvey, G. Hynd, D. Ohara, W. Picoul, R. Robiette, C. Smith, J.-L. Vasse and C. L. Winn, *J. Am. Chem. Soc.*, 2006, **128**, 2105–2114; (g) V. K. Aggarwal, G. Hynd, W. Picoul and J.-L. Vasse, *J. Am. Chem. Soc.*, 2002, **124**, 9964–9965; (h) O. Illa, M. Namutebi, C. Saha, M. Ostovar, C. C. Chen, M. F. Haddow, S. Nocquet-Thibault, M. Lusi, E. M. McGarrigle and V. K. Aggarwal, *J. Am. Chem. Soc.*, 2003, **135**, 11951–11966; (i) A. Solladie-Cavallo and A. Diep-Vohuule, *J. Org. Chem.*, 1995, **60**, 3494–3498.
- For the use of sulfur ylides in aziridine synthesis see: (a) A.-H. Li, L.-X. Dai and X.-L. Hou, *J. Chem. Soc., Perkin Trans. 1*, 1996, 867–869; (b) V. K. Aggarwal, M. Ferrara, C. J. O'Brien, A. Thompson, R. V. H. Jones and R. Fieldhouse, *J. Chem. Soc., Perkin Trans. 1*, 2001, 1635–1643; (c) R. Robiette, *J. Org. Chem.*, 2006, **71**, 2726–2734; (d) T. Saito, M. Sakairi and D. Akiba, *Tetrahedron Lett.*, 2001, **42**, 5451–5454; (e) V. K. Aggarwal, E. Alonso, G. Fang, M. Ferrara, G. Hynd and M. Porcelloni, *Angew. Chem., Int. Ed.*, 2001, **40**, 1433–1436; (f) V. K. Aggarwal and J.-L. Vasse, *Org. Lett.*, 2003, **5**, 3987–3990; (g) A. Solladie-Cavallo, M. Roje, R. Welter and V. Sunjić, *J. Org. Chem.*, 2004, **69**, 1409–1412.
- For ammonium ylide mediated cyclopropanations see: (a) M. J. Gaunt and C. C. C. Johansson, *Chem. Rev.*, 2007, **107**, 5596–5605; (b) C. C. C. Johansson, N. Bremeyer, S. V. Ley, D. R. Owen, S. C. Smith and M. J. Gaunt, *Angew. Chem., Int. Ed.*, 2006, **45**, 6024–6028; (c) C. D. Papageorgiou, M. A. Cubillo de Dios, S. V. Ley and M. J. Gaunt, *Angew. Chem., Int. Ed.*, 2004, **43**, 4641–4644; (d) N. Bremeyer, S. C. Smith, S. V. Ley and M. J. Gaunt, *Angew. Chem., Int. Ed.*, 2004, **43**, 2681–2684; (e) C. D. Papageorgiou, S. V. Ley and M. J. Gaunt, *Angew. Chem., Int. Ed.*, 2003, **42**, 828–831; (f) S. Kojima, K. Hiroike and K. Ohkata, *Tetrahedron Lett.*, 2004, **45**, 3565–3568; (g) S. Yamada, J. Yamamoto and E. Ohta, *Tetrahedron Lett.*, 2007, **48**, 855–858; (h) N. Kanomata, R. Sakaguchi, K. Sekine, S. Yamashita and H. Tanaka, *Adv. Synth. Catal.*, 2010, **352**, 2966–2978.
- For benzylic ammonium ylide mediated epoxidations see: (a) T. Kimachi, H. Kinoshita, K. Kusaka, Y. Takeuchi, M. Aoe and M. Ju-ichi, *Synlett*, 2005, 842–844; (b) H. Kinoshita, A. Ihoriya, M. Ju-ichi and T. Kimachi, *Synlett*, 2010, 2330–2334; (c) R. Robiette, M. Conza and V. K. Aggarwal, *Org. Biomol. Chem.*, 2006, **4**, 621–623.



- 7 For amide-stabilised ammonium ylide mediated epoxidations see: (a) M. Waser, R. Herchl and N. Müller, *Chem. Commun.*, 2011, **47**, 2170–2172; (b) R. Herchl, M. Stiftinger and M. Waser, *Org. Biomol. Chem.*, 2011, **9**, 7023–7027.
- 8 For cyano-stabilised ammonium ylide mediated epoxidations see: (a) A. Kowalkowska, D. Sucholbiak and A. Jonczyk, *Eur. J. Org. Chem.*, 2005, 925–933; (b) A. Jonczyk and A. Konarska, *Synlett*, 1999, 1085–1087; (c) A. Alex, B. Larmanjat, J. Marrot, F. Couty and O. David, *Chem. Commun.*, 2007, 2500–2502.
- 9 For ammonium ylide-mediated aziridinations see: (a) L. D. S. Yadav, R. Kapoor and Garima, *Synlett*, 2009, 3123–3126; (b) S. Aichhorn, G. N. Gururaja, M. Reisinger and M. Waser, *RSC Adv.*, 2013, **3**, 4552–4557.
- 10 For selected reviews see: (a) A. H. Hoveyda, D. A. Evans and G. C. Fu, *Chem. Rev.*, 1993, **93**, 1307–1370; (b) P. Arya and H. Qin, *Tetrahedron*, 2000, **56**, 917–947; (c) D. A. Evans and J. T. Shaw, *Actual. Chim.*, 2003, 35–38; (d) D. J. Ager, I. Prakash and D. R. Schaad, *Chem. Rev.*, 1996, **96**, 835–875.
- 11 For chiral amide containing sulfonium ylides in epoxidations see: (a) F. Sarabia, S. Chammaa, M. Garica-Castro and F. Martin-Galvez, *Chem. Commun.*, 2009, 5763–5765; (b) D. M. Aparicio, J. L. Teran, D. Gnecco, A. Galindo, J. R. Juarez, M. L. Orea and A. Mendoza, *Tetrahedron: Asymmetry*, 2009, **20**, 2764–2768; (c) F. Sarabia, C. Vivar-Garcia, M. Garica-Castro and J. Martin-Ortiz, *J. Org. Chem.*, 2011, **76**, 3139–3150; (d) D. M. Aparicio, D. Gnecco, J. R. Juarez, M. L. Orea, A. Mendoza, N. Waksman, R. Salazar, M. Fores-Alamo and J. L. Teran, *Tetrahedron*, 2012, **68**, 10252–10256; (e) P. G. Gordillo, D. M. Aparicio, M. Flores, A. Mendoza, L. Orea, J. R. Juarez, G. Huelgas, D. Gnecco and J. L. Teran, *Eur. J. Org. Chem.*, 2013, 5561–5565.
- 12 Seminal report describing the use of these auxiliaries: D. A. Evans, J. Bartroli and T. L. Shih, *J. Am. Chem. Soc.*, 1981, **103**, 2127–2129.
- 13 Seminal reports about the use as a chiral auxiliary: (a) M. Larcheveque, E. Ignatova and T. Cuvigny, *Tetrahedron Lett.*, 1978, **19**, 3961–3964; (b) A. G. Myers, B. H. Yang, H. Chen and J. L. Gleason, *J. Am. Chem. Soc.*, 1994, **116**, 9361–9362.
- 14 These conditions were found to be useful for the parent epoxidation of ammonium salts **1** with different aldehydes: S. Aichhorn, *On the Diastereoselective Synthesis of Aziridines and Epoxides using Nitrogen Ylides*, Diploma Thesis, Institute of Organic Chemistry, Johannes Kepler University Linz, 2012.
- 15 (a) CCDC 1023711 (Cambridge Crystallographic Data Centre, <http://www.ccdc.cam.ac.uk>) contains the supplementary crystallographic data for **17a** (b) CCDC 1030561 (Cambridge Crystallographic Data Centre, <http://www.ccdc.cam.ac.uk>) contains the supplementary crystallographic data for **19f**.
- 16 Calculations were carried out at the B3LYP-D3/6-311+G** (CH₂Cl₂)//B3LYP-D3/6-31G*(CH₂Cl₂) level using Jaguar, version 8.0, Schrodinger, LLC, New York, NY, 2011. Reported energies are free energies (kcal mol⁻¹) relative to reactants in their most stable conformation.
- 17 Systematic more general computational studies delineating ammonium ylide mediated epoxidation reactions will be an integral part of our future research in this field.
- 18 For computational studies showing preference of (*Z*)-configuration in sulfonium and phosphonium ylides see ref. 3f and: R. Robiette, J. Richardson, V. K. Aggarwal and J. N. Harvey, *J. Am. Chem. Soc.*, 2005, **127**, 13468–13469.
- 19 The aziridines were generally less stable than the epoxides especially at higher temperature or under acidic conditions.
- 20 S. Kanemasa and K. Onimura, *Tetrahedron*, 1992, **48**, 8631–8644.
- 21 R. J. R. Lumby, P. M. Joensuu and H. W. Lam, *Tetrahedron*, 2008, **64**, 7729–7740.

