Cite this: *Org. Biomol. Chem.*, 2015, **13**, 3787

Gold-catalyzed allene cycloisomerization for pyrrole synthesis: towards highly fluorinated BODIPY dyes†

Linda Lempke,^a Tobias Fischer,^b Jérémy Bell,^b Werner Kraus,^b Knut Rurack^{*b} and Norbert Krause^{*a}

A novel synthetic strategy toward highly fluorinated BODIPY dyes with exceptional photostabilities relying on sustainable gold catalysis has been developed. A key to the tailored pyrrole precursors is the gold catalysis performed in ionic liquids as the reaction medium, allowing a facile recycling of the catalysts. The dyes prepared are well-matching with the spectral windows of popular rhodamine dyes and possess high brightness while showing a distinctly higher photostability than the rhodamines especially in aprotic solvents.

Received 26th December 2014,
Accepted 5th February 2015

DOI: 10.1039/c4ob02671c

www.rsc.org/obc

Introduction

Sustainable metal catalysis has attracted general attention during the last few years. Meanwhile also for gold catalysis powerful sustainable methods are known.¹ The use of ionic liquids (ILs) or micelles as the reaction medium enables the stabilization of catalysts, preventing their undesired reduction to metallic gold. Previously, the unique stability and recyclability of these catalyst solutions were demonstrated.²

Novel sustainable synthetic routes are often a key to success in advancing compounds of a certain class of specialty chemicals, for instance with unprecedented substitution patterns or improved reaction yields. In this regard, boron-dipyrromethene (BODIPY) dyes as one of the fastest growing classes of dyes³ are very attractive targets. The popularity of BODIPY dyes is rooted in their outstanding absorption and fluorescence properties and high photo- and chemical stabilities, having resulted in a wide variety of applications. Besides bio-imaging,⁴ fluorescent indicators,⁵ photodynamic therapy,⁶ and organic photovoltaics,⁷ highly fluorinated BODIPYs have been recently found to constitute a very promising class of multifunctional labeling agents.⁸ One of the bottlenecks for novel

BODIPYs is the availability of tailor-made pyrroles which constitute their core building blocks.

The expertise of our groups in sustainable metal catalysis² and the development of high-performance dyes⁹ prompted us to embark on a collaborative approach toward new fluorinated BODIPY dyes **4** by gold-catalyzed cycloisomerization of α -aminoallenes **1** to 2,5-dihydropyrroles **2** in ionic liquids (Scheme 1). Although fluorinated BODIPYs possess many promising features, only a very limited number of fluorinated BODIPYs is known in the literature.^{8,10} Furthermore, the synthesis of pyrroles by cyclization reactions allows extensive structural variation and represents a valuable alternative compared to common strategies.¹¹

Scheme 1 Synthesis of BODIPY dyes.

^aOrganic Chemistry, Dortmund University of Technology, Otto-Hahn-Str. 6, D-44227 Dortmund, Germany. E-mail: norbert.krause@tu-dortmund.de; Fax: (+49 231 755 3884

^bDept. 1 Analytical Chemistry; Reference Materials, BAM Federal Institute for Materials Research and Testing, Richard-Willstätter-Str. 11, D-12489 Berlin, Germany. E-mail: knut.rurack@bam.de; Fax: (+49 30 8104 1157

†Electronic supplementary information (ESI) available: Full experimental details, including X-ray structure analysis, optical spectroscopy, computational studies, and NMR spectra. CCDC 1040186 and 1040187. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c4ob02671c

Results and discussion

The cycloisomerization was carried out in ionic liquids (Tables 1–3). Different gold catalysts and ionic liquids were examined for this purpose (Scheme 2).

The cyclization conditions were optimized with allene **1a** and cationic gold catalyst **A** (Table 1). Best results were achieved with [BMIM][PF₆] and [BMIM][HSO₄].

With 1 mol% of catalyst **A** in [BMIM][PF₆], complete conversion was observed after 24 h, while 2 mol% led to a shorter reaction time of 2 h (entries 1 & 2). The acidic ILs [BMIM][HSO₄] and [EMIM][HSO₄] were used at 60 °C to attain a lower

Table 1 Influence of ionic liquids on the gold-catalyzed cycloisomerization of α -aminoallene **1a**

Entry	Ionic liquid	A (mol%)	T (°C)	t (h)	Conv. ^a (%)
1	[BMIM][PF ₆]	1	25	24	100
2	[BMIM][PF ₆]	2	25	2	100
3	[BMIM][PF ₆]	2	60	0.25	100
4	[BMIM][HSO ₄]	2	60	0.25	100
5	[EMIM][HSO ₄]	2	60	1	100
6	[BMIM][HSO ₄]	—	60	24	0
7	[EMIM][HSO ₄]	—	60	24	0
8	[BMIM][BF ₄]	2	25	24	0
9	[EMIM][BF ₄]	2	25	24	0

^a Conversion was determined by ¹H-NMR spectroscopy.

Table 2 Influence of the gold catalyst on the cycloisomerization of α -aminoallene **1a** in [BMIM][PF₆]

Entry	[Au]	Additive	t	Conv. ^a (%)
1	A	—	2 h	100
2	AuBr ₃	—	48 h	30
3	Ph ₃ PAuNTf ₂	—	1 h	100
4 ^b	Ph ₃ PAuNTf ₂	—	5 min	100
5	Ph ₃ PAuCl	AgSbF ₆	1 h	100
6	Ph ₃ PAuCl	—	24 h	0
7	B	AgSbF ₆	30 min	100
8	B	—	24 h	0
9 ^b	B	AgSbF ₆	5 min	100
10 ^c	B	AgSbF ₆	10 min	100
11	—	AgSbF ₆	24 h	20

^a Conversion was determined by ¹H-NMR spectroscopy. ^b Toluene (ca. 20 vol%) was added. ^c Reaction temperature of 50 °C.

Table 3 Recycling of the catalyst/solvent system

Entry	Allene	Run	Yield ^a (%)
1	1a	1	91
2	1a	2	91
3	1a	3	98
4	1a	4	95
5	1a	5	80 ^b
6	1b	1	90
7	1b	2	92

^a Isolated yield. ^b 12% of **1a** was recovered.

Scheme 2 Gold catalysts and ionic liquids used in this work.

viscosity. Short reaction times of 15 min and 1 h were obtained (entries 4 & 5), which are comparable to [BMIM][PF₆] at the same temperature (entry 3). The acidic ILs do not mediate the cycloisomerization in the absence of the gold catalyst (entries 6 & 7).^{12,13} In [BMIM][BF₄] and [EMIM][BF₄] no reaction was observed after 24 h at RT (entries 8 & 9). This may be due to a limited solubility of the gold catalyst in these more hydrophilic ionic liquids.

Allene **1a** and ionic liquid [BMIM][PF₆] were used for screening the gold catalyst (Table 2). Except for AuBr₃ (entry 2), all catalysts examined gave short reaction times between 30 min and 2 h (entries 1, 3, 5 & 7). Both precatalyst **B** and Ph₃PAuCl were transformed into the corresponding cationic gold species with AgSbF₆. Without the addition of a silver salt no reaction was observed (entries 6 & 8), whereas the silver salt alone induces only a very slow cyclization (entry 11). Beside the possibility of warming up (entry 10), the addition of a certain amount of toluene to form a solvent film led to decreased reaction times of 5 min (entries 4 vs. 3, 9 vs. 7). This acceleration is probably caused by the two-phasic character of the reaction mixture. While the catalyst/solvent system is a viscous solution, the allene is a solid and affords an emulsion during the reaction. The toluene seems to act as a phase-transfer agent. These conditions were chosen for recycling experiments (Table 3).

Previous investigations with AuBr₃ in [BMIM][PF₆] showed an extremely low catalyst leaching using hexane as an extraction solvent.^{2a} The pyrrolines **2** are not soluble in nonpolar solvents like hexane. Unfortunately, polar solvents like Et₂O cause a dramatic catalyst leaching. Therefore, the recycling was tested for **1a** with toluene as the extraction solvent and Ph₃PAuNTf₂ as the gold catalyst. Excellent reactivity was

Fig. 1 ORTEP representation of **4a** (left) and **4b** (right) with atomic labeling shown with 30% probability displacement ellipsoids.

rotational freedom. The comparatively small spectral differences between **4a** and **4b** despite the introduction of two additional CF₃ groups in the first one are tentatively ascribed to the enhanced steric hindrance in the case of 3,5-phenyl substitution on the BODIPY core.¹⁷ That results in considerable (average) torsion angles around the C1–C1' bond ($\langle\theta_{C2-C1-C1'-C2'}\rangle = 39^\circ$ for **4a** and 38° for **4b** (for the atom numbering pattern, see Scheme 1, bottom panel) and hence reduced electronic coupling between the BODIPY core and phenyl rings, diminishing the electron withdrawing effect of these substituents. These experimental findings are well reflected by quantum chemical calculations employing the density functional theory (DFT) method, revealing differences in the S₁←S₀ transition in the gas phase of only ca. 5 nm (Table S2, ESI†). The latter are also the only oscillator-strong transitions in the 350–500 nm region for both dyes, involving HOMO and LUMO which are both centered almost exclusively on the 3,5-diphenyl-dipyrrin fragments for **4a,b** (Fig. S2, ESI†). Furthermore, with the corresponding triplet transitions being shifted for ca. 1.25 eV, the theoretical studies fully support the favorable spectroscopic properties found experimentally.

Dyes **4a** and **4b** absorb and emit in the green visible range, matching well the spectral region of prominent rhodamine dyes such as rhodamine 101, 6G, B or TRITC¹⁸ and fitting perfectly to the output of green laser sources (e.g., the prominent 532 or 543 nm lasers, Fig. 3). Compared to rhodamines the zwitterionic yet net uncharged BODIPYs are commonly well soluble in organic solvents across the entire polarity range,^{3,19} broadening the areas of applications and offering potent alternatives to the search for hydrophobic rhodamine derivatives.²⁰ The most important feature of the newly synthesized fluorinated BODIPYs **4a,b** with regard to application is their excellent photostability, in particular when used in aprotic solvents. Fig. 4 exemplarily combines the photobleaching curves of **4a,b** and rhodamine 101 upon irradiation at 532 nm.

Fig. 4 Photobleaching curves for irradiation at 532 nm (1 W laser power) in acetonitrile (**4a**: orange, **4b**: blue, rhodamine 101: red) and ethanol (**4a**: magenta, **4b**: green, rhodamine 101: black).

Conclusions

In the work presented here, two highly fluorinated and photostable BODIPY dyes with excellent spectroscopic properties have been prepared *via* gold catalysis in ionic liquids. While the use of sustainable gold catalysis is often strictly limited to the applied system, the possibility of recycling the catalyst solution was extended to more challenging allenes. Here, [BMIM][PF₆] was found to be an excellent solvent system for gold catalysis and recycling. The fluorinated dyes possess an enormous potential for further applications such as functional group labeling or the staining of lipophilic compartments, for instance in cell imaging. Current work in our laboratories is directed toward both the synthesis of other functional BODIPYs and the application of the title dyes.

Acknowledgements

Financial support by SusChemSys (which is co-financed by the Regional Development Fund (Investing in Your Future) of the European Union and the state of North Rhine-Westphalia), the European Metrology Research Programme (EMRP) (project IND15 SurfChem) and the Adolf-Martens-Fund is gratefully acknowledged. The EMRP is jointly funded by the EMRP participating countries within EURAMET and the European Union.

Notes and references

- (a) X. Liu, L. He, Y.-M. Liu and Y. Cao, *Acc. Chem. Res.*, 2014, **47**, 793; (b) C. Winter and N. Krause, *Green Chem.*, 2009, **11**, 1309; (c) C. Hviid Christensen and J. K. Nørskov, *Science*, 2010, **327**, 278.
- (a) Ö. Aksin and N. Krause, *Adv. Synth. Catal.*, 2008, **350**, 1106; (b) S. R. K. Minkler, B. H. Lipshutz and N. Krause, *Angew. Chem., Int. Ed.*, 2011, **50**, 7820; (c) S. R. K. Minkler, N. A. Isley, D. J. Lippincott, N. Krause and B. H. Lipshutz, *Org. Lett.*, 2014, **16**, 724; (d) X. Liu, Z. Pan, X. Shu, X. Duan and Y. Liang, *Synlett*, 2006, 1962; (e) X. Moreau, A. Hours, L. Fensterbank, J.-P. Goddard, M. Malacria and S. Thorimbert, *J. Organomet. Chem.*, 2009, **594**, 561.
- (a) G. Ulrich, R. Ziessel and A. Harriman, *Angew. Chem. Int. Ed.*, 2008, **47**, 1184; (b) A. Loudet and K. Burgess, *Chem. Rev.*, 2007, **107**, 4891.
- K. Umezawa, D. Citterio and K. Suzuki, *Anal. Sci.*, 2014, **30**, 327.
- N. Boens, V. Leen and W. Dehaen, *Chem. Soc. Rev.*, 2012, **41**, 1130.
- A. Kamkaew, S. H. Lim, H. B. Lee, L. V. Kiew, L. Y. Chung and K. Burgess, *Chem. Soc. Rev.*, 2013, **42**, 77.
- A. Bessette and G. S. Hanan, *Chem. Soc. Rev.*, 2014, **43**, 3342.
- (a) M. Hecht, T. Fischer, P. Dietrich, W. Kraus, A. Descalzo, W. Unger and K. Rurack, *ChemistryOpen*, 2013, **2**, 25; (b) T. Fischer, P. M. Dietrich, C. Streeck, S. Ray, A. Nutsch,

- A. Shard, B. Beckhoff, W. E. S. Unger and K. Rurack, *Anal. Chem.*, 2015, DOI: 10.1021/ac503850f.
- 9 (a) P. Ashokkumar, H. Weißhoff, W. Kraus and K. Rurack, *Angew. Chem. Int. Ed.*, 2014, **53**, 2225; (b) Y.-H. Yu, A. B. Descalzo, Z. Shen, H. Röhr, Q. Liu, Y.-W. Wang, M. Spieles, Y.-Z. Li, K. Rurack and X.-Z. You, *Chem. – Asian J.*, 2006, **1**, 176; (c) M. Kollmannsberger, K. Rurack, U. Resch-Genger and J. Daub, *J. Phys. Chem. A*, 1998, **102**, 10211.
- 10 For pentafluorophenyl-substituted BODIPYs see: (a) G. Vives, C. Giansante, R. Bofinger, G. Raffy, A. Del Guerzo, B. Kauffmann, P. Batat, G. Jonusauskas and N. D. McClenaghan, *Chem. Commun.*, 2011, **47**, 10425; (b) O. Galangau, C. Dumas-Verdes, R. Méallet-Renault and G. Clavier, *Org. Biomol. Chem.*, 2010, **8**, 4546. For aryl- and fluorine-substituted BODIPYs see: (c) V. Lakshmi and M. Ravikanth, *Chem. Phys. Lett.*, 2013, **564**, 93; (d) V. Lakshmi and M. Ravikanth, *J. Org. Chem.*, 2011, **76**, 8466; (e) A. Burghart, H. Kim, M. B. Welch, L. H. Thoresen, J. Reibenspies and K. Burgess, *J. Org. Chem.*, 1999, **64**, 7813; (f) L. H. Thoresen, H. Kim, M. B. Welch, A. Burghart and K. Burgess, *Synlett*, 1998, **11**, 1276.
- 11 (a) V. Estévez, M. Villacampa and J. C. Menéndez, *Chem. Soc. Rev.*, 2010, **39**, 4402; (b) V. Estévez, M. Villacampa and J. C. Menéndez, *Chem. Soc. Rev.*, 2014, **43**, 4633. For 2,4-substituted pyrroles see: (c) Z. Chen, B. Lu, Z. Ding, K. Gao and N. Yoshikai, *Org. Lett.*, 2013, **15**, 1966; (d) H.-Y. Wang, D. S. Mueller, R. M. Sachwani, H. N. Londino and L. L. Anderson, *Org. Lett.*, 2010, **12**, 2290–2293; (e) L. Meng, K. Wu, C. Liua and A. Lei, *Chem. Commun.*, 2013, **49**, 5853; (f) Z. Shi, M. Suri and F. Glorius, *Angew. Chem., Int. Ed.*, 2013, **52**, 4892.
- 12 N. Krause, M. Laux and A. Hoffmann-Röder, *Tetrahedron Lett.*, 2000, **41**, 9613.
- 13 N. Krause, A. Hoffmann-Röder and J. Canisius, *Synthesis*, 2012, **12**, 1759.
- 14 T. Xu, X. Mu, H. Peng and G. Liu, *Angew. Chem., Int. Ed.*, 2011, **50**, 8176.
- 15 (a) C. L. Picou, E. D. Stevens, M. Shah and J. H. Boyer, *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.*, 1990, **46**, 1148; (b) D.-C. Wang, C. He, J.-L. Fan, W.-W. Huang and X.-J. Peng, *Acta Crystallogr., Sect. E: Struct. Rep. Online*, 2007, **63**, o2900; (c) A. Wakamiya, T. Murakami and S. Yamaguchi, *Chem. Sci.*, 2013, **4**, 1002.
- 16 Y.-W. Wang, A. B. Descalzo, Z. Shen, X.-Z. You and K. Rurack, *Chem. – Eur. J.*, 2010, **16**, 2887.
- 17 In addition it has to be kept in mind that the phenyl ring attenuates the effects of a CF₃ group, *i.e.*, the step from 4-CF₃C₆H₅ to 3,5-(CF₃)₂C₆H₄ has much less influence in terms of for instance the Hammett constant than the step from a *p*-CF₃ to two *m*-CF₃ groups, *cf. e.g.* G. A. Olah, A. L. Berrier, M. Arvanaghi and G. K. S. Prakash, *J. Am. Chem. Soc.*, 1981, **103**, 1122; Y. Borguet, X. Sauvage, G. Zaragoza, A. Demonceau and L. Delaude, *Organometallics*, 2010, **29**, 6675.
- 18 (a) P. C. Beaumont, D. G. Johnson and B. J. Parsons, *J. Chem. Soc., Faraday Trans.*, 1993, **89**, 4185; (b) H. Brismar, O. Trepte and B. Ulfhake, *J. Histochem. Cytochem.*, 1995, **43**, 699.
- 19 Another advantage of BODIPY dyes is the straightforwardness of tailoring water solubility through substitution of the fluorine atoms of the BF₂ group, see, *e.g.*: (a) A. M. Courtis, S. A. Santos, Y. Guan, J. A. Hendricks, B. Ghosh, D. M. Szantai-Kis, S. A. Reis, J. V. Shah and R. Mazitschek, *Bioconjugate Chem.*, 2014, **25**, 1043; (b) S.-L. Niu, G. Ulrich, P. Retailleau, J. Harrowfield and R. Ziessel, *Tetrahedron Lett.*, 2009, **50**, 3840.
- 20 L. F. Mottram, S. Forbes, B. D. Ackley and B. R. Peterson, *Beilstein J. Org. Chem.*, 2012, **8**, 2156.