



Cite this: *Org. Biomol. Chem.*, 2015, **13**, 4706

## Investigations of possible prodrug structures for 2-(2-mercaptophenyl)tetrahydropyrimidines: reductive conversion from anti-HIV agents with pyrimidobenzothiazine and isothiazolopyrimidine scaffolds†

Shiho Okazaki,<sup>a</sup> Shinya Oishi,<sup>\*a</sup> Tsukasa Mizuhara,<sup>a</sup> Kazuya Shimura,<sup>b</sup> Hiroto Murayama,<sup>b</sup> Hiroaki Ohno,<sup>a</sup> Masao Matsuoka<sup>b</sup> and Nobutaka Fujii<sup>\*a</sup>

3,4-Dihydro-2*H*,6*H*-pyrimido[1,2-*c*][1,3]benzothiazin-6-imine (PD 404182) and 3,4-dihydro-2*H*-benzo[4,5]isothiazolo[2,3-*a*]pyrimidine are the heterocyclic antiretroviral agents against human immunodeficiency virus type 1 (HIV-1) infection. On the basis of similar structure–activity relationships of anti-HIV activities toward the early-stage of viral infection between these unique scaffolds, the transformations under the bioassay conditions were investigated. The distinctive S–N bond in the isothiazolopyrimidine scaffold was immediately cleaved under reductive conditions in the presence of GSH to generate a thiophenol derivative. A similar rapid conversion of PD 404182 into the same thiophenol derivative was observed, suggesting that pyrimidobenzothiazine and isothiazolopyrimidine scaffolds may work as prodrug forms of the common bioactive thiophenol derivatives.

Received 12th February 2015,  
Accepted 9th March 2015

DOI: 10.1039/c5ob00301f

[www.rsc.org/obc](http://www.rsc.org/obc)

## Introduction

PD 404182 (**1a**, 3,4-dihydro-2*H*,6*H*-pyrimido[1,2-*c*][1,3]benzothiazin-6-imine)<sup>1–4</sup> is an antiviral agent against human hepatitis C virus (HCV),<sup>5,6</sup> HIV,<sup>6–10</sup> simian immunodeficiency virus (SIV),<sup>6</sup> and herpes simplex virus (HSV) (Fig. 1).<sup>10</sup> 3,4-Dihydro-2*H*-benzo[4,5]isothiazolo[2,3-*a*]pyrimidine **2a** with a unique heterocyclic scaffold also exhibits potent anti-HIV activity ( $EC_{50} = 0.29 \pm 0.09 \mu M$ ) (Fig. 1).<sup>11</sup> In our recent studies, we

revealed that modification of the 9-position of PD 404182 derivatives with hydrophobic aryl groups improved the anti-HIV activity.<sup>7–9</sup> For example, the introduction of an aryl (**1b**, **1c** and **1f**) or bromo (**1d**) substituent at the 9-position led to a two- to three-fold increase in the anti-HIV activity compared with compound **1a** (Table 1). This improvement was duplicated by the modification of benzo[4,5]isothiazolo[2,3-*a*]pyrimidine derivatives at the 8-position (**2b**, **2c**, **2d** and **2f**), which is the equivalent position of the 9-position in pyrimido[1,2-*c*][1,3]benzothiazin-6-imine. Similarly, introduction of acetamido group at the equivalent positions resulted in a complete loss of the anti-HIV activity of both the PD 404182 derivative (**1e**) and isothiazolopyrimidine derivative (**2e**). Furthermore, a complete loss of anti-HIV activity was observed by the introduction of the 8-bromo and 7-bromo groups to the parent compounds (**1g** and **2g**, respectively). These similar structure–activity relationships implied that the same target molecule(s) or mechanism(s) of action could possibly be involved in the anti-HIV activities of PD 404182 and the isothiazolopyrimidine derivatives.

In addition, heterocyclic compounds **1a** and **2a** both inhibited early-stage HIV infection including virus attachment and membrane fusion to host cells as exemplified in comparison with DS 5000 (adsorption inhibitor)<sup>12</sup> and enfuvirtide (fusion inhibitor).<sup>13</sup> Although it was suggested that the antiviral effects of compound **1a** are derived from its virucidal effect against viral particles,<sup>14</sup> its antiviral mechanism of action has not yet




Fig. 1 Structures of PD 404182 (**1a**) and benzo[4,5]isothiazolo[2,3-*a*]pyrimidine (**2a**).

<sup>a</sup>Graduate School of Pharmaceutical Sciences, Kyoto University, Sakyo-ku, Kyoto 606-8501, Japan. E-mail: soishi@pharm.kyoto-u.ac.jp, nfujii@pharm.kyoto-u.ac.jp; Fax: +81-75-753-4570; Tel: +81-75-753-4561

<sup>b</sup>Institute for Virus Research, Kyoto University, Sakyo-ku, Kyoto, 606-8507, Japan

† Electronic supplementary information (ESI) available. CCDC 1037500. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c5ob00301f



**Table 1** Comparison of the structure–activity relationships of PD 404182 derivatives and benzo[4,5]isothiazolo[2,3-*a*]pyrimidines

Cmpd		EC <sub>50</sub> <sup>a,7,8</sup> (μM)	Cmpd	EC <sub>50</sub> <sup>b,11</sup> (μM)	
<b>1a</b> (PD 404182)		0.44 ± 0.08 <sup>c</sup>	<b>2a</b>	R = H 0.29 ± 0.09	
<b>1b</b>	R = Ph	0.24 ± 0.04	<b>2b</b>	R = Ph 0.30 ± 0.10	
<b>1c</b>	R = <i>m</i> -anisyl	0.15 ± 0.05	<b>2c</b>	R = <i>m</i> -anisyl 0.10 ± 0.03	
<b>1d</b>	R = Br	0.25 ± 0.09 <sup>c</sup>	<b>2d</b>	R = Br 0.22 ± 0.07	
<b>1e</b>	R = NHAc	>10	<b>2e</b>	R = NHAc >10	
<b>1f</b>		0.15 ± 0.03	<b>2f</b>	R = NHAc 0.16 ± 0.02	
<b>1g</b>		>10 <sup>c</sup>	<b>2g</b>		>10 <sup>d</sup>

<sup>a</sup> EC<sub>50</sub> values represent the concentration of the compound required to inhibit the HIV-1 infection by 50%. The data were obtained from three independent experiments by the MAGI assay. <sup>b</sup> EC<sub>50</sub> values were obtained by the NCK assay. <sup>c</sup> EC<sub>50</sub> values were obtained by the NCK assay, compound **1a**: 0.30 ± 0.06 μM; compound **1d**: 0.14 ± 0.02 μM; compound **1g**: >10 μM. <sup>d</sup> Cytotoxicity was observed at 10 μM.

been sufficiently detailed. In the current study, we investigated the chemical transformations of pyrimido[1,2-*c*][1,3]benzothiazin-6-imine derivatives **1** and benzo[4,5]isothiazolo[2,3-*a*]pyrimidine derivatives **2** under the bioassay conditions.


## Results and discussion

### Synthesis

PD 404182 derivatives **1** and benzo[4,5]isothiazolo[2,3-*a*]pyrimidine derivatives **2** were synthesized according to the protocols in our previous studies (Scheme 1).<sup>7,8,11</sup> 2-(2-Mercaptophenyl)tetrahydropyrimidine derivatives **4** were obtained by trifluoroacetic acid (TFA)-mediated ethanalysis of the imino group in pyrimido[1,2-*c*][1,3]benzothiazin-6-imines **1**. The alternative route *via* alkaline hydrolysis of pyrimido[1,2-*c*][1,3]-benzothiazin-6-thione **3** also generated thiophenol derivatives **4**. Some thiophenol derivatives **4** could be isolated as stable crystals, such as 2-(2-mercaptophenyl)-, 2-(4-bromo-2-mercaptophenyl)- or 2-(3-bromo-2-mercaptophenyl)-tetrahydropyrimidine (**4a**, **d**, **g**). Thioanisole derivative **7** was obtained by the oxidative amidination of the corresponding benzaldehyde **5**.<sup>15,16</sup> Sulfuric acid-mediated deprotection of *N*-tosylated tetrahydropyrimidine **8**, derived from benzaldehyde **6** by the identical protocol,<sup>7</sup> afforded aniline derivative **9**.

### Identification of the inactive ingredient by transformation from PD 404182 in aqueous media


Recently, Chamoun-Emanuelli *et al.* reported that the antiviral activity of PD 404182 (**1a**) disappeared during storage under some conditions (for example, in Dulbecco's phosphate-buffered saline (pH 7) at 42 °C).<sup>10</sup> During our studies on PD



**Scheme 1** Synthesis of benzo[4,5]isothiazolo[2,3-*a*]pyrimidine and 2-phenyl-1,4,5,6-tetrahydropyrimidine derivatives. Reagents and conditions: (a) TFA, CHCl<sub>3</sub>-EtOH or EtOH, rt; (b) NaOH, MeOH-H<sub>2</sub>O, reflux; (c) PIDA, CHCl<sub>3</sub> and/or EtOH, rt; (d) 1,3-propanediamine, I<sub>2</sub>, K<sub>2</sub>CO<sub>3</sub>, *t*-BuOH, 70 °C; (e) R' = NHTs (6) or R' = NHTs (8), conc. H<sub>2</sub>SO<sub>4</sub>, 100 °C.

404182 derivatives, we also found that the anti-HIV activity from the stock solution of **1a** in DMSO was often attenuated. X-ray crystallography of the predominant component from **1a** revealed that the resulting product was 3,4-dihydro-2*H*,6*H*-





**Fig. 2** Transformations of PD 404182 under assay conditions and under reductive conditions in the presence of GSH at 37 °C. EC<sub>50</sub> values represent the concentration of the compound required to inhibit the HIV-1 infection by 50%. The data were obtained from three independent experiments by the NCK assay.


benzo[*e*]pyrimido[2,1-*b*][1,3]thiazin-6-one **10** (Fig. 2). An apparent rearrangement of the cyclic amidine substructure in **1a** provided pyrimidothiazinone **10** with no anti-HIV activity. In contrast, benzo[4,5]isothiazolo[2,3-*a*]pyrimidine **2a** was stable in acidic and basic solutions, and the assay medium containing fetal calf serum.

#### Transformation of PD 404182 and benzo[4,5]isothiazolo[2,3-*a*]pyrimidine derivatives under reductive conditions in the presence of GSH

We investigated the conversion of PD 404182 (**1a**) in the presence of high glutathione (GSH) concentrations that mimic the


intracellular environment.<sup>17</sup> Interestingly, the imino group in **1a** decomposed in 10 mM GSH solution at 37 °C, and was converted into thiophenol **4a** within 1 h (Fig. 2 and 3).<sup>18</sup> This thiophenol derivative **4a** exhibited equipotent anti-HIV activity as the parent compound **1a** (EC<sub>50</sub> = 0.32 ± 0.06 μM). The mechanism of conversion from **1a** into thiophenol **4a** could be *via* the release of cyanylated glutathione, which was detected by ESI-MS analysis.<sup>19</sup>

Our previous structure–activity relationship study of **1a** demonstrated that the 6-imino group and the 7-sulfur atom in **1a** were indispensable for the anti-HIV activity (Table 2).<sup>7</sup> For example, substitution of the imino group in compound **1a** with a carbonyl (**13**) or thiocarbonyl (**3a**) group resulted in a significant decrease or loss of activity. Neither pyrimido[1,2-*c*]-[1,3]benzoxazines (**14a–c**) nor pyrimido[1,2-*c*]quinazolines (**15a–c**) exhibited anti-HIV activity (Table 2). In the presence of GSH, the carbonyl (**13**) and thiocarbonyl (**3a**) derivatives were slowly converted to compound **4a** in 26 and 15% yield, respectively, after 24 h (Table 2, Fig. 3). This is in contrast with the rapid and quantitative conversion from **1a** within 1 h, indicating that a more reactive imino group in **1a** was a prerequisite for the efficient generation of **4a** with potent anti-HIV activity. In the absence of GSH, compound **13** was converted into **4a** in only 5% yield for 24 h, and **3a** did not produce **4a**, indicating that degradation of **13** and **3a** was mainly mediated by GSH.<sup>20</sup> Compound **14a** was unstable in the presence or absence of GSH to give a cyclic carbamate **14b** and a phenol derivative **16**,<sup>7,21</sup> which showed no anti-HIV activity. The carbamate **14b** and thiocarbamate **14c** were also degraded into an inactive phenol **16** in 78 and 71% yield, respectively, in GSH solution over 24 h. The pyrimido[1,2-*c*]quinazolines (**15a–c**) were stable under identical conditions, and thus, did not produce the possible aniline **9**. In light of these results, the absent or diminished anti-HIV activity of **3a** and **13–15** is likely attributable to decreased conversion to the potent thiophenol **4a** or the poor anti-HIV activity of the parent compounds and ring-opened products.



**Fig. 3** Transformation of compounds **1a** (A), **13** (B) and **3a** (C) under reductive conditions in the presence of GSH. <sup>a</sup> The concentrations of compounds were calculated by HPLC analysis using previously determined calibration curves.




**Table 2** Reactions of PD 404182 derivatives under reductive conditions in the presence of GSH

Cmpd	X	EC <sub>50</sub> <sup>b,7</sup> (μM)	Conversion (time, yield <sup>c</sup> )
PD 404182 (1a)	NH	0.44 ± 0.08	4a (1 h, quant.)
13	O	8.94 ± 1.07	4a (24 h, 26%) <sup>d</sup>
3a	S	>10	4a (24 h, 15%) <sup>d</sup>
14a	NH	>10	16 + 14b <sup>e</sup>
14b	O	>10	16 (24 h, 78%)
14c	S	>10	16 (24 h, 71%)
15a	NH	>10	No reaction
15b	O	>10	No reaction
15c	S	>10	No reaction

<sup>a</sup> EC<sub>50</sub> values represent the concentration of the compound required to inhibit the HIV-1 infection by 50%. The data were obtained from three independent experiments by the NCK assay. <sup>b</sup> EC<sub>50</sub> values were obtained by the MAGI assay. <sup>c</sup> The product yields were calculated by HPLC analysis using previously determined calibration curves. <sup>d</sup> The reactions were also carried out in the absence of GSH: 5% conversion from 13 after 24 h; no reaction from 3a. <sup>e</sup> The accurate product yields of 14b and 16 by GSH-mediated conversion were not calculated because of the chemical instability of the substrate 14a.


We next focused on the isothiazolopyrimidine scaffold of 2a with a characteristic S–N covalent bond, which is contained in several anti-HIV agents against HIV-1 nucleocapsid protein 7 (NCp7).<sup>22–24</sup> We assessed the stability of compound 2a under identical GSH-containing conditions (Fig. 4). In 10 mM GSH solution at 37 °C, compound 2a was also immediately converted to a thiophenol derivative 4a with potent anti-HIV activity (<3 min). The two-fold more potent pyrimidobenzothiazine 1d or isothiazolopyrimidine 2d was also converted under identical reductive conditions into the corresponding thiophenol derivative 4d, which has similar anti-HIV activity to those of 1d and 2d.<sup>25</sup> The thiophenol derivative 4g, which was obtained by conversion from the inactive pyrimidobenzothiazine 1g or isothiazolopyrimidine 2g under the GSH-mediated conditions, exhibited no anti-HIV activity. Thus, the anti-HIV activities of the PD 404182 derivatives (1a, 1d and 1g) and iso-

**Fig. 4** Reactions of potent and inactive pyrimido[1,2-c][1,3]benzothiazin-6-imine and benzo[4,5]isothiazolo[2,3-a]pyrimidine derivatives under reductive conditions in the presence of GSH at 37 °C. EC<sub>50</sub> values represent the concentration of the compound required to inhibit the HIV-1 infection by 50%. The data were obtained from three independent experiments by the NCK assay.

thiazolopyrimidine derivatives (2a, 2d and 2g) corresponded, at least in part, to those of the thiophenol derivatives (4a, 4d and 4g). These results suggest that the tricyclic structures of 1a and 2a are the prodrug forms, which could be transformed into 4a under intracellular conditions of high GSH concentrations in host cells. Interestingly, inactive analogues with scaffolds similar to PD 404182 (1a) or compound 2a were not converted into thiophenol 4a (Fig. 5). For example, compound 10,<sup>26</sup> thioanisole derivative 7, benzo[4,5]thiazolo[3,2-a]pyrimidine 11 (the structural isomer of 2a),<sup>27</sup> and saccharin-like derivative 12<sup>28</sup> were completely stable under high concentrations of GSH for 24 hours. Therefore, the pyrimidobenzothiazine scaffold in 1a and the isothiazolopyrimidine scaffold in 2a structurally satisfied two criteria, which would make them good prodrugs of 4a.

Taken together, the structure–activity relationships and antiviral profiles were similar between the PD 404182 derivatives 1 and benzo[4,5]isothiazolo[2,3-a]pyrimidine derivatives 2. Under the reductive conditions corresponding to the intracellular environment, these scaffolds were efficiently converted to the common ring-opened thiophenol derivatives with potent anti-HIV activity. In addition, these two scaffolds inhibited not only HIV-1 infection at an early stage, but also several RNA and DNA viruses as we and others recently reported.<sup>5–8,10,11</sup> In light of these findings, it was suggested that PD 404182 and benzo[4,5]isothiazolo[2,3-a]pyrimidine derivatives may achieve their antiviral activities by acting





**Fig. 5** Reactions of PD 404182, benzo[4,5]isothiazolo[2,3-*a*]pyrimidines and the inactive analogues under reductive conditions in the presence of GSH at 37 °C. EC<sub>50</sub> values represent the concentration of the compound required to inhibit the HIV-1 infection by 50%. The data were obtained from three independent experiments by the NCK assay.

through an identical intermediate or *via* an interaction with a similar target molecule(s) or mechanism of action in an intracellular compartment. A viral target molecule(s) or mechanism (s) has not yet been determined,<sup>14</sup> but these heterocyclic compounds could conceivably regulate the defense mechanism(s) of the host cells.

## Conclusions

We have unveiled the chemical transformations of pyrimido-[1,2-*c*][1,3]benzothiazin-6-imine and benzo[4,5]isothiazolo[2,3-*a*]pyrimidine scaffolds, which yield compounds with potent antiviral activities. Under the reductive conditions meant to mimic intracellular GSH levels, the covalent S–N bond of the isothiazole substructure in **2a** was cleaved to provide the ring-opened 2-(2-mercaptophenyl)tetrahydropyrimidine **4a** with potent anti-HIV activity, which was also generated from PD 404182 (**1a**) under identical conditions. The similar structure–activity relationships and antiviral profiles between compounds **1a** and **2a** suggested that these could possibly be the prodrug forms for the common bioactive substance **4a** that binds with the potential target molecule(s) in host cells.

## Experimental section

### General

<sup>1</sup>H NMR spectra were recorded using a JEOL ECA-500 spectrometer. Chemical shifts are reported in  $\delta$  (ppm) relative to

Me<sub>4</sub>Si as an internal standard. <sup>13</sup>C NMR spectra were referenced to the residual solvent signal. Exact mass (HRMS) spectra were recorded on a JMS-HX/HX 110A mass spectrometer or Shimadzu LC-ESI-IT-TOF-MS equipment. For flash chromatography, Wakogel C-300E (Wako) and CHROMATOREX NH DM1080 (Fuji Silysys) were employed. For analytical HPLC, a Cosmosil 5C<sub>18</sub>-ARII column (4.6 × 250 mm, Nacalai Tesque, Inc.) was employed with a linear gradient of CH<sub>3</sub>CN containing 0.1% (v/v) aq. TFA at a flow rate of 1 cm<sup>3</sup> min<sup>-1</sup>, and eluting products were detected by UV at 254 nm. Preparative HPLC was performed using a Cosmosil 5C<sub>18</sub>-ARII preparative column (20 × 250 mm, Nacalai Tesque, Inc.) with a linear gradient of CH<sub>3</sub>CN containing 0.1% (v/v) aq. TFA at a flow rate of 8 cm<sup>3</sup> min<sup>-1</sup>. The purity of the compounds was determined as no less than 95% by combustion analysis or HPLC analysis. Preparation of the compounds **1–3**, **4d**, **6**, **8**, **13–15** was already reported from our laboratory.<sup>7,8,11,15</sup>

### 2-(2-Mercaptophenyl)-1,4,5,6-tetrahydropyrimidine (4a).

**Method A:** TFA (3.70 cm<sup>3</sup>, 50.0 mmol) was added to a suspension of PD 404182 (**1a**) (1.09 g, 5.00 mmol) in CHCl<sub>3</sub> (50.0 cm<sup>3</sup>) and EtOH (75.0 cm<sup>3</sup>) dropwise. After being stirred at room temperature for 1 h, the mixture was quenched with Et<sub>3</sub>N (7.20 cm<sup>3</sup>, 50.0 mmol) at 0 °C and was concentrated. The crude product was purified by flash chromatography over NH silica gel with CHCl<sub>3</sub>–MeOH (10 : 0 to 9 : 1) to give the title compound **4a** as yellow crystals (999 mg, quant.); mp 208–210 °C (from *n*-hexane–CHCl<sub>3</sub>–MeOH); IR (neat)  $\nu_{\text{max}}$ /cm<sup>-1</sup>: 3185–3092 (SH), 1590 (C=N);  $\delta$ <sub>H</sub> (500 MHz, CD<sub>3</sub>OD): 2.05–2.09 (2H, m, CH<sub>2</sub>), 3.55 (4H, t, *J* 5.7, CH<sub>2</sub>), 6.87–6.91 (1H, m, Ar), 7.06 (1H, ddd, *J*<sub>1</sub> = *J*<sub>2</sub> 8.0, *J*<sub>3</sub> 1.7, Ar), 7.28 (1H, dd, *J*<sub>1</sub> 7.7, *J*<sub>2</sub> 1.4, Ar), 7.56 (1H, dd, *J*<sub>1</sub> 8.0, *J*<sub>2</sub> 1.1, Ar);  $\delta$ <sub>C</sub> (125 MHz, DMSO-*d*<sub>6</sub>): 18.1, 38.1 (2C), 118.7, 122.8, 126.7, 128.8, 137.8, 158.9, 159.8; HRMS (FAB): *m/z* calcd for C<sub>10</sub>H<sub>13</sub>N<sub>2</sub>S [M + H]<sup>+</sup> 193.0799; found: 193.0801.

**Method B:** 3,4-dihydro-2*H*,6*H*-pyrimido[1,2-*c*][1,3]benzothiazine-6-thione **3a**<sup>15</sup> (1.15 g, 4.90 mmol) was suspended in 0.1 N NaOH in MeOH–H<sub>2</sub>O (9 : 1, 98.0 cm<sup>3</sup>). After being stirred under reflux for 12 h, the mixture was quenched with 1 N HCl until pH was adjusted to 7. The whole mixture was extracted with CHCl<sub>3</sub>–MeOH (95 : 5), and dried over MgSO<sub>4</sub>. After concentration, the resulting solid was recrystallized from Et<sub>2</sub>O–CHCl<sub>3</sub>–MeOH to give the title compound **4a** as an yellow solid (655 mg, 70%); spectral data were in good agreement with those of the compound that was synthesized using method A.

**2-(3-Bromo-2-mercaptophenyl)-1,4,5,6-tetrahydropyrimidine (4g).** By using a procedure similar to that described for the preparation of the compound **4a** from **1a**, compound **1g**<sup>7</sup> (45.0 mg, 0.152 mmol) was converted to the title compound **4g** as a colorless solid (29.6 mg, 72%); mp 260–262 °C (from *n*-hexane–CHCl<sub>3</sub>–MeOH); IR (neat)  $\nu_{\text{max}}$ /cm<sup>-1</sup>: 2895–2795 (SH), 1626 (C=N);  $\delta$ <sub>H</sub> (500 MHz, DMSO-*d*<sub>6</sub>): 1.88–1.93 (2H, m, CH<sub>2</sub>), 3.44 (4H, t, *J* 5.7, CH<sub>2</sub>), 6.58 (1H, dd, *J*<sub>1</sub> = *J*<sub>2</sub> 7.7, Ar), 7.29 (1H, d, *J* 8.0, Ar), 7.53 (1H, d, *J* 8.0, Ar), 10.85 (2H, br s);  $\delta$ <sub>C</sub> (125 MHz, DMSO-*d*<sub>6</sub>): 17.8, 38.2 (2C), 117.5, 126.2, 128.1, 132.8, 133.4, 160.1, 160.7; anal. calcd for



$C_{10}H_{11}BrN_2S$ : C, 44.29; H, 4.09; N, 10.33. Found: C, 44.12; H, 4.08; N, 10.31.

**2-[2-(Methylthio)phenyl]-1,4,5,6-tetrahydropyrimidine** (7). 2-(Methylthio)benzaldehyde 5 (90% purity, 0.430  $\text{cm}^3$ , 3.00 mmol) was reacted with propylenediamine (0.300  $\text{cm}^3$ , 3.60 mmol),  $K_2CO_3$  (1.24 g, 9.00 mmol) and  $I_2$  (952 mg, 3.75 mmol). The crude compound was purified by flash chromatography over NH silica with  $CHCl_3$ -MeOH (10:0 to 10:1) followed by recrystallization from *n*-hexane- $CHCl_3$  to give the title compound 7 as colorless crystals (294 mg, 48%): mp 110–112  $^{\circ}\text{C}$ ; IR (neat)  $\nu_{\text{max}}/\text{cm}^{-1}$ : 1621 (C=N);  $\delta_H$  (500 MHz,  $CDCl_3$ ): 1.82–1.87 (2H, m,  $CH_2$ ), 2.43 (3H, s,  $CH_3$ ), 3.45 (4H, t,  $J$  5.7,  $CH_2$ ), 4.68 (1H, br s, NH), 7.12 (1H, ddd,  $J_1 = J_2$  7.4,  $J_3$  1.1, Ar), 7.20 (1H, d,  $J$  6.9, Ar), 7.29 (1H, ddd,  $J_1 = J_2$  7.6,  $J_3$  1.5, Ar), 7.36 (dd,  $J_1$  7.4,  $J_2$  1.7, 1H, Ar);  $\delta_C$  (125 MHz,  $CDCl_3$ ): 16.0, 20.6, 42.1 (2C), 124.6, 125.5, 128.3, 129.1, 136.7, 137.0, 154.8; HRMS (ESI):  $m/z$  calcd for  $C_{11}H_{15}N_2S$  [M + H]<sup>+</sup> 207.0956; found: 207.0954.

**2-(2-Aminophenyl)-1,4,5,6-tetrahydropyrimidine** (9). A mixture of 2-[2-*N*-(*p*-toluenesulfonylamino)phenyl]-1,4,5,6-tetrahydropyrimidine 8<sup>7</sup> (494 mg, 1.50 mmol) in conc.  $H_2SO_4$  (10.0  $\text{cm}^3$ ) was stirred at 100  $^{\circ}\text{C}$  for 30 min, the mixture was cooled to 0  $^{\circ}\text{C}$ , and then pH was adjusted to 12–14 with 2 N NaOH. The whole mixture was extracted with  $CHCl_3$ , and dried over  $MgSO_4$ . After concentration, the resulting solid was recrystallized from *n*-hexane- $CHCl_3$  to give the title compound 9 as colorless crystals (137 mg, 52%): mp 83–85  $^{\circ}\text{C}$ ; IR (neat)  $\nu_{\text{max}}/\text{cm}^{-1}$ : 2931 (NH<sub>2</sub>), 2851 (NH<sub>2</sub>), 1620 (C=N);  $\delta_H$  (500 MHz,  $CDCl_3$ ): 1.81–1.86 (2H, m,  $CH_2$ ), 3.48 (4H, t,  $J$  5.7,  $CH_2$ ), 4.71 (1H, br s, NH), 5.72 (2H, br s, NH<sub>2</sub>), 6.61–6.66 (2H, m, Ar), 7.07–7.10 (1H, m, Ar), 7.23–7.26 (1H, m, Ar);  $\delta_C$  (125 MHz,  $CDCl_3$ ): 20.7, 42.0 (2C), 116.56, 116.61, 119.3, 126.3, 129.9, 147.1, 155.1; anal. calcd for  $C_{10}H_{14}N_3$ : C, 68.54; H, 7.48; N, 23.98. Found: C, 68.28; H, 7.57; N, 23.75.

**3,4-Dihydro-2*H*,6*H*-benzo[e]pyrimido[2,1-*b*][1,3]thiazin-6-one** (10). PD 404182 (1a) (20.0 mg, 0.0920 mmol) in DMSO (0.920  $\text{cm}^3$ ) was dissolved in  $H_2O$  (92.0  $\text{cm}^3$ ). After being stirred at 37  $^{\circ}\text{C}$  for 26 h, the mixture was purified by HPLC. The resulting solid was dissolved in EtOAc, and washed with sat.  $NaHCO_3$ . The extract was dried over  $MgSO_4$ , and concentrated to give the title compound 10 as colorless crystals (5.2 mg, 27%); mp 132–134  $^{\circ}\text{C}$  (from toluene); IR (neat)  $\nu_{\text{max}}/\text{cm}^{-1}$ : 1662 (C=O), 1608 (C=N);  $\delta_H$  (500 MHz,  $CDCl_3$ ): 1.98–2.03 (2H, m,  $CH_2$ ), 3.57 (2H, t,  $J$  5.7,  $CH_2$ ), 4.02 (2H, t,  $J$  6.0,  $CH_2$ ), 7.17 (1H, d,  $J$  8.0, Ar), 7.26–7.29 (1H, m, Ar), 7.47 (1H, ddd,  $J_1 = J_2$  7.7,  $J_3$  1.6, Ar), 8.24 (1H, dd,  $J_1$  8.0,  $J_2$  1.1, Ar);  $\delta_C$  (125 MHz,  $CDCl_3$ ): 21.1, 42.3, 46.1, 122.7, 124.0, 126.0, 131.0, 133.2, 134.3, 147.0, 161.7; HRMS (FAB):  $m/z$  calcd for  $C_{11}H_{11}N_2OS$  [M + H]<sup>+</sup> 219.0592; found: 219.0584. Spectral data were in good agreement with those previously reported.<sup>26</sup>

*Crystal structure:*  $C_{11}H_{10}N_2OS$ ,  $M_w$ : 218.27, primitive monoclinic,  $a = 10.7456(6)$ ,  $b = 10.9466(6)$ ,  $c = 16.6473(7)$   $\text{\AA}$ ,  $\beta = 97.422(2)^{\circ}$ ,  $V = 1941.78(17)$   $\text{\AA}^3$ , space group  $P2_1/n$  (no. 14),  $Z = 8$ . The data were collected with a Rigaku R-AXIS RAPID diffractometer using graphite monochromated Mo-K $\alpha$  radiation at –93 K. 18 534 Reflections were measured, 4420 unique

( $R_{\text{int}} = 0.0220$ ) which were used in all calculations. The final  $wR$  was 0.0796. The substance was crystallized from toluene. The CCDC deposition number: 1037500.

**3,4-Dihydro-2*H*-benzo[4,5]thiazolo[3,2-*a*]pyrimidine** (11). Following the reported procedure,<sup>27</sup> 2-chloro-1,3-benzothiazole (0.619  $\text{cm}^3$ , 5.00 mmol) was converted to the title compound 11, as a pale yellow solid (407 mg, 2 steps 73%), by the reaction with 3-amino-1-propanol (0.381  $\text{cm}^3$ , 5.00 mmol). The product (80.8 mg) was purified by HPLC to give the compound 11 as a colorless solid (72.3 mg, TFA salt): IR (neat)  $\nu_{\text{max}}/\text{cm}^{-1}$ : 3274–3181 (OH), 1672 (C=O), 1632 (C=N);  $\delta_H$  (500 MHz,  $CD_3OD$ ): 2.31–2.36 (2H, m,  $CH_2$ ), 3.70 (2H, t,  $J$  5.7,  $CH_2$ ), 4.27 (2H, t,  $J$  6.0,  $CH_2$ ), 7.42–7.45 (1H, m, Ar), 7.57–7.61 (2H, m, Ar), 7.84 (1H, d,  $J$  8.0, Ar);  $\delta_C$  (125 MHz,  $CD_3OD$ ): 19.2, 41.3, 44.0, 113.5, 118.0 (q,  $J$  291.5), 123.1, 124.1, 126.7, 129.2, 140.0, 162.4 (q,  $J$  36.0), 166.0; HRMS (FAB):  $m/z$  calcd for  $C_{10}H_{11}N_2S$  [M + H]<sup>+</sup> 191.0643; found: 191.0641.

**3,4-Dihydro-2*H*-benzo[4,5]isothiazolo[2,3-*a*]pyrimidine 6,6-dioxide** (12). 50%  $H_2O_2$  (0.0391  $\text{cm}^3$ , 0.638 mmol) was added to a suspension of compound 2a (48.5 mg, 0.159 mmol, TFA salt) in TFA (0.610  $\text{cm}^3$ ) dropwise. After being stirred at room temperature for 24 h, the mixture was quenched with  $Et_3N$  (1.50  $\text{cm}^3$ ), and the whole mixture was extracted with EtOAc. The extract was washed with sat.  $NaHCO_3$ , brine, and dried over  $MgSO_4$ . After concentration, the residue was purified by flash chromatography over aluminum oxide with *n*-hexane-EtOAc (3:2 to 1:1) to give the title compound 12 as colorless crystals (25.2 mg, 71%); mp 146–148  $^{\circ}\text{C}$  (from  $Et_2O$ -MeOH); IR (neat)  $\nu_{\text{max}}/\text{cm}^{-1}$ : 1670 (C=N);  $\delta_H$  (500 MHz,  $CDCl_3$ ): 2.01–2.05 (2H, m,  $CH_2$ ), 3.70 (2H, t,  $J$  5.7,  $CH_2$ ), 3.79 (2H, t,  $J$  6.0,  $CH_2$ ), 7.70–7.76 (2H, m, Ar), 7.85–7.88 (1H, m, Ar), 7.98–8.01 (1H, m, Ar);  $\delta_C$  (125 MHz,  $CDCl_3$ ): 19.5, 36.5, 44.2, 120.7, 122.8, 129.6, 132.4, 133.6, 134.8, 142.7; anal. calcd for  $C_{10}H_{10}N_2O_2S$ : C, 54.04; H, 4.54; N, 12.60. Found: C, 53.89; H, 4.74; N, 12.36.

**2-(2-Hydroxyphenyl)-1,4,5,6-tetrahydropyrimidine** (16).<sup>21</sup> A mixture of methyl salicylate (1.29  $\text{cm}^3$ , 10.0 mmol) and propylenediamine (2.56  $\text{cm}^3$ , 30.0 mmol) was refluxed for 16 h. The crude product was dissolved in MeOH, and crystallized with  $Et_2O$ . The precipitate was filtered, and the unreacted propylenediamine was removed by washing with  $Et_2O$ . The resulting solid was purified by flash chromatography over aluminum oxide with  $CHCl_3$ -MeOH (10:0 to 10:1) to give the title compound 16 as colorless crystals (497 mg, 28%); mp 259–261  $^{\circ}\text{C}$  (from  $Et_2O$ -MeOH); IR (neat)  $\nu_{\text{max}}/\text{cm}^{-1}$ : 3212–3050 (OH), 1620 (C=N);  $\delta_H$  (500 MHz,  $DMSO-d_6$ ): 1.83–1.88 (2H, m,  $CH_2$ ), 3.40 (4H, t,  $J$  5.7,  $CH_2$ ), 6.27–6.30 (1H, m, Ar), 6.46 (1H, d,  $J$  8.6, Ar), 7.04–7.08 (1H, m, Ar), 7.45 (1H, dd,  $J_1$  8.0,  $J_2$  1.7, Ar), 12.10 (1H, br s, OH);  $\delta_C$  (125 MHz,  $CD_3OD$ ): 20.0, 39.3 (2C), 111.2, 114.5, 124.4, 126.3, 135.1, 160.9, 172.5; anal. calcd for  $C_{10}H_{12}N_2O$ : C, 68.16; H, 6.86; N, 15.90. Found: C, 68.13; H, 7.03; N, 16.09.

### Determination of anti-HIV activity

The anti-HIV activity of a series of compounds against HIV-1<sub>IIIB</sub> was determined by the NCK assay.<sup>29</sup> The target



cells (NCK45- $\beta$ -Gal;  $10^4$  cells per well) were plated in 96-well flat microtiter culture plates. On the following day, the cells were inoculated with HIV-1 (60 NCK U per well, giving 60 blue cells after 48 h of incubation) and cultured in the presence of various concentrations of the test compounds in fresh medium. Forty-eight hours after viral exposure, all the blue cells stained with X-Gal (5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactopyranoside) were counted in each well. Cytotoxic effects were not observed at 10  $\mu$ M except for compound **2g**.

The activity of test compounds was determined as the concentration that blocked HIV-1 infection by 50% (50% effective concentration [ $EC_{50}$ ]).  $EC_{50}$  was determined by using the following formula:

$$EC_{50} = 10^{[\log(A/B) \times (50-C)/(D-C) + \log(B)]},$$

where  $A$ , of the two points on the graph that bracket 50% inhibition, the higher concentration of the test compound;  $B$ , of the two points on the graph that bracket 50% inhibition, the lower concentration of the test compound;  $C$ , inhibitory activity (%) at the concentration  $B$ ;  $D$ , inhibitory activity (%) at the concentration  $A$ .

### Glutathione-mediated transformation of pyrimidobenzothiazine and isothiazolopyrimidine derivatives

The compound (1 mM) was incubated in 10 mM GSH (in a mixture of 50 mM phosphate buffer (pH 7.4) and MeCN [75 : 25 (v/v) or 1 : 1 (v/v)]) at 37 °C. The sample was analyzed by HPLC and the peak area was recorded by UV detection at 254 nm. The concentrations of the products were calculated using previously determined calibration curves. Of note, the GSH-mediated transformation from **1a** was slightly accelerated by the addition of MeCN in phosphate buffer.

### Acknowledgements

This work was supported by Grants-in-Aid for Scientific Research from JSPS, Japan; Platform for Drug Discovery, Informatics, and Structural Life Science from MEXT, Japan; a Grant-in-Aid for Research on HIV/AIDS from the Ministry of Health and Welfare of Japan. T.M. is grateful for JSPS Research Fellowships for Young Scientists.

### References

- PD 404182 (**1a**) was previously reported to be an enzyme inhibitor against 3-deoxy-D-manno-octulosonic acid 8-phosphate synthase<sup>2</sup> and phosphopantetheinyl transferase.<sup>3</sup> In addition, it was recently reported that PD 404182 (**1a**) inhibits human dimethylarginine dimethylaminohydrolase isoform 1 (DDAH).<sup>4</sup>
- M. R. Birck, T. P. Holler and R. W. Woodard, *J. Am. Chem. Soc.*, 2000, **122**, 9334–9335.
- C. R. Vickery, N. M. Kosa, E. P. Casavant, S. Duan, J. P. Noel and M. D. Burkart, *ACS Chem. Biol.*, 2014, **9**, 1939–1944.
- Y. T. Ghebremariam, D. A. Erlanson and J. P. Cooke, *J. Pharmacol. Exp. Ther.*, 2014, **348**, 69–76.
- K. Chockalingam, R. L. Simeon, C. M. Rice and Z. Chen, *Proc. Natl. Acad. Sci. U. S. A.*, 2010, **107**, 3764–3769.
- A. M. Chamoun, K. Chockalingam, M. Bobardt, R. Simeon, J. Chang, P. Gallay and Z. Chen, *Antimicrob. Agents Chemother.*, 2012, **56**, 672–681.
- T. Mizuhara, S. Oishi, H. Ohno, K. Shimura, M. Matsuoka and N. Fujii, *Org. Biomol. Chem.*, 2012, **10**, 6792–6802.
- T. Mizuhara, S. Oishi, H. Ohno, K. Shimura, M. Matsuoka and N. Fujii, *Bioorg. Med. Chem.*, 2012, **20**, 6434–6441.
- T. Mizuhara, S. Oishi, H. Ohno, K. Shimura, M. Matsuoka and N. Fujii, *Bioorg. Med. Chem.*, 2013, **21**, 2079–2087.
- A. M. Chamoun-Emanuelli, M. Bobardt, B. Moncla, M. K. Mankowski, R. G. Ptak, P. Gallay and Z. Chen, *Antimicrob. Agents Chemother.*, 2014, **58**, 687–697.
- S. Okazaki, T. Mizuhara, K. Shimura, H. Murayama, H. Ohno, S. Oishi, M. Matsuoka and N. Fujii, *Bioorg. Med. Chem.*, 2015, **23**, 1447–1452.
- M. Baba, D. Schols, R. Pauwels, H. Nakashima and E. De Clercq, *J. Acquired Immune Defic. Syndr.*, 1990, **3**, 493–499.
- T. Matthews, M. Salgo, M. Greenberg, J. Chung, R. DeMasi and D. Bolognesi, *Nat. Rev. Drug Discovery*, 2004, **3**, 215–225.
- Of note, it was reported that PD 404182 showed the anti-viral activities by the virucidal effect *via* the physical disruption of virions, see ref. 6 and 10.
- T. Mizuhara, S. Oishi, N. Fujii and H. Ohno, *J. Org. Chem.*, 2010, **75**, 265–268.
- M. Ishihara and H. Togo, *Tetrahedron*, 2007, **63**, 1474–1480.
- For a review on the function of glutathione, see: R. Franco, O. J. Schoneveld, A. Pappa and M. I. Panayiotidis, *Arch. Physiol. Biochem.*, 2007, **113**, 234–258 and references cited therein.
- Compound **4a** was also identified from HTS as an anti-bacterial agent that inhibits localization of lipoproteins (Lol) during cell wall biosynthesis, see: H. Ito, A. Ura, Y. Oyamada, H. Yoshida, J. Yamagishi, S. Narita, S. Matsuyama and H. Tokuda, *Microbiol. Immunol.*, 2007, **51**, 263–270.
- During the course of our investigations, an initial report describing GSH-mediated degradation of PD 404182 appeared, see ref. 4.
- In the absence of GSH, compound **1a** was converted into pyrimidothiazinone **10** as discussed above.
- R. Mitsuhashi, T. Suzuki and Y. Sunatsuki, *Inorg. Chem.*, 2013, **52**, 10183–10190.
- J. A. Loo, T. P. Holler, J. Sanchez, R. Gagliotti, L. Maloney and M. D. Reily, *J. Med. Chem.*, 1996, **39**, 4313–4320.
- T. Vercruyse, B. Basta, W. Dehaen, N. Humbert, J. Balzarini, F. Debaene, S. Sanglier-Cianfrani,



C. Pannecouque, Y. Mély and D. Daelemans, *Retrovirology*, 2012, **9**, 95–108.

24 C. Pannecouque, B. Szafarowicz, N. Volkova, V. Bakulev, W. Dehaen, Y. Mély and D. Daelemans, *Antimicrob. Agents Chemother.*, 2010, **54**, 1461–1468.

25 The transformations from **1f** and **2f** into the ring-opened thiophenol derivative were also observed in the presence of GSH.

26 D. Chen, J. Wu, J. Yang, L. Huang, Y. Xiang and W. Bao, *Tetrahedron Lett.*, 2012, **53**, 7104–7107.

27 P. A. Woods, L. C. Morrill, T. Lebl, A. M. Z. Slawin, R. A. Bragg and A. D. Smith, *Org. Lett.*, 2010, **12**, 2660–2663.

28 S. Sivaramakrishnan, A. H. Cummings and K. S. Gates, *Bioorg. Med. Chem. Lett.*, 2010, **20**, 444–447.

29 K. Kajiwara, E. Kodama, Y. Sakagami, T. Naito and M. Matsuoka, *J. Clin. Microbiol.*, 2008, **46**, 792–795.

