RSC Advances



PAPER

View Article Online



Cite this: RSC Adv., 2015, 5, 18894

Building polyfunctional piperidines: a stereoselective strategy of a three-component Mannich reaction inspired by biosynthesis and applications in the synthesis of natural alkaloids (+)-241D; (-)-241D; isosolenopsin A and (-)-epimyrtine†

Yang Yang*

A general method to assemble multi-substituted chiral piperidines was developed, inspired by the biosynthesis of piperidine natural products. In biosynthesis, Δ^1 -piperideine 4 plays a key role as a common intermediate giving rise to a variety of piperidine-based natural alkaloids. Nature uses L-lysine as a building block, enzymatically transforming it into a δ -amino carbonyl intermediate 3 as the precursor to cyclize into Δ^1 -piperideine 4. We envisioned that such a process could be accomplished by a vinylogous type Mannich reaction if a functionalized dienolate was employed. A stereoselective three-component vinylogous Mannich-type reaction (VMR) of 1,3-bis-trimethylsily enol ether 7 was therefore investigated and was found to give cyclized chiral dihydropyridinone compound 9 as an adduct. Like Δ^1 -piperideine in biosynthesis, the chiral 2,3-dihydropyridinone compound 9 from VMR is a versatile intermediate for building a variety of new chiral piperidine compounds. The method was showcased by concise two-step approaches in the synthesis of the bioactive natural alkaloids (+)-241D; (-)-241D and isosolenopsin A. Furthermore, when properly functionalized substrate aldehyde 24 was employed, the corresponding dihydropyridinone adduct 25 cyclized to form a second piperidine ring, leading to a chiral polyfunctional quinolizidine enaminone 27. This versatile intermediate was used to prepare a variety of new chiral quinolizidine compounds, including natural alkaloid (-)-epimyrtine.

Received 12th November 2014 Accepted 6th February 2015

DOI: 10.1039/c4ra14418j

www.rsc.org/advances

Introduction

Functionalized piperidine rings are common moieties incorporated in a variety of natural alkaloids and pharmaceutical molecules. In fact, piperidine is the most frequently used non-aromatic ring in small molecule drugs listed in the FDA orange book. Developing synthetic approaches for the stereoselective construction of these ring systems has been an area of intense research in synthetic organic chemistry for decades. Among the various piperidine derivatives, 2 and/or 6 substituted piperidines are particularly common and interesting since such substitution patterns block the metabolism of the piperidine ring and potentially have a significant impact on the ring's 3D conformation. For such reasons, installation of α substitutions adjacent to the piperidine nitrogen are commonly employed as a strategy in medicinal chemistry research to tune either

substituents serving this purpose. Interestingly, α-methyl multisubstituted piperidines are also commonly found in naturally occurring piperidine alkaloids such as (–)-pinidinol, (+)-241D and isosolenopsin A *etc.* (Fig. 1). Some of these natural alkaloids have demonstrated interesting pharmacological properties and served as valuable starting points for new drug discovery.⁵ The biosynthetic pathway of many piperidine-based natural

biological activities or pharmacological properties. In practice,

the methyl group is one of the most common and simplest

The biosynthetic pathway of many piperidine-based natural alkaloids has been studied. Δ^1 -Piperideine 4, which forms from an intramolecular imine cyclization of a δ-amino pentanal precursor 3, was believed to be a key common intermediate in the pathway. Studies have shown that further transformations on this prototype piperidine ring lead to a variety of structurally diversified piperidine, quinolizidine and indolizidine alkaloids in nature. The basic starting building block in this pathway is L-lysine, which undergoes several enzymatically catalyzed transformations, including decarboxylation by LDC (lysine decarboxylase) and oxidative deamination by CuAO (copper amine oxidase). The resulting δ-amino pentanal 3 then gives rise to the key Δ^1 -piperideine ring (Fig. 2). However, without nature's

Genomics Institute of the Novartis Research Foundation, 10675 John Jay Hopkins Drive, San Diego, California 92121, USA. E-mail: yyang@gnf.org

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c4ra14418i

Paper RSC Advances

Fig. 1 Examples of natural alkaloids incorporating α -methy substituted piperidines.

powerful enzyme tools, chemical synthesis of Δ^1 -piperideine is tedious7 due to its instability and such intermediate is therefore not practical to be widely applied in synthesis lab like its role in biosynthesis.8 We envisioned however that similar δ-amino carbonyl precursor for Δ^1 -piperideine can be assembled conveniently via a vinylogous Mannich-type reaction (VMR) with an aldimine if a properly functionalized dienolate was employed. As shown in Fig. 2, cyclization of the initial δ -amino carbonyl adduct would lead to a 2,3-dihydropyridinone, which could also be viewed as a tautomeric form of cyclic imine, but more stable and easier to handle (Fig. 2). In fact, the synthetic utility of dihydropyridinones has been extensively investigated by the Comins group, but to date the methodology for preparation of these intermediates has been limited.9 Here we report the successful implementation of the VMR strategy to generate useful chiral dihydropyridone intermediates, and their subsequent transformation to a variety of interesting piperidinecontaining natural products and compounds of medicinal interest.

Results and discussion

The simple 1,3-bis-trimethylsily enol ether 7 has been employed as a vinylogous nucleophilic reagent in several organic

transformations such as cyclization with 1,2-dielectrophiles, bromination, and vinylogous aldol reaction. ¹⁰ Surprisingly, the use of 7 as dienolate in a Mannich-type reaction has never been reported. ¹¹ To ensure stereoselective control in VMR, inexpensive commercially available chiral α -methyl benzylamine 6 was employed to form chiral aldimines *in situ*. The three-component VMR reaction of 6 and 7 with various aldehydes 5 was carried out in the presence of Sn(OTf)₂ in DCM at -78 °C to 0 °C. Corresponding adducts 8 were observed from reaction LC-MS analysis, however in a mixture with cyclized 2,3-dihydropyridinone products 9. Treatment of the crude mixture with a catalytic amount of acetic acid in DCM led to complete conversion of acylic adducts 8 into 9 (Scheme 1).

The results of the VMR reaction of 7 with various aldehydes are summarized in Table 1. Most of the reactions showed moderate to good yields. A variety of functional groups were well tolerated. The reactions showed excellent diastereoselectivities since in all cases only single isomers were observed and isolated from the reaction mixtures. In order to confirm that the stereoselectivities of the reaction were auxiliary directed, compounds 9d-I and 9d-II were prepared from the same chiral substrate aldehyde 5d, in the presence of chiral amine auxiliary 6a and its enantiomer 6b. The proton NMR spectra of these compounds showed that the $J_{\text{Ha/Hb}}$ value for **9d-I** was 8.80 Hz while the corresponding value for 9d-II was 9.2 Hz, suggesting that 9d-I and 9d-II were the erythro and threo isomers respectively, based on literature precedent.12 These results confirmed auxiliary directed stereoselectivities and further supported the established sense of stereochemical induction in such VMR13 (Fig. 3).

To examine the synthetic utility of 2,3-dihydropyridinones obtained from the VMR, adduct compound **9h** was selected to probe further transformations. When the compound **9h** was treated with TFA at room temperature, the chiral benzyl directing group was cleaved to give cyclic enaminone **10** in quantitative yield (Scheme 2). We also found that the corresponding chiral substituted piperidine could be obtained from **9h** *via* palladium catalyzed hydrogenation. Interestingly, under different hydrogenation conditions, the reduction of **9h** yielded




Fig. 2 VMR strategy for piperidine synthesis inspired by biosynthesis.

RSC Advances Paper

Scheme 1

different major piperidine products. When hydrogenation was performed in MeOH in presence of palladium on carbon at room temperature, the reaction cleaved the chiral benzyl group and saturated the 2,3-dihydro-4-pyridinone simultaneously to give cis-3-hydroxy 2,6-disubstituted piperidine compound 11 stereospecifically as the major product, accompanied by deoxygenated piperidine compound 12 as the minor product (11/12, ratio 10:1)14 (Scheme 2). However, when the hydrogenation was performed in a Parr hydrogenator under 40 psi hydrogen pressure in a mixture of methanol and acetic acid (1/1), the major product was deoxygenated cis-2,6-dialkylated piperidine 12 accompanied by 11 as the minor product (12/11, 5:1)(Scheme 2). The results could be explained by a shift in the equilibrium between 2,3-dihydropridinone and 2,3-dihydropridinium under different conditions. 15 Presumably, 2,3dihydropridinone is the major species present under neutral conditions and hydrogenation led to 4-hydroxy piperidine product 11. However, under acidic conditions, the protonated 2,3-dihydropridinium species is the major (or more reactive) species present, and hydrogenation gives the corresponding deoxygenated piperidine 12 as the major product¹⁶ (Fig. 4). Similar hydrogenations of 2,3-dihydropyridinones have been reported to yield 4-hydroxy piperidine compounds stereospecifically.14 However, to our knowledge the direct deoxygenative reduction of 2,3-dihydropridinones is rarely reported. The results allowed accessing different substitution type piperidine compounds from 2,3-dihydropyridinones 9 by simply switching different reduction conditions.

We further probed the utility of our chiral piperidine intermediates by applying the VMR methodology to the asymmetric synthesis of natural piperidine-containing alkaloids. Dendrobate alkaloid (+)-241D and its enantiomer (–)-241D were among our first targets. Dendrobate alkaloid (+)-241D was isolated from the methanolic skin extracts of the Panamanian poison frog *Dendrobates speciosus*. The alkaloid shows

interesting bioactivity as a potent non-competitive blocker of acetylcholine and ganglionic nicotinic receptor channels.18 The structure of (+)-241D features an all-cis 2,4,6-trisubstituted piperidine core bearing three chiral centers. The asymmetric synthesis of (+)-241D has been reported by multiple research groups via a variety of synthesis routes employing between eight and eighteen steps.19 We were delighted to find that using the newly developed VMR strategy, the asymmetric synthesis of (+)-241D and its enantiomer could be accomplished simply in two steps from inexpensive commercial materials. Using chiral α-methyl benzylamines 6a & 6b to control stereochemistry, the reaction of bis-trimethylsily enol ether 7 with decanal 13 yielded chiral adducts 14 & 15 respectively. Subsequent reduction of 2,3dihydro-4-pyridones 14 & 15 by palladium-catalyzed hydrogenation in methanol gave (+)-241D and (-)-241D in good yield (Scheme 3).

The versatile utility of such VMR approach in assembling piperidine was further exampled in asymmetric synthesis of another natural alkaloid isosolenopsin A which incorporate cis-2,6-dialkylpiperidine as a core. isosolenopsin A was isolated from the venom of the fire ant solenopsis and was found to have a variety of interesting bioactivities including antibiotic, antifungal, anti-HIV, blockade of neuromuscular transmission and potent and selective inhibition of the neuronal nitric oxide synthase.20 By the similar strategy, corresponding VMR adduct 2,3-dihydro-4-pyridones 17 was obtained when dodecanal 16 and chiral amine 6b were employed. The palladium-catalyzed reduction on 2,3-dihydro-4-pyridone 17 was carried out in methanol in presence of acetic acid (50%) under 40 psi hydrogen pressure in a Parr hydrogenator. Corresponding deoxygenated product isosolenopsin A was obtained as the major product in moderate yield (45%) (Scheme 4). The current approach presented the shortest route for asymmetric synthesis of isosolenopsin A than any other reported methods.21

Table 1 Asymmetric three-component vinylogous Mannich reactions of 1,3-bis-trimethylsily enol ether 7

Entry	Substrate 5	6	9 (yield)
1	CHO 5a	6 b	OMe 9a (71%)
2	CHO 5b	6b	OMe 9b (65%)
3	∕∕✓∕CHO 5c	6b	$^{\circ}$ OMe $^{\circ}$ $^{\circ$
4	СHO ОМе 5d	6b	Meo H Meo
5	CHO OMe 5d	6 a	MeO H Me OMe 9d-II (64%)
6	OBn 5e	6a	OBn NMe OMe 9e (66%)
7	OBn 5f	ба	OBn Me OMe 9f (75%)
8	Sg CHO	6a	OMe 9g (75%)
9	CHO 5h	6b	OMe 9h (53%)

MeO, N Me MeO, N Me Ph N Me Ph

Fig. 3 NMR analysis of 9d-I and 9d-II.

Beyond applying to building simple multi-substituted piperidine compounds, current VMR strategy also provides potentials in synthesizing chiral quinolizidine compounds. Quinolizidine compounds structurally incorporate two fused piperidine rings sharing common nitrogen. Like piperidine, quinolizidine represent both a class compound of pharmaceutical interest and an important family of natural alkaloids. In nature, several hundred structurally related quinolizidine compounds have been identified from a variety of natural sources, predominately from plants and amphibian skin.22 Some natural quinolizidine alkaloids exhibit interesting pharmacological properties23 and serve as important starting points for the drugs discovery.24 Interestingly, the biosynthesis of some quinolizidine alkaloids shares the same pathway of the natural piperidine alkaloids that undergo the same Δ^1 -piperideine intermediate 4, enzymatically starting from L-lysine.25 As an example, in the biosynthesis of quinolizidine alkaloids lupinine, Δ^1 -piperideine is also the key intermediate to assemble the first piperidine ring for the quinolizidine core. To build the second piperidine ring, two Δ^1 -piperideine intermediates undergo a cross aldol-type coupling and one of the imine systems gets hydrolyzed after coupling and undergoes oxidation resulting in primary amine function 20. Ultimately the formation of the quinolizidine nucleus in biosynthesis is accomplished by another intramolecular imine formation (Fig. 5). We however envisioned that in the VMR we developed, if the aldimine substrate has been properly functionalized, piperidinelike adducts arising from the asymmetric VMR may further conveniently cyclize to form second piperidine ring to give the desired quinolizidine product. As shown in Fig. 5, if a δ-leaving group is incorporated in the aldimine substrate and can be tolerated in the asymmetric VMR for the first piperidine ring construction, the quinolizidine structure 22 should be readily formed by a subsequent intramolecular SN2 cyclization (Fig. 5).

To test this idea 5-chloropentanal 24 was prepared from the oxidation of 5-chloropentan-1-ol 23 and the corresponding three-component VMR reaction was carried out. The reaction gave adduct 25 in expected excellent diastereoselectivity as a single stereoisomer. The δ chloride group which serve as a future leaving group on the substrate, was well tolerated (Scheme 5). With dihydropyridinone 25 in hand we set out to construct the second ring for a quinolizidine core. The α-methyl benzyl group was cleaved cleanly upon the treatment with TFA at room temperature overnight to give compound 26 in quantitative yield. In presence of sodium hydride in DMF, intramolecular SN2 cyclization by 26 led to a quinolizidine intermediate 27 as a cyclic enaminone (Scheme 5). We envisaged that such cyclic enaminone 27 could be a valuable polyfunctional quinolizidine intermediate since different organic transformations can be carried out at different positions on this molecule. It provides convenient entries to access different types chiral quinolizidine compounds (Fig. 6).

The reduction of cyclic enanminone 27 was first explored. It was found that under the conditions of either palladium-catalyzed hydrogenation in methanol or treating L-selectride (LiBu $^{i}_{3}$ BH) in THF, both alkene and carbonyl were reduced affording *cis*-2-hydroxyl-4-methly quinolizidine 28 as product (Scheme 6, eqn (1)). Similarly as in the reduction of 2,3-dihydropyridinone, the reduction on quinolizidine enanminone also proceeded in stereoselective manner which is in agree with literature precedents. He had the reduction was carried out with "super hydride" (LiEt₃BH) in presence of BF₃·Et₂O in THF, the alkene functionality was selectively reduced, giving 29 as natural quinolizidine alkaloid (—)-epimyrtine as the product in good yield²⁷ (Scheme 6, eqn (2)). The results provided a concise approach for the enantioselective synthesis of such natural alkaloid.

Conjugate additions to quinolizidine enaminone 27 were also explored. Although direct conjugate addition of Grignard reagents to cyclic enaminones has been previously reported,²⁹ in our hands, treatment of 27 with methyl magnesium bromide

Paper RSC Advances

Fig. 4 Equilibrium between 2,3-dihydropridinone and 2,3-dihydropridinium.

did not yield expected product (Table 1, entry 1). Similarly, when methyl cuprate was employed, only a trace amount of product 30a was observed (Table 1, entry 2). However, when methyl Grignard addition was carried out in the presence of TMS-Cl, 1,4-conjugate addition went smoothly giving adduct 30 in good yield³⁰ (Table 2, entry 3). Under similar conditions, conjugate additions by vinyl and allyl Grignard reagents were also performed (Table 2, entries 4 & 5). As the similar examples reported in literature, such conjugate addition on quinolizidine enaminone proceeded in stereoselective manner by generating a quaternary chiral carbon in the product (Scheme 7).³¹

Finally, to further probe structural diversification, alkylation reaction on the methyl side chain of 27 was investigated. It was found that a corresponding enolate can be generated by treating 27 with LiN(SiMe₃)₂ (LiHMDS) in THF at low temperature. By subsequently treating such enolate with alkylating agents 31, corresponding alkylation products 32 could be obtained smoothly. The results of such reaction were summarized in Table 3. The reactions gave moderate to good yields by showing the tolerance toward different functional groups. No

epimerization was observed in such enolate alkylation. Such alkylation reaction led to the side chain extension and provided opportunities to synthesize more structural diversified quinolizatione-based compound beyond α methyl substituted type (Scheme 8).

Conclusion

In summary, inspired by the biosynthesis pathway of natural piperidine-based alkaloids, a general and practical approach to synthesize multi-substituted chiral piperidine was developed via a stereoselective three-component vinylogous Mannich-type reaction (VMR) by using 1,3-bis-trimethylsily enol ether 7 as a dienolate. The corresponding VMR adduct was chiral 2,3dihydropyridinones 9 which played the role of cornerstone in building new targeted chiral piperidine compounds. The efficiency of such stereoselective synthesis approach was exampled in developing novel synthesis of bioactive natural alkaloids: dendrobate alkaloids (+)-241D; (-)-241D, and isosolenopsin A in highly concise manners. Beyond simple piperidine compound synthesis, the method also provided rapid route for chiral quinolizidine construction. When prefunctionalized substrate aldehyde 24 was employed, the corresponding VMR adduct could cyclize to give versatile quinocyclic enaminone 27. The different types transformations carried out on such polyfunctional intermediate gave rise a variety of new chiral quinolizidine compounds, including natural alkaloid (-)-epimyrtine. We believe the presented VMR approach offers a general; stereoselective and efficient way to assemble multi-substituted chiral piperidine-based compound in organic synthesis.

Experimental section

General methods

All commercial reagents and solvents were used without purification. ¹H and ¹³C NMR spectra were recorded on a Bruker 400 MHz spectrometer using TMS as the internal standard (0 ppm). TLC analyses were carried out on aluminum sheets precoated with silica gel 60 F254, and UV radiation was used for detection.

Scheme 3 Enantioselective synthesis of (+)-241D and (-)-241D.

Scheme 4 Enantioselective synthesis of natural alkaloid isosolenopsin A.

Flash column chromatography was performed on silica gel (SiliaFlash F60, 230-400 mesh). LC/MS analysis was performed on an Agilent 1100 series system equipped with an Agilent 1100 series binary pump, Agilent 1100 series autosampler, Agilent 1100 series DAD UV detector, Agilent 1100 series single quadrupole mass spectrometer with ESI source, and a SEDEX 75 ELSD. The mass spectrometer was set to scan from 100 to 1000 AMU. Mass spectrometric data were acquired in the positive ionization mode. The mobile-phase solvents used were (A) 0.05% aq. TFA; and (B) 0.035% TFA in MeCN. The total mobile phase flow rate was 1.0 mL min⁻¹. The gradient was 10-90% in 3 min with an isocratic hold of 100% mobile-phase B for 0.49 min at the end of the gradient. A Waters Atlantis T3 (5 μ m, 2.1 \times 50 mm) column was used. Chemical shifts of NMR spectra are reported in ppm, multiplicities are indicated by s (singlet), d (doublet), t (triplet), q (quartet), and m (multiplet). All vinylogous Mannich reactions were carried out in oven-dried glassware under air atmosphere. 1,3-Bis-trimethylsily enol ether 7 was prepared freshly following the procedure from literature. 10d Diastereo-selectivities of VMR described in the manuscript were determined both by HPLC and NMR.

(*S*)-2-Cyclohexyl-1-((*S*)-1-(4-methoxyphenyl)ethyl)-6-methyl-2,3-dihydropyridin-4(1*H*)-one (9a). To a round-bottom flask contains (*R*)-1-(4-methoxyphenyl)ethanamine (6b) (151 mg, 1 mmol, 1 eq.) in dried DCM (0.1 M) solution added 4-AMS (500 mg mmol⁻¹) followed by cyclohexane-carbaldehyde (5a) (112 mg, 1 mmol, 1 eq.). After stirring at room temperature for

10 min, 1,3-bis-trimethylsily enol ether 3 (292 mg, 1.2 mmol, 1.5 eq.) was added and the mixture solution was cooled to -78 °C. Tin(II) triflate (412 mg, 1 mmol, 1 eq.) was then added and the reaction was stirred at this temperature for 8 h. The reaction temperature was raised to 0 °C and kept the same temperature overnight. LC-MS analysis showed the mixture of 8a and 9a as new products ($\sim 2/1$). The reaction was guenched with saturated aqueous solution of sodium bicarbonate and removed solid via filtration. The reaction mixture was then extracted with DCM (15 mL \times 5). The combined organic phase was treated with acetic acid (0.1 mL) and the resulting solution was stirred at room temperature for 1 h until 8a disappeared from LC-MS analysis. The solution was then basified by treating with saturated aqueous solution of sodium bicarbonate and washed with brine and dried over sodium sulfate. After concentration, the crude product was purified by ISCO silica gel chromatography (ethyl acetate/hexane) to afford the product as colorless oil (232 mg, 71% yield). ¹H-NMR (400 MHz, CDCl₃, 25 °C): $\delta = 0.87$ (m, 1H); 1.30 (m, 4H); 1.41 (m, 1H); 1.55 (d, 3H, I = 6.8 Hz, CH₃-); 1.70 (m, 5H); 2.02 (m, 2H); 2.07 (s, 3H, CH₃-); 2.97 (m, 1H); 3.70 (s, 3H, OCH₃); 4.87 (s, 1H, CH=); 4.94 (q, 1H, J = 6.8 Hz); 6.80 (m, 2H); 7.13 (d, 2H, J = 8.4Hz). ¹³C-NMR (100 MHz, CDCl₃, 25 °C): $\delta = 18.8$; 21.8; 26.2; 26.5; 26.6; 29.7; 30.6; 36.6; 40.9; 55.3; 558; 57.3; 102.5; 114.1; 127.5; 133.8; 159.1; 161.6; 191.9. LC-MS: 100% (purity), m/z: 328 (M + 1). Calcd for C₂₁H₃₀NO₂ (M + H): 328.22765; found: 328.2272.

A VMR strategy for quinolizidine synthesis

Fig. 5 VMR strategy for quinolizidine synthesis.

Paper **RSC Advances**

Scheme 5

Fig. 6 Synthesis versatility of quinolizidine enaminone 27.

Scheme 6 Reduction of guinolizidine enaminone 27.

Similar procedure was applied to synthesize compound 9ah; 14; 15; 17.

(R)-1-((S)-1-(4-Methoxyphenyl)ethyl)-6-methyl-2-pentyl-2,3dihydropyridin-4(1H)-one (9b). Colorless oil (yield, 65%). ¹H-NMR (400 MHz, CDCl₃, 25 °C): $\delta = 0.78$ (t, 3H, J = 6.8 Hz, CH_3 ; 1.05 (m, 1H); 1.18 (m, 5H); 1.25 (m, 2H); 1.56 (d, 3H, J =6.8 Hz, CH₃-); 2.01 (d, 1H, J = 16.58 Hz); 2.20 (dd, 1H, $J_1 = 6$ Hz, $J_2 = 16.51 Hz$); 2.07 (s, 3H, CH_3 -); 3.16 (m, 1H); 3.75 (s, 3H, OCH_3); 4.87 (s, 1H, CH =); 4.98 (q, 1H, J = 6.8 Hz); 6.83 (m, 2H); 7.16 (d, 2H, J = 8.4 Hz). ¹³C-NMR (100 MHz, CDCl₃, 25 °C): $\delta = 14.0$; 17.5; 21.5; 22.6; 25.7; 30.1; 31.7; 38.0; 52.9; 55.3; 55.4; 99.9; 100.0; 114.1; 127.6; 133.4; 159.1; 159.9; 190.6. LC-MS: 100% (purity), m/z: 316 (M + 1). Calcd for $C_{20}H_{30}NO$ (M + H): 316.2276; found: 316.2271.

(R)-2-Heptyl-1-((S)-1-(4-methoxyphenyl)ethyl)-6-methyl-2,3dihydropyridin-4(1H)-one (9c). Colorless oil (yield, 68%). ¹H-NMR (400 MHz, CDCl₃, 25 °C): $\delta = 0.79$ (t, 3H, J = 6.88 Hz, CH_3); 1.16 (m, 12H); 1.56 (d, 3H, J = 7.06 Hz, CH_3 -); 1.98 (d, 1H, J = 16.52 Hz); 2.19 (dd, 1H, $J_1 = 5.94$ Hz, $J_2 = 16.49$ Hz); 2.08 (s, 3H, CH₃-); 3.16 (m, 1H); 3.74 (s, 3H, OCH₃); 4.86 (s,

1H, CH=); 4.97 (q, 1H, J = 6.8 Hz); 6.83 (m, 2H); 7.16 (d, 2H, J= 8.69 Hz). ¹³C-NMR (100 MHz, CDCl₃, 25 °C): δ = 14.1; 17.5; 21.5; 22.6; 25.9; 29.2; 29.5; 30.1; 31.7; 38.0; 52.9; 55.3; 55.4; 99.9; 114.1; 127.7; 133.3; 159.1; 159.9; 190.6. LC-MS: 100% (purity), m/z: 344 (M + 1). Calcd for $C_{22}H_{34}NO_2$ (M + H): 344.2589; found: 344.2584.

(S)-2-((S)-Methoxy(phenyl)methyl)-1-((S)-1-(4-methoxyphenyl)ethyl)-6-methyl-2,3-dihydropyridin-4(1H)-one (9d-I). Colorless oil (yield, 59%). ¹H-NMR (400 MHz, CDCl₃, 25 °C): $\delta = 0.45$ (d, J =7.14 Hz, 3H); 1.33 (dd, J = 5.53 Hz, $J_2 = 16.92$ Hz, 1H); 2.04 (s, 3H); 2.41 (d, J = 16.89 Hz, 1H); 3.11 (s, 3H); 3.26 (ddd, $J_1 = 1.62$ $Hz; J_2 = 5.52 Hz; J_3 = 8.80 Hz, 1H); 3.70 (s, 3H, OCH_3); 4.50 (d, J = 3.50 Hz); 3.70 (s, 3H, OCH_3); 4.50 (d, J = 3.50 Hz); 3.70 (s, 3H, OCH_3); 4.50 (d, J = 3.50 Hz); 3.70 (s, 3H, OCH_3); 4.50 (d, J = 3.50 Hz); 3.70 (s, 3H, OCH_3); 4.50 (d, J = 3.50 Hz); 3.70 (s, 3H, OCH_3); 4.50 (d, J = 3.50 Hz); 3.70 (s, 3H, OCH_3); 4.50 (d, J = 3.50 Hz); 3.70 (s, 3H, OCH_3); 4.50 (d, J = 3.50 Hz); 4.50 (d, J = 3.50 Hz$ 8.80 Hz, 1H); 4.54 (q, J = 7.14 Hz, 1H); 5.09 (s, 1H); 6.74 (m, 2H); 6.98 (m, 2H); 7.18 (dd, $J_1 = 7.25$, $J_2 = 8.83$ Hz; 2H); 7.26 (m, 1H); 7.30 (m, 2H). ¹³C-NMR (100 MHz, CDCl₃, 25 °C): δ = 16.1; 21.7; 36.9; 55.3; 56.1; 57.8; 79.2; 104.4; 114.0; 127.5; 127.6; 128.1; 128.3; 133.5; 140.2; 159.1; 161.5; 192.6. LC-MS: 100% (purity), m/z: 366 (M + 1). Calcd for $C_{23}H_{28}NO_3$ (M + H): 366.2069; found: 366.2065.

(R)-2-((S)-Methoxy(phenyl)methyl)-1-((R)-1-(4-methoxyphenyl)ethyl)-6-methyl-2,3-dihydropyridin-4(1H)-one (9d-II). Colorless oil (yield, 64%). ¹H-NMR (400 MHz, CDCl₃, 25 °C): $\delta = 1.47$ (dd, $J_1 =$ 12.13 Hz; $J_2 = 29.37$ Hz, 1H); 1.69 (d, J = 7.21 Hz, 3H); 1.97 (m, 1H); 2.08 (s, 3H); 3.08 (s, 3H); 3.44 (dd, $J_1 = 6.88$ Hz; $J_2 = 8.23$ Hz, 1H); 3.71 (s, 3H); 4.49 (d, J = 9.19 Hz, 1H); 4.95 (s, 1H); 5.01 (q, J =7.21 Hz, 1H); 6.78 (m, 2H); 7.20 (m, 7H). ¹³C-NMR (100 MHz, CDCl₃, 25 °C): δ = 19.3; 21.7; 36.5; 55.3; 56.9; 57.1; 58.4; 79.6; 102.0; 114.1; 127.6; 127.8; 128.2; 128.6; 134.3; 139.1; 159.0; 161.0; 190.5. LC-MS: 100% (purity), m/z: 366 (M + 1). Calcd for $C_{23}H_{28}NO_3$ (M + H): 366.20692; found: 366.2065.

(R)-2-(4-(Benzyloxy)butyl)-1-((R)-1-(4-methoxyphenyl)ethyl)-6-methyl-2,3-dihydropyridin-4(1H)-one (9e). Colorless oil (yield, 66%). ¹H-NMR (400 MHz, CDCl₃, 25 °C): $\delta = 1.19-1.41$ (m, 4H); 1.48 (m, 2H); 1.52 (d, J = 7.04 Hz, 3H); 1.95 (d, J = 5.09)Hz, 1H); 1.99 (m, 1H); 2.08 (s, 3H); 2.18 (dd, $J_1 = 5.94$ Hz; $J_2 =$ 16.54 Hz); 3.17 (m, 1H); 3.34 (m, 2H); 3.71 (s, 3H, OCH₃); 4.39 (s, 2H); 4.86 (s, 1H); 4.95 (q, J = 7.04 Hz, 1H); 6.81 (m, 2H); 7.13 (m, 2H); 7.23 (m, 5H). ¹³C-NMR (100 MHz, CDCl₃, 25 °C): δ = 14.2; 17.5; 21.5; 22.7; 29.6; 29.8; 37.9; 52.8; 55.3; 55.5; 70.1; 73.0; 100.0; 114.1; 127.5; 127.64; 127.67; 127.70; 127.74; 128.4; 133.3; 138.5; 159.1; 159.9; 190.5. LC-MS: 100% (purity), m/z: 408 (M + 1). Calcd for C₂₆H₃₄NO₃ (M + H): 408.2538; found: 408.2533.

(R)-2-(3-(Benzyloxy)propyl)-1-((R)-1-(4-methoxyphenyl)ethyl)-6-methyl-2,3-dihydropyridin-4(1H)-one (9f). Colorless oil (yield, 75%). ¹H-NMR (400 MHz, CDCl₃, 25 °C): δ = 1.39–1.24 (m, 1H); 1.47-1.40 (m, 1H); 1.55 (d, J = 7.20 Hz, 3H); 1.98-1.94 (m, 2H);

Table 2 Conjugate addition reactions of quinolizidine enaminone 27

Entry	RM	Conditions	Products (yield)
1	MeMgBr	THF (0-RT)	No reaction
2	MeMgBr/CuI (Me₂CuLi)	THF (0-RT)	30a (<5%)
3	MeMgBr	THF/TMS-Cl (3 eq.) (0-RT)	30a (65%)
4	VinylMgBr	THF/TMS-Cl (3 eq.) (0-RT)	30b (57%)
5	AllylMgBr	THF/TMS-Cl (3 eq.) (0-RT)	30c (51%)

2.09 (s, 3H); 2.26–2.13 (m, 1H); 3.29–3.08 (m, 1H); 3.50–3.29 (m, 2H); 3.72 (s, 3H, OCH₃); 4.38 (s, 2H); 4.87 (s, 1H); 4.94 (q, J=7.20 Hz, 1H); 6.79 (m, 2H); 7.10 (m, 2H); 7.21 (m, 5H). ¹³C-NMR (100 MHz, CDCl₃, 25 °C): $\delta=190.4$; 160.0; 159.1; 138.4; 133.3; 128.4; 127.7; 127.6; 127.5; 114.1; 100.0; 72.9; 69.8; 55.5; 55.3; 52.5 38.0; 26.7; 25.9; 21.5; 17.4. LC-MS: 100% (purity), m/z: 394 (M + H). Calcd for $C_{25}H_{32}NO_3$ (M + H): 394.2382 found: 394.2377.

(*R*)-1-((*R*)-1-(4-Methoxyphenyl)ethyl)-6-methyl-2-phenyl-2,3-dihydropyridin-4(1*H*)-one (9g). Colorless oil (yield, 75%). ¹H-NMR (400 MHz, CDCl₃, 25 °C): $\delta = 1.24$ (d, J = 7.2 Hz, 3H), 2.18 (d, J = 16.3 Hz, 1H); 2.23 (s, 3H); 2.71 (dd, $J_1 = 7.53$ Hz; $J_2 = 16.35$ Hz, 1H); 3.739 (s, 3H); 4.42 (m, 1H); 4.96 (s, 1H); 5.12 (q, J = 6.97 Hz, 1H); 6.85 (m, 2H); 7.11 (m, 3H); 7.18 (m, 4H). ¹³C-NMR (100 MHz, CDCl₃, 25 °C): $\delta = 17.7$; 21.6; 42.6;

55.4; 56.0; 56.1; 101.5; 114.2; 126.0; 127.3; 127.6; 128.6; 133.3; 140.6; 159.2; 161.6; 188.9. LC-MS: 100% (purity), *m/z*: 322 (M + 1). Calcd for C₂₁H₂₄NO₂ (M + H): 322.1807; found: 322.1802.

(*S*)-1-((*S*)-1-(4-Methoxyphenyl)ethyl)-6-methyl-2-phenethyl-2,3-dihydropyridin-4(1*H*)-one (9h). Colorless oil (yield, 53%).

¹H-NMR (400 MHz, CDCl₃, 25 °C): δ = 7.19 (m, 2H); 7.13 (m, 1H); 7.04 (d, J = 7.16 Hz, 2H); 6.98 (d, J = 8.65 Hz, 2H); 6.76 (d, J = 8.71 Hz, 2H); 4.90 (q, J = 7.01 Hz, 1H); 4.87 (s, 1H); 3.73 (s, 1H, OCH₃); 3.19 (m, 1H); 2.61 (m, 1H); 2.44 (m, 1H); 2.30 (m, 1H); 2.19 (m, 1H); 2.08 (s, 3H); 1.58 (m, 1H); 1.38 (d, J = 7.07 Hz, 3H, -CH₃).

¹³C-NMR (100 MHz, CDCl₃, 25 °C): δ = 190.3; 160.0; 159.1; 140.9; 132.9; 128.5; 127.7; 126.1; 114.0; 99.9; 55.6; 55.3; 51.5; 37.7; 31.9; 31.7; 21.5; 17.2. LC-MS: 100% (purity), m/z: 335 (M + 1). Calcd for C₂₃H₂₈NO₂ (M + H): 350.2120; found: 350.2115.

(*S*)-1-((*S*)-1-(4-Methoxyphenyl)ethyl)-6-methyl-2-nonyl-2,3-dihydropyridin-4(1*H*)-one (14). Colorless oil (yield, 76%). ¹H-NMR (400 MHz, CDCl₃, 25 °C): δ = 7.16 (d, J = 8.57 Hz, 2H); 6.82 (d, J = 8.79 Hz, 2H); 4.97 (q, J = 6.98 Hz, 1H); 4.86 (s, 1H); 3.73 (s, 3H); 3.16 (m, 1H); 2.19 (dd, J_a = 5.74 Hz, J_b = 16.53 Hz 1H); 2.08 (s, 3H); 1.97 (m, 1H); 1.55 (d, J = 6.98 Hz, 3H); 1.31–1.03 (m, 14H); 0.79 (t, J = 6.90 Hz, 3H). ¹³C-NMR (100 MHz, CDCl₃, 25 °C): δ = 190.6; 159.9; 159.1; 133.3; 127.6; 114.0;

Table 3 Alkylation of quinolizidine enaminone 27

	, any tanton of quinouzing	inte charminone Er	
Entry	RX	32 (yield)	
1	✓ 31a	32a (82%)	
2	Ph∕Br 31b	Ph 32b (78%)	
3	≫ ^{Br} 31c	32c (55%))
4	t-BuOOC ✓ Br 31d	32d (79%)	%)
5	MeBr 31e	Me 32 e (61%)	(o)

99.8; 55.4; 55.3; 52.8; 38.0; 31.8; 30.1; 29.6; 29.5; 29.2; 25.9; 22.6; 21.5; 17.5; 14.1. LC-MS: 100% (purity), m/z: 372 (M + 1). Calcd for $C_{24}H_{38}NO_2$ (M + H): 372.2902; found: 372.2897.

(*R*)-1-((*R*)-1-(4-Methoxyphenyl)ethyl)-6-methyl-2-nonyl-2,3-dihydropyridin-4(1*H*)-one (15). Colorless oil (yield, 69%). ¹H-NMR (400 MHz, CDCl₃, 25 °C): δ = 7.13 (d, J = 8.66 Hz, 2H); 6.80 (d, J = 8.78 Hz, 2H); 4.95 (q, J = 6.98 Hz, 1H); 4.84 (s, 1H); 3.71 (s, 3H); 3.14 (m, 1H); 2.17 (dd, J_a = 6.02 Hz, J_b = 16.62 Hz 1H); 2.06 (s, 3H); 1.95 (m, 1H); 1.53 (d, J = 6.98 Hz, 3H); 1.35–0.94 (m, 14H); 0.77 (t, J = 6.89 Hz, 3H). ¹³C-NMR (100 MHz, CDCl₃, 25 °C): δ = 190.6; 159.9; 159.1; 133.3; 127.6; 114.0; 99.8; 55.4; 55.3; 52.9; 38.0; 31.8; 30.1; 29.6; 26.5; 29.4; 29.2; 25.9; 22.6; 21.5; 17.5; 14.1. LC-MS: 100% (purity), m/z: 372 (M + 1). Calcd for C₂₄H₃₈NO₂ (M + H): 372.2902; found: 372.2897.

(*R*)-1-((*R*)-1-(4-Methoxyphenyl)ethyl)-6-methyl-2-undecyl-2,3-dihydropyridin-4(1*H*)-one (17). Colorless oil (yield, 66%). ¹H-NMR (400 MHz, CDCl₃, 25 °C): $\delta = 7.21$ (d, J = 8.62 Hz, 2H); 6.88 (d, J = 8.62 Hz, 2H); 5.03 (q, J = 6.99 Hz, 1H); 4.92 (s, 1H);

3.97 (s, 3H, OMe); 3.22 (m, 1H); 2.25 (dd, J_1 = 5.88 Hz, J_2 = 16.49 Hz 1H); 2.14 (s, 3H); 2.04 (m, 2H); 1.61 (d, J = 6.99 Hz, 3H); 1.35–1.10 (m, 20H); 0.86 (t, J = 6.86 Hz, 3H). ¹³C-NMR (100 MHz, CDCl₃, 25 °C): δ = 190.6; 159.9; 159.1; 133.3; 127.6; 114.0; 99.9; 55.4; 55.3; 52.9; 38.0; 31.9; 30.1; 29.6; 29.5; 29.4; 29.3; 25.9; 22.7; 21.5; 17.5; 14.1. LC-MS: 100% (purity), m/z: 400 (M + 1). Calcd for $C_{15}H_{42}NO_2$ (M + H): 400.32155; found: 400.3210.

(S)-6-Methyl-2-phenethyl-2,3-dihydropyridin-4(1H)-one (10). To a flask contains (S)-1-((S)-1-(4-methoxyphenyl)ethyl)-6methyl-2-phenethyl-2,3-dihydropyridin-4(1H)-one (9h) (75 mg, 0.21 mmol) added TFA (1.5 mL) and the solution was stirred at room temperature overnight. TFA was removed by vacuum and the residue was re-dissolved in DCM (10 mL). The solution was washed with saturated aqueous solution of sodium bicarbonate and brine and dried over sodium sulfate. After concentration, the crude product was purified by ISCO silica gel chromatography (DCM/methanol) to afford the product as colorless oil (46 mg, 100% yield). 1 H-NMR (400 MHz, CDCl₃, 25 ${}^{\circ}$ C): $\delta = 7.22$ (m, 2H); 7.14 (m, 3H); 4.59 (s, 1H); 4.84 (s, 1H); 3.57 (sex, J = 6.18 Hz, 1H); 2.67 (m, 1H); 2.61 (m, 1H); 2.34 (m, 1H); 2.22 (dd, $J_a = 12.5$ Hz, $J_b = 16.08$ Hz, 1H); 1.93 (m, 1H); 1.85 (m, 1H); 1.81 (s, 3H). ¹³C-NMR (100 MHz, CDCl₃, 25 °C): $\delta = 192.3$; 161.8; 140.8; 128.7; 128.3; 126.4; 99.1; 53.0; 41.1; 35.7; 31.9; 21.2. LC-MS: 100% (purity), m/z: 216 (M + 1). Calcd for $C_{14}H_{18}NO$ (M + H): 216.1388; found: 216.1383.

(2R,4S,6S)-2-Methyl-6-phenethylpiperidin-4-ol (11). To a flask contains methanol solution (0.1 M) of 1(S)-1-((S)-1-(4methoxyphenyl)ethyl)-6-methyl-2-phenethyl-2,3-dihydropyridin-4(1H)-one (9h) (120 mg, 0.34 mmol) added powder of palladium on carbon (12 mg) cautiously under nitrogen stream. The flask was then washed and filled with hydrogen gas. The reaction was stirred at room temperature for 4 h under hydrogen atmosphere and checked by LC-MS until starting material disappeared. After filtration to remove catalyst, the solvent was removed by vacuum and the crude product was purified by ISCO silica gel chromatography (DCM/methanol) to afford the product as gray powder (35 mg, 77% yield). NMR (400 MHz, CD₃OD, 25 °C): δ = 7.10 (m, 2H); 7.14 (m, 2H); 7.05 (t, J = 7.13 Hz, 1H); 3.46 (ddd, J_a = 4.54 Hz, J_b = 7.83 Hz, J_c = 11.08 Hz); 2.56 (m, 3H); 2.44 (m, 1H); 1.93 (m, 1H); 1.79 (m, 1H); 1.68 (m, 1H); 1.58 (m, 1H); 1.01 (d, J = 6.37 Hz, 3H, -Me); 0.91 (m, 2H). ¹³C-NMR (100 MHz, CD₃OD, 25 °C): δ = 143.3; 129.5; 129.4; 126.9; 69.6; 55.6; 51.4; 43.9; 41.5; 39.3; 33.3; 22.1. LC-MS: 100% (purity), m/z: 220 (M + 1). Calcd for C₁₄H₂₂NO (M + H): 220.1701; found: 220.1696.

(2*R*,6*R*)-2-Methyl-6-phenethylpiperidine (12). To Parr shaker reaction vessel contains (0.1 M) of 1(*S*)-1-((*S*)-1-(4-methoxyphenyl) ethyl)-6-methyl-2-phenethyl-2,3-dihydropyridin-4(1*H*)-one (5h) (120 mg, 0.34 mmol) in mixed solvent of methanol (5 mL) and acetic acid (5 mL) added powder of palladium on carbon (36 mg) cautiously under nitrogen stream. The reaction was then carried out by Parr shaker hydrogenation apparatus under 40 psi overnight. After filtration to remove catalyst, the solvent was removed by vacuum and the crude product was redissolved in mixture of chloroform and 2-propanol (3/1, 15 mL). The solution was washed with saturated aqueous solution of sodium bicarbonate and brine and dried over sodium sulfate. After removing the solvent, the crude product was purified by

RSC Advances

ISCO silica gel chromatography (DCM/methanol) to afford the product as colorless oil (yield, 61%) (29 mg, 43% yield). ¹H-NMR (400 MHz, CD₃OD, 25 °C): $\delta = 7.26-6.21$ (m, 5H); 3.03 (m, 1H); 2.93 (m, 1H); 2.72-2.42 (m, 2H); 2.01 (m, 1H); 1.89 (m, 1H); 1.81 (m, 3H); 1.63 (m, 1H); 1.46 (m, 1H); 1.28 (m, 1H); 1.21 (d, J =6.41 Hz, 3H, Me). ¹³C-NMR (100 MHz, CD₃OD, 25 °C): δ = 141.9; 129.7; 129.4; 129.3; 127.4; 58.3; 54.8; 36.9; 32.3; 31.8; 29.2; 23.6; 19.8. LC-MS: 100% (purity), m/z: 204 (M + 1). Calcd for $C_{14}H_{22}N$ (M + H): 204.1752; found: 204.1747.

(2S,4R,6R)-2-Methyl-6-nonylpiperidin-4-ol [(-)-241D]. Starting from 15 similar hydrogenation procedure as in preparing 11 was applied to give (-)-241D as gray solid (yield, 69%). ¹H-NMR (400 MHz, CD₃OD, 25 °C): $\delta = 3.66$ (tt, $J_1 = 4.51$ Hz, $J_2 = 11.09$ Hz); 3.02 (m, 1H); 2.90 (m, 1H); 2.06 (m, 1H); 1.98 (m, 1H); 1.58 (m, 1H); 1.43 (m, 1H); 1.36–1.14 (m, 17H); 1.09 (dd, $J_1 = 12.35$ Hz, $J_2 = 24.34$ Hz, 2H); 0.80 (t, J = 6.85 Hz, 3H, Me). ^{13}C -NMR (100 MHz, CD₃OD, 25 °C): $\delta = 65.8$; 55.2; 51.1; 40.1; 37.6; 33.7; 31.7; 29.2; 29.1; 29.0; 25.0; 22.4; 18.5; 13.1. LC-MS: 100% (purity), m/z: 242 (M + 1). Calcd for $C_{15}H_{32}NO_2$ (M + H): 242.2483; found: 242.2478 $[\alpha]_D = -5.5^{\circ}$ (C, 0.62, MeOH).

(2R,4S,6S)-2-Methyl-6-nonylpiperidin-4-ol [(+)-241D]. Starting from 14 similar hydrogenation procedure as in preparing 11 was applied to give (+)-241D as gray solid (yield, 72%). ¹H-NMR (400 MHz, CD₃OD, 25 °C): $\delta = 3.72$ (tt, $J_1 = 4.42$ Hz, $J_2 = 11.06$ Hz); 3.16 (m, 1H); 3.05 (m, 1H); 2.13 (m, 1H); 2.05 (m, 1H); 1.62 (m, 1H); 1.47 (m, 1H); 1.41-1.18 (m, 17H); 1.14 (m, 2H); 0.80 (t, J=6.84 Hz, 3H, Me). 13 C-NMR (100 MHz, CD $_3$ OD, 25 $^\circ$ C): $\delta=$ 66.5; 56.8; 52.9; 40.6; 38.1; 34.4; 33.1; 30.6; 30.52; 30.55; 30.4; 26.2; 23.8; 19.2; 14.5. LC-MS: 100% (purity), m/z: 242 (M + 1). Calcd for $C_{15}H_{32}NO_2$ (M + H): 242.2483; found: 242.2478 $[\alpha]_D$ = +5.8° (C, 0.69, MeOH).

(2S,6R)-2-Methyl-6-undecylpiperidine (isosolenopsin Starting from 17 similar hydrogenation procedure as in preparing 12 was applied to give isosolenopsin A as colorless oil (yield, 45%). ¹H-NMR (400 MHz, CD₃OD, 25 °C): $\delta = 3.07$ (m, 1H); 2.94 (m, 1H); 1.92 (m, 1H); 1.83 (m, 2H); 1.75-1.52 (m, 1H); 1.52-1.37 (m, 2H); 1.37-1.09 (m, 23H); 0.80 (t, J = 6.73 Hz, 3H, Me). ¹³C-NMR (100 MHz, CD₃OD, 25 °C): $\delta = 57.4$; 53.4; 33.5; 31.7; 30.3; 29.3; 29.2; 29.0; 27.7; 24.7; 22.3; 22.1; 18.1; 13.0. LC-MS: 100% (purity), m/z: 254 (M + 1). Calcd for $C_{17}H_{36}N$ (M + H): 254.2847; found: 254.2842 [α]_D = +9.7° (C, 0.71, CHCl₃).

(R)-2-(4-Chlorobutyl)-1-((R)-1-(4-methoxyphenyl)ethyl)-6-methyl-2,3-dihydropyridin-4(1H)-one (25). To a round-bottom flask containing (R)-1-(4-methoxyphenyl)ethanamine (6a) (151 mg, 1 mmol, 1 eq.) in dried DCM (0.1 M) solution was added 4-AMS (500 mg mmol⁻¹) followed by 5-chloropentanal (24) (120 mg, 1 mmol, 1 eq.). After stirring at room temperature for 10 min, (1-methoxybuta-1,3-dienyloxy)-trimethyl-silane (7) (292 mg, 1.2 mmol, 1.2 eq.) was added and the solution was cooled to -78 °C. Tin(II) triflate (412 mg, 1 mmol, 1 eq.) was then added and the reaction was stirred at this temperature for 8 h. The reaction temperature was raised to 0 °C and kept at the same temperature overnight. LC-MS analysis of the reaction showed the disappearance of the starting materials. The reaction was quenched with saturated aqueous solution of sodium bicarbonate and solids were removed via filtration. The reaction mixture was then extracted with DCM (5 imes15 mL). The combined organic phase was treated with acetic acid

and the resulting solution was stirred at room temperature for 1 h until only compound 25 was observed from LC-MS analysis. The solution was then basified by treating with saturated aqueous solution of sodium bicarbonate, washed with brine and dried over sodium sulfate. After concentration, the crude product was purified by ISCO silica gel chromatography (ethyl acetate/hexane) to afford the product 25 (ethyl acetate/hexane = 5/1) as colorless oil (184 mg, 55% yield). ¹HNMR (400 MHz, CDCl₃, 25 °C): δ = 7.23 (d, I= 8.45 Hz, 2H; 6.90 (d, J = 8.80 Hz, 2H); 5.06 (q, J = 7.01 Hz, 1H); 4.95 (s, 1H); 3.82 (s, 3H); 3.50 (m, 2H); 3.25 (m, 1H); 2.28 (dd, $J_a =$ 5.94 Hz, $J_b = 16.58 \text{ Hz}$ 1H); 2.17 (s, 3H); 2.04 (m, 1H); 1.72 (m, 2H)1.63 (d, J = 7.07 Hz, 3H); 1.38 (m, 3H). ¹³C-NMR (100 MHz, CDCl₃, 25 °C): δ = 190.4; 160.0; 159.2; 133.1; 127.6; 114.1; 100.1; 55.4; 55.3; 52.7; 44.7; 38.0; 32.3; 29.4; 23.4; 21.5; 17.5. LC-MS: 100% (purity), m/z: 336 (M + 1). Calcd for C₁₉H₂₇ClNO₂ (M + H): 336.1730; found: 336.1725.

(R)-2-(4-Chlorobutyl)-6-methyl-2,3-dihydropyridin-4(1H)-one (26). To the flask containing compound 25 (120 mg, 0.36 mol) was added TFA (2 mL, 99%) at room temperature and the resulting solution was stirred at room temperature overnight. TFA was removed by vacuum and the residue was re-dissolved in DCM (10 mL). The solution was washed with saturated aqueous solution of sodium bicarbonate and brine and dried over sodium sulfate. After concentration, the crude product was purified by ISCO silica gel chromatography (DCM/methanol) to afford the product as colorless oil (72 mg, 100% yield). ¹H-NMR (400 MHz, CDCl₃, 25 °C): $\delta = 4.59$ (s, 1H); 4.85 (s, 1H); 3.56 (m, 1H); 2.32 (dd, $J_a = 5.07$ Hz, $J_b = 16.10$ Hz); 2.19 (dd, $J_a = 5.07$ Hz, $J_{\rm b} = 16.10 \text{ Hz}$; 1.91 (s, 3H); 1.74 (m, 2H); 1.63 (m, 2H); 1.53 (m, 1H); 1.47 (m, 2H). ¹³C-NMR (100 MHz, CDCl₃, 25 °C): δ = 190.4; 162.5; 98.8; 53.0; 44.7; 40.9; 33.3; 32.2; 22.7; 21.2. LC-MS: 100% (purity), m/z: 202 (M + 1). Calcd for $C_{10}H_{17}ClNO$ (M + H): 202.0998; found: 202.0993.

(R)-4-Methyl-7,8,9,9a-tetrahydro-1H-quinolizin-2(6H)-one (27). To the DMF (5 mL) solution contained 26 (95 mg, 0.47 mmol) added sodium hydride (56 mg, 60%, 3 eq.) at 0 °C. The reaction was stirred at this temperature for 2 h before quenched with saturated aqueous solution of ammonium chloride. The mixture was extracted with ethyl acetate (10 mL \times 3) and the combined organic phase was washed with brine and dried over sodium sulfate. After concentration, the crude product was purified by ISCO silica gel chromatography (ethyl acetate/hexane) to afford the product as colorless oil (74 mg, 96% yield). ¹H-NMR (400 MHz, CDCl₃, 25 °C): δ = 4.93 (s, 1H); 3.72 (m, 1H); 3.29 (m, 1H); 2.75 (td, $J_1 = 2.91 \text{ Hz}, J_2 = 12.76 \text{ Hz}, 1\text{H}$; 1.92 (s, 3H); 1.80 (m, 1H); 1.68 (m, 1H); 1.58 (m, 2H); 1.43 (m, 2H). ¹³C-NMR (100 MHz, CDCl₃, 25 °C): $\delta = 191.5$; 163.0; 101.8; 58.6; 48.1; 42.9; 31.4; 25.8; 23.7; 21.2. LC-MS: 100% (purity), m/z: 166 (M + 1). Calcd for $C_{10}H_{16}NO$ (M + H): 166.1231; found: 166.1226.

(2R,4S,9aR)-4-Methyloctahydro-1H-quinolizin-2-ol (28). To flask contained 27 (85 mg, 0.51 mmol) in THF (3 mL) added L-selectride (1 M in THF, 2.5 mL, 5 eq.) at 0 °C. The reaction was stirred at this temperature for 2 h before quenching with saturated aqueous solution of sodium bicarbonate. The mixture was extracted with mixed solvent (chloroform/isopropanol 3/1, 10 mL \times 3) and the combined organic phase was washed with brine and dried over sodium sulfate. After concentration, the

crude product was purified by ISCO silica gel chromatography (chloroform/methanol) to afford the product as colorless oil (70 mg, 81% yield). $^1\text{H-NMR}$ (400 MHz, CDCl $_3$, 25 °C): $\delta=4.13$ (s, 1H, OH); 3.70 (m, 1H); 3.63 (m, 1H); 3.40 (m, 1H); 3.31 (m, 1H); 2.45 (m, 1H); 2.25 (t, J=13.21 Hz, 1H); 2.15 (t, J=12.77 Hz, 2H); 1.94 (m, 3H); 1.83 (m, 2H); 1.69 (m, 1H); 1.44 (m, 1H); 1.42 (d, J=6.30 Hz, 3H, Me). $^{13}\text{C-NMR}$ (100 MHz, CDCl $_3$, 25 °C): $\delta=61.8$; 59.2; 56.1; 50.9; 38.6; 37.4; 30.1; 23.3; 22.6; 17.2. LC-MS: 100% (purity), m/z: 170 (M+1). Calcd for $\mathrm{C_{10}H_{20}NO}$ (M+H): 170.1544; found: 170.1539.

(4S,9aR)-4-Methylhexahydro-1H-quinolizin-2(6H)-one (29). To flask contained 27 (68 mg, 0.41 mmol) in THF (2 mL) added BF₃·OEt₂ (57 μ L, 1.1 eq.) at -78 °C. The reaction was stirred at this temperature for 10 min before adding lithium triethylborohydride "super hydride" (1 M in THF, 0.49 mL, 1.2 eq.). After stirring at -78 °C for 2 h, the reaction was quenched with saturated aqueous solution of sodium bicarbonate. The mixture was extracted with mixed solvent (chloroform/isopropanol 3/1, 10 mL \times 3) and the combined organic phase was washed with brine and dried over sodium sulfate. After concentration, the crude product was purified by ISCO silica gel chromatography (chloroform/methanol) to afford the product as colorless oil (37 mg, 54% yield). ¹H-NMR (400 MHz, CDCl₃, 25 °C): $\delta = 3.27$ (m, 1H); 2.33 (m, 3H): 2.22 (m, 2H): 2.12 (t, J = 10.95 Hz, 1H); 1.78 (m, 1H); 1.68 (m, 2H); 1.58 (m, 2H); 1.36 (m, 1H); 1.20 (m, 1H); 1.14 (d, J = 5.72 Hz, 3H). ¹³C-NMR (100 MHz, CDCl₃, 25 °C): $\delta =$ 207.9; 61.6; 58.8; 50.5; 49.2; 48.1; 33.6; 25.4; 23.4; 20.2. LC-MS: 100% (purity), m/z: 168 (M + 1). Calcd for $C_{10}H_{18}NO$ (M + H): 168.1388; found: 168.1383 $[\alpha]_D = -16.9^\circ$ (C, 0.28, CHCl₃).

(R)-4,4-Dimethylhexahydro-1H-quinolizin-2(6H)-one To the flask contained 27 (42 mg, 0.25 mmol) in dried THF (2 mL) added TMS-Cl (80 μ L, 3 eq.) at 0 °C. The reaction was stirred at this temperature for 10 min before adding methyl magnesium bromide (2 M in ethyl ether, 0.18 mL, 1.5 eq.). After stirring at 0 °C for 2 h, the reaction temperature was allowed to raise to room temperature overnight. Quenched by saturated aqueous solution of sodium bicarbonate the reaction mixture was extracted with mixed solvent (chloroform/isopropanol 3/1, 10 mL \times 3) and the combined organic phase was washed with brine and dried over sodium sulfate. After concentration, the crude product was purified by ISCO silica gel chromatography (chloroform/methanol) to afford the product as colorless oil (28 mg, 65% yield). 1 H-NMR (400 MHz, CDCl₃, 25 ${}^{\circ}$ C): $\delta = 3.0$ (m, 1H); 2.48 (m, 2H); 2.15 (m, 2H); 1.99 (m, 2H); 1.60 (m, 3H); 1.45 (m, 1H); 1.27 (m, 1H); 1.10 (m, 1H); 1.13 (s, 3H); 0.81 (s, 3H). ¹³C-NMR (100 MHz, CDCl₃, 25 °C): δ = 209.1; 58.0; 55.8; 55.4; 48.2; 45.4; 34.6; 29.6; 25.8; 23.8; 15.8. LC-MS: 100% (purity), m/z: 182 (M + 1). Calcd for $C_{11}H_{20}NO(M + H)$: 182.1544; found: 182.1539.

(4*S*,9*aR*)-4-Methyl-4-vinylhexahydro-1*H*-quinolizin-2(6*H*)-one (30b). Starting from 27 similar addition procedure as in preparing 30a was applied to give 30b as colorless oil (yield, 57%). 1 H-NMR (400 MHz, CDCl₃, 25 $^{\circ}$ C): δ = 5.82 (dd, J = 17.6, 11.1 Hz, 1H), 5.16 (d, J = 11.0 Hz, 1H), 5.01 (d, J = 17.5 Hz, 1H), 3.07 (m, 1H), 2.84–2.42 (m, 2H), 2.37 (m, 1H), 2.21 (m, 3H), 1.93–1.72 (m, 1H), 1.63 (m, 3H), 1.57–1.41 (m, 1H), 1.27 (s, 3H), 1.18 (m, 2H). 13 C-NMR (100 MHz, CDCl₃, 25 $^{\circ}$ C): δ = 209.1; 137.1; 116.6; 61.2; 55.1; 51.9; 47.5; 44.9; 34.3; 26.5; 25.8; 23.3. LC-MS:

100% (purity), m/z: 194 (M + 1). Calcd for $C_{12}H_{19}NO$ (M + H): 194.1544; found: 194.1540.

(4*R*,9*aR*)-4-Allyl-4-methylhexahydro-1*H*-quinolizin-2(6*H*)-one (30c). Starting from 27 similar addition procedure as in preparing 30a was applied to give 14 as colorless oil (yield, 57%).
¹H NMR (400 MHz, CDCl₃, 25 °C) δ = 5.81–5.44 (m, 1H), 5.15–4.86 (m, 2H), 3.22–2.97 (m, 1H), 2.80–2.59 (m, 1H), 2.31 (s, 2H), 2.27–2.11 (m, 3H), 2.08–1.97 (m, 2H), 1.66 (d, *J* = 9.9 Hz, 2H), 1.31 (m, 2H), 1.18 (s, 2H), 1.15 (s, 3H).
¹³C NMR (101 MHz, CDCl₃, 25 °C) δ = 209.33; 133.84; 118.66; 61.04; 55.72; 51.76; 48.41; 45.38; 35.28; 34.28; 26.91; 26.43; 24.13. LC-MS: 100% (purity), *m/z*: 208 (M + 1). Calcd for C₁₃H₂₂NO (M + H): 208.1701; found: 208.1696.

(1R,9aR)-1-Benzyl-4-methyl-7,8,9,9a-tetrahydro-1H-quinolizin-2(6H)-one (32b). To the flask contained 27 (42 mg, 0.25 mmol) in dried THF (2 mL) added LiHMDS (0.3 mL, 1 M in THF) at -78 °C. The reaction was stirred at this temperature for 1 h before adding benzyl bromide (88 μL, 3 eq.). After stirring at 0 °C for 2 h, the reaction temperature was allowed to raise to room temperature overnight. Quenched by saturated aqueous solution of sodium bicarbonate the reaction mixture was extracted with mixed solvent (ethyl acetate, 10 mL \times 3) and the combined organic phase was washed with brine and dried over sodium sulfate. After concentration, the crude product was purified by ISCO silica gel chromatography (hexane/ethyl acetate) to afford the product as colorless oil (49 mg, 78% yield). H NMR (400 MHz, CDCl₃, 25 °C) δ = 7.27-7.19 (m, 2H); 7.19-7.08 (m, 3H); 5.00 (s, 1H); 3.91-3.57 (m, 1H); 3.29 (dd, J = 5.9, 4.2 Hz, 1H); 2.97-2.62 (m, 3H); 2.62-2.38 (m, 3H);2.23 (dd, J = 16.5, 10.7 Hz, 1H); 1.86–1.73 (m, 1H); 1.73–1.49 (m, 3H); 1.49–1.31 (m, 2H). 13 C NMR (101 MHz, CDCl₃, 25 $^{\circ}$ C) δ = 191.2; 165.2; 139.7; 128.1; 127.8; 126.1; 100.5; 76.9; 58.2;47.6; 42.2; 35.1; 33.9; 31.0; 25.5; 23.3. LC-MS: 100% (purity), m/ z: 256 (M + 1). Calcd for C₁₇H₂₂NO (M + H): 256.1701; found: 256.1696.

Starting from 27 similar alkylation procedure as in preparing 32b was applied to give 32a; 32c; 32d and 32e.

(*R*)-4-Ethyl-7,8,9,9*a*-tetrahydro-1*H*-quinolizin-2(6*H*)-one (32a). Colorless oil (yield, 82%). ¹H NMR (400 MHz, CDCl₃, 25 °C) δ = 4.99 (s, 1H); 3.81–3.63 (m, 1H); 3.39–3.19 (m, 1H); 2.74 (td, J = 12.7, 2.9 Hz, 1H); 2.47 (dd, J = 16.6, 5.9 Hz, 1H); 2.29–2.18 (m, 2H); 2.17 (m, 1H); 1.79 (m 1H); 1.74–1.66 (m, 1H); 1.66–1.54 (m, 2H); 1.54–1.32 (m, 2H); 1.09 (t, J = 7.5 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃, 25 °C) δ = 191.2; 167.6; 99.4; 58.2; 47.3; 42.0; 31.0; 26.4; 25.5; 23.3; 11.9. LC-MS: 100% (purity), m/z: 180 (M + 1). Calcd for C₁₁H₁₈NO (M + H): 180.1388; found: 180.1383.

(*R*)-4-(But-3-en-1-yl)-7,8,9,9*a*-tetrahydro-1*H*-quinolizin-2(6*H*)-one (32c). Colorless oil (yield, 55%). ¹H NMR (400 MHz, CDCl₃, 25 °C) δ = 5.96–5.72 (m, 1H); 5.21–4.93 (m, 3H); 3.94–3.64 (m, 1H); 3.49–3.18 (m, 1H); 2.82 (td, J = 12.7, 2.8 Hz, 1H); 2.53 (dd, J = 16.4, 5.8 Hz, 1H); 2.45–2.19 (m, 3H); 1.92–1.83 (m, 1H); 1.83–1.74 (m, 3H); 1.69 (dd, J = 6.8, 3.6 Hz, 2H); 1.61–1.41 (m, 2H). ¹³C NMR (101 MHz, CDCl₃, 25 °C) δ = 191.8; 165.7; 136.5; 115.9; 101.1; 58.6; 48.1; 42.8; 33.2; 32.1; 31.5; 26.0; 23.8. LC-MS: 100% (purity), m/z: 205 (M + 1). Calcd for C₁₃H₂₀NO (M + H): 206.1545; found: 206.1540.

(*R*)-tert-Butyl-3-(2-oxo-2,6,7,8,9,9*a*-hexahydro-1*H*-quinolizin-4-yl)propanoate (32d). Colorless oil (yield, 79%). ¹H NMR (400 MHz, CDCl₃, 25 °C) δ = 4.92 (s, 1H); 3.89–3.63 (m, 1H); 3.47–3.17 (m, 1H); 2.88–2.63 (m, 1H); 2.56–2.41 (m, 3H); 2.41–2.34 (m, 2H); 2.21 (dd, *J* = 16.4, 10.4 Hz, 1H); 1.86–1.75 (m, 1H); 1.75–1.65 (m, 1H); 1.65–1.52 (m, 2H); 1.52–1.39 (m, 2H); 1.38 (s, 9H, *t*-Bu). ¹³C NMR (101 MHz, CDCl₃, 25 °C) δ = 190.7; 170.2; 163.5; 99.6; 80.2; 57.7; 47.0; 41.7; 32.4; 30.3; 27.6; 27.1; 27.1; 24.9; 22.7. LC-MS: 100% (purity), *m/z*: 280 (M + 1). Calcd for C₁₆H₂₆NO₃ (M + H): 280.1913; found: 280.1908.

(*R*)-4-(Pent-3-yn-1-yl)-7,8,9,9*a*-tetrahydro-1*H*-quinolizin-2(6*H*)-one (32e). Colorless oil (yield, 61%). ¹H NMR (400 MHz, CDCl₃, 25 °C) δ = 4.95 (s, 1H); 3.91–3.57 (m, 1H); 3.39–3.17 (m, 1H); 2.74 (td, J = 12.8, 2.8 Hz, 1H); 2.50–2.33 (m, 3H); 2.33–2.27 (m, 2H); 2.23 (dd, J = 16.5, 10.7 Hz, 1H); 1.89–1.74 (m, 1H); 1.71 (s, 3H); 1.69–1.65 (m, 1H); 1.65–1.51 (m, 2H); 1.51–1.33 (m, 2H). ¹³C NMR (101 MHz, 25 °C) δ = 191.9; 164.5; 101.0; 77.2; 76.9; 58.6; 48.1; 42.8; 33.2; 31.4; 25.9; 23.8; 17.9; 3.5. LC-MS: 100% (purity), m/z: 218 (M + 1). Calcd for $C_{14}H_{20}NO$ (M + H): 218.1545; found: 218.1541.

Acknowledgements

Dedicated to ICCAS Emeritus Professor Dong Wang on the occasion of his 74th birthday. The author thanks Dr Perry Gordon and Mr Kevin Johnson for providing high resolution mass spectrometry data (HRMS) of all reported compounds. The author sincerely appreciates the management supports from Dr John Tellew; Dr Shifeng Pan; Dr Dean P. Phillips and Dr Valentina Molteni. The author is also grateful to Dr John Tellew and Dr Yahu A. Liu for helpful discussions and proof reading.

References

- 1 G. M. Struntz and J. A. Findlay, in *The Alkaloids*, ed. A. Brossi, Academic, New York, 1985, vol. 26, pp. 89–193.
- 2 R. D. Taylor, M. MacCoss and A. D. G. Lawson, J. Med. Chem., 2014, 57, 5845–5859.
- 3 For recent reviews see: (a) S. Kallstrom and R. Leino, *Bioorg. Med. Chem.*, 2008, 681; (b) M. G. P. Buffat, *Tetrahedron*, 2004, 1701; (c) F.-X. Felpin and J. Lebreton, *Eur. J. Org. Chem.*, 2003, 3693; (d) S. Laschat and T. Dickner, *Synthesis*, 2000, 1781.
- 4 Y. Ying, H. Kim and J. Hong, *Org. Lett.*, 2011, **13**, 796–799 and references therein.
- 5 G. M. Strunz and J. A. Findlay, Pyridine and piperidine alkaloids, in *The Alkaloids*, ed. A. Brossi, Academic Press, New York, 1985, vol. 26, pp. 89–174.
- 6 For general reviews see: (a) P. M. Dewick, in *Medicinal Natural Products, A Biosynthetic Approach*, Wiley, 3rd revised edn, 2002; (b) E. Szoke, E. Lemberkovics and L. Kursinszki, in *Natural Products, Alkaloids Derived from Lysine: Piperidine Alkaloids*, ed. K. G. Ramawat and J. M. Merillon, Springer, 2013, pp. 301–343.
- 7 Δ^{1} -Piperideine is unstable and exists in solution as a complex diastereoisomeric mixture of a trimeric (major component) and monomeric form: (*a*) A. Rouchaud and J. C. Braekman, *Eur. J. Org. Chem.*, 2009, 2666; (*b*)

- C. Darwich, M. Elkhatib, G. Steinhauser and H. Delalu, *Helv. Chim. Acta*, 2009, **92**, 98; (*c*) J. C. Guillemin, J. M. Denis, M. C. Lasne and J. L. Ripoll, *Tetrahedron*, 1988, **44**, 4447–4455.
- 8 Biomimetic synthesis of piperidine-type alkaloids by using Δ^1 -piperideine was reported by Bella *et al.* see: M. R. Monaco, P. Renzi, D. M. Scarpino Schietroma and M. Bella, *Org. Lett.*, 2011, 13, 4546–4549.
- 9 (a) D. L. Comins and J. D. Brown, Tetrahedron Lett., 1986, 27, 4549; (b) D. L. Comins and L. A. Morgan, Tetrahedron Lett., 1991, 32, 5919; (c) D. L. Comins and H. Hong, J. Org. Chem., 1993, 58, 5035; (d) D. L. Comins and N. R. Benjelloun, Tetrahedron Lett., 1994, 35, 829; (e) D. L. Comins and Y. M. Zhang, J. Am. Chem. Soc., 1996, 118, 12248; (f) D. L. Comins, A. H. Libby, R. S. Al-awar and C. J. Foti, J. Org. Chem., 1999, 64, 2184; (g) D. L. Comins, M. J. Sandelier and T. A. Grillo, J. Org. Chem., 2001, 66, 6829; (h) R. Badorrey, C. Cativiela, M. D. Diaz de Villegas and J. A. Galvez, Tetrahedron, 2002, 58, 341; (i) N. S. Josephsohn, M. L. Snapper and A. H. Hoveyda, *J. Am.* Chem. Soc., 2003, 125, 4018; (j) D. L. Comins and J. J. Sahn, Org. Lett., 2005, 7, 5227; (k) R. T. Yu and T. Rovis, J. Am. Chem. Soc., 2006, 128, 12370; (l) B. J. Turunen and G. I. Georg, J. Am. Chem. Soc., 2006, 128, 8702; (m) C. A. Newman, J. C. Antilla, P. Chen, A. V. Predeus, L. Fielding and W. D. Wulff, J. Am. Chem. Soc., 2007, 129, 7216; (n) T. Andreassen, T. Haaland, L. K. Hansen and O. R. Gautun, Tetrahedron Lett., 2007, 48, 8413; (o) R. K. Friedman and T. Rovis, J. Am. Chem. Soc., 2009, 131, 10775; (p) M. J. Niphakis, B. J. Turunen and G. I. Georg, J. Org. Chem., 2010, 75, 6793; (q) For a recent review: J. A. Bull, J. J. Mousseau, G. Pelletier and A. B. Charette, Chem. Rev., 2012, 112, 2642-2713.
- 10 As a dianion synthons with 1,2-dielectrophiles (a) P. Langer and V. Kohler, Chem. Commun., 2000, 1653-1654; (b) E. Ulah, B. Appel, C. Fischer and P. Langer, Tetrahedron Lett., 2006, 62, 9694-9700; (c) P. Langer and K. Valentin, Org. Lett., 2000, 2, 1597-1599; (d) E. Ullah, B. Appel, C. Fischer and P. Langer, Tetrahedron, 2006, 62, 9694-9700; (e) E. Holtz, V. Koehler, B. Appel and P. Langer, Eur. J. Org. Chem., 2005, 532-542; (f) I. Hussain, M. A. Yawer, M. Lau, T. Pundt, C. Fischer, H. Gorls and P. Langer, Eur. J. Org. Chem., 2008, 503-518; (g) G. Bose, E. Ullah and P. Langer, Chem.-Eur. J., 2004, 10, 6015-6028; (h) M. Lubbe, R. Klassen, T. Trabhardt, A. Villinger and P. Langer, Synlett, 2008, 15, 2331; (i) E. Bellur, H. Goerls and P. Langer, J. Org. Chem., 2005, 70, 4751-4761. In vinylogous aldol reaction: (j) S. Danishefsky, D. F. Harvey, G. Qualich and B. J. Uang, J. Org. Chem., 1984, 49, 393-395; (k) V. Henryon, L. W. Liu, R. Lopez, J. Prunet and J.-P. Ferezou, Synthesis, 2001, 2401–2414; (l) G. Bose and P. Langer, Synlett, 2005, 1021-1023; (m) S. E. Denmark and J. R. Heemstra Jr, J. Org. Chem., 2007, 72, 5668-5688. In bromination: (n) J. L. Shamshina and T. S. Snowden, Tetrahedron Lett., 2007, 48(22), 3767-3769; (o) A. Barbero and F. J. Pulido, Synthesis, 2004, 401-404.

- 11 Instead of using 7, the addition to aldimines by the dianion generated from the 1,3-diketones was once reported see:
 J. Jiao, Y. Zhang and R. A. Flowers II, *Synlett*, 2006, 3355–3357.
- 12 A. Solladi-Cavallo, C. Marsol, M. Yakoub, K. Azyat, A. Klein, M. Roje, C. Suteu, T. B. Freedman, X. Can and L. A. Nafie, J. Org. Chem., 2003, 68, 7308.
- 13 The absolute stereochemistry of VMR was determined by comparing spectrum and optical rotation data of known compounds in literature. See: M. J. Niphakis, B. J. Turunen and G. I. Georg, *J. Org. Chem.*, 2010, 75, 6793–6805. See also ref. 18.
- 14 Hydrogenations on 2,6- disubstituted 2,3-dihydropyridone are known to be stereospecific in literature. For some examples: (*a*) D. Ma and H. Sun, *Org. Lett.*, 2000, 2, 2503–2505; (*b*) N. Gouault, M. Le Roch, G. de. C. Pinto and M. David, *Org. Biomol. Chem.*, 2012, **10**, 5541–5546.
- 15 (a) E. S. Tasber and R. M. Garbaccio, *Tetrahedron Lett.*, 2003, 44, 9185–9188; (b) E. M. Fry, *J. Org. Chem.*, 1963, 1869–1874.
- 16 J. A. Findlay, W. H. J. Tam and J. Krepinsky, Can. J. Chem., 1978, 56, 613.
- 17 M. W. Edwards, J. W. Daly and C. W. Myer, J. Nat. Prod., 1988, 51, 1188.
- 18 N. Gouault, M. L. Roch, G. C. Pinto and M. David, *Org. Biomol. Chem.*, 2012, **10**, 5541-5546 and reference therein.
- 19 For other reported asymmetric synthesis on (+)-241D and/or (-)-241D in literature: (a) V. H. Vu, F. Louafi, N. Girard, R. Marion, T. Roisnel, V. Dorcet and J.-P. Hurvois, J. Org. Chem., 2014, 79, 3358-3373; (b) I. Abrunhosa-Thomas, A. Plas, A. Vogrig, N. Kandepedu, P. Chalard and Y. Troin, J. Org. Chem., 2013, 78, 2511-2526; (c) N. Saha and S. K. Chattopadhyay, J. Org. Chem., 2012, 77, 11056-11063; (d) N. Gouault, M. Le Roch, G. de. C. Pinto and M. David, Org. Biomol. Chem., 2012, 10, 5541-5546; (e) N. Gouault, M. Le Roch, G. de. C. Pinto and M. David, Org. Biomol. Chem., 2012, 10, 5541-5546; (f) R. V. N. S. Murali and S. Chandrasekhar, Tetrahedron Lett., 2012, 53, 3467-3470; (g) K. Damodar and B. Das, Synthesis, 2012, 44, 83-86; (h) C. K. R. Sateesh, R. G. Venkateswar, G. Shankaraiah, K. Suresh Babu and J. Madhusudana Rao, Tetrahedron Lett., 2010, 51, 1114-1116; (i) C. Gnamm, C. M. Krauter, K. Broedner and G. Helmchen, Chem.-Eur. J., 2009, 15, 2050-2054; (j) J. Monfray, Y. Gelas-Mialhe, J.-C. Gramain and R. Remuson, Tetrahedron: Asymmetry, 2005, 16, 1025-1034; (k) F. A. Davis, B. Chao and A. Rao, Org. Lett., 2000, 2, 3169-3171; (l) D. Ma and H. Sun, Org. Lett., 2000, 2, 2503-2505.
- 20 For some bioactivity studies on isosolenpsin A see: (a) S. Leclercq, D. Daloze and J. C. Braekman, Org. Prep. Proced. Int., 1996, 28, 499; (b) G. Howell, J. Butler, R. D. de Shazo, J. M. Farley, H.-L. Liu, N. P. D. Nanayakkara, A. Yates, G. B. Yi and R. W. Rockhold, Ann. Allergy, Asthma, Immunol., 2005, 94, 380–386; (c) G. B. Yi, D. McClendon, D. Desaiah, J. Goddard, A. Lister, J. Moffitt, R. K. Vander Meer, R. Deshazo, K. S. Lee and R. W. Rockhold, Int. J. Toxicol., 2003, 22, 81–86.

- 21 For other reported asymmetric synthesis of isosolenpsin A see: (a) R. S. C. Kumar, E. Sreedhar, G. Venkateswar Reddy, K. Suresh Babu and J. Madhusudana Rao, Tetrahedron: Asymmetry, 2009, 20, 1160-1163; (b) J. C. Gonzalez-Gomez, F. Foubelo and M. Yus, Synlett, 2008, 2777-2780; (c) H. M. T. B. Herath and N. P. D. Nanayakkara, J. Heterocycl. Chem., 2008, 45, 129-136; (d) N. Girard, J.-P. Hurvois, L. Toupet and C. Moinet, Synth. Commun., 2005, 35, 711-723; (e) J. Monfray, Y. Gelas-Mialhe, J.-C. Gramain and R. Remuson, Tetrahedron: Asymmetry, 2005, 16, 1025-1034; (f) S. Ciblat, P. Besse, G. I. Papastergiou, H. Veschambre, J.-L. Canet and Y. Troin, Tetrahedron: Asymmetry, 2000, 11, 2221–2229; (g) H. Poerwono, K. Higashiyama, T. Yamauchi, H. Kubo, S. Ohmiya and H. Takahashi, Tetrahedron, 1998, 54, 13955-13970; (h) C. W. Jefford and J. B. Wang, Tetrahedron Lett., 1993, 34, 2911-2914.
- 22 (a) A. S. Howard and J. P. Michael, Simple indolizidine and quinolizidine alkaloids, in *The Alkaloids*, ed. A. Brossi, Academic Press, New York, 1986, pp. 183–308; (b) J. W. Daly, H. M. Garraffo and T. F. Spande, Alkaloids from amphibian skins, in *Alkaloids: Chemical and Biological Perspectives*, ed. S. W. Pelletier, Pergamon Press, Amsterdam, 1999, pp. 1–161; (c) I. Garnett, The quinolizidine alkaloids, in *Rodd's Chemistry of Carbon Compounds*, ed. M. Sainsbury, Elsevier, Amsterdam, 1998, pp. 181–239.
- 23 A. M. Lourenço, P. Máximo, L. M. Ferreira and M. M. A. Pereira, Indolizidine and quinolizidine alkaloids: structure and bioactivity, in *Studies in Natural Product Chemistry Bioactive Natural Products Part H*, ed. A. Rahman, Elsevier, Amsterdam, 2002, vol. 27, pp. 233–298.
- 24 As a notable example, quinolizidine alkaloid (—)-cytisine was once identified as a potent partial nicotinic acetylcholine receptor agonist. Inspired by the bioactivity of (—)-cytisine, researchers at Pfizer successfully developed novel smoking cessation drug varenicline approved by FDA in 2006. See J. W. Coe, P. R. Brooks, M. G. Vetelino, M. C. Wirtz, E. P. Arnold, J. Huang, S. B. Sands, T. I. Davis, L. A. Lebel, C. B. Fox, A. Shrikhande, J. H. Heym, E. Schaeffer, H. Rollema, Y. Lu, R. S. Mansbach, L. K. Chambers, C. C. Rovetti, D. W. Schulz, F. D. Tingley and B. T. O'Neill, *J. Med. Chem.*, 2005, 48, 3474.
- 25 (a) M. Wink and T. Hartmann, *Plant Physiol.*, 1982, 70, 74–77; (b) For a review on quinolizidine alkaloid biosynthesis:
 S. Bunsupa, M. Yamazaki and K. Saito, *Front. Plant Sci.*, 2012, 3, 239.
- 26 The stereochemistry results of the reduction on cyclic enaminone were established by literature precedents see: (a) W. Wysocka and A. Przybyl, Monatsh. Chem., 2001, 132, 973–984; (b) D. L. Comins, X. Zheng and R. R. Goehring, Org. Lett., 2002, 4, 1611–1613; (c) R. T. Yu and T. Rovis, J. Am. Chem. Soc., 2006, 128, 12370–12371; (d) T. Nagasaka, H. Yamamoto, H. Hayashi, M. Watanabe and F. Hamaguchi, Heterocycles, 1989, 29, 155–164.
- 27 Similar conditions have also been applied in the regionselective reduction of 2,3-dihydropyridones, M. Dax,

- R. Frohlich, D. Schepmann and B. Wunsch, Eur. J. Org. Chem., 2008, 6015–6028.
- 28 For the recent asymmetric synthesis of (-)-epimyrtine see: (a) T. T. H. Trinh, K. H. Nguyen, P. de A. Amaral and N. Gouault, *Beilstein J. Org. Chem.*, 2013, **9**, 2042–2047; (b) Y. Ying, H. Kim and J. Hong, *Org. Lett.*, 2011, **13**, 796–799; (c) S. M. Amorde, A. S. Judd and S. F. Martin, *Org. Lett.*, 2005, 7, 2031–2033; (d) F. A. Davis, Y. Zhang and G. Anilkumar, *J. Org. Chem.*, 2003, **68**, 8061–8064; (e) D. Gardette, Y. Gelas-Mialhe, J.-C. Gramain, B. Perrin and R. Remuson, *Tetrahedron: Asymmetry*, 1998, **9**, 1823–1828.
- 29 D. L. Comins and D. H. LaMunyon, J. Org. Chem., 1992, 57, 5807–5809.
- 30 For other examples on promoting conjugate addition by TMS-Cl: (*a*) S. Hanessian and K. Sumi, *Synthesis*, 1991, 1083–1089; (*b*) M. P. Jennings and K. B. Sawant, *Eur. J. Org. Chem.*, 2004, 3201–3204.
- 31 The stereochemistry was established by literature precedents as in ref. 28. See also: J. Thiel, W. Wysocka and W. Boczon, *Monatsh. Chem.*, 1995, 126, 233–239.