

CrossMark
click for updatesCite this: *RSC Adv.*, 2015, 5, 22075

DOI: 10.1039/c5ra90018b

www.rsc.org/advances

Correction: From micron to nano-curcumin by sophorolipid co-processing: highly enhanced bioavailability, fluorescence, and anti-cancer efficacy

Pradeep Kumar Singh,^{ab} Kirtee Wani,^c Ruchika Kaul-Ghanekar,^c Asmita Prabhune^{*b} and Satishchandra Ogale^{*a}

Correction for 'From micron to nano-curcumin by sophorolipid co-processing: highly enhanced bioavailability, fluorescence, and anti-cancer efficacy' by Pradeep Kumar Singh *et al.*, *RSC Adv.*, 2014, 4, 60334–60341.

There are errors in the structures shown in Fig. 1 of the paper. The correct Fig. 1 is shown below.




Fig. 1 Structure of SL(A) + Cur self-assembly. SL(A) is seen to completely encapsulate curcumin through hydrophobic part because of its hydrophobic nature. The hydrophilic part of SL(A) makes this assembly soluble in aqueous environment and prevents it from degradation.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aPhysical and Materials Chemistry Division, National Chemical Laboratory (NCL), Council of Scientific and Industrial Research (CSIR), Dr Homi Bhabha Road, Pashan, Pune 411008, India. E-mail: sb.ogale@ncl.res.in

^bBiochemical Sciences Division, National Chemical Laboratory (NCL), Council of Scientific and Industrial Research (CSIR), Dr Homi Bhabha Road, Pashan, Pune 411008, India. E-mail: aa.prabhune@ncl.res.in

^cCell and Translational Research Lab, Interactive Research School for Health Affairs (IRSHA), Bharati Vidyapeeth Deemed University, Pune, India