

Cite this: *Chem. Sci.*, 2015, **6**, 3044

Received 20th January 2015

Accepted 13th March 2015

DOI: 10.1039/c5sc00221d

[www.rsc.org/chemicalscience](http://www.rsc.org/chemicalscience)

## Iron(II)-catalyzed asymmetric intramolecular olefin aminochlorination using chloride ion<sup>†</sup>

Cheng-Liang Zhu,<sup>‡,a</sup> Jun-Shan Tian,<sup>‡,a</sup> Zhen-Yuan Gu,<sup>ab</sup> Guo-Wen Xing<sup>b</sup> and Hao Xu<sup>\*a</sup>

An iron-catalyzed enantioselective and diastereoselective intramolecular olefin aminochlorination reaction is reported (ee up to 92%, dr up to 15 : 1). In this reaction, a functionalized hydroxylamine and chloride ion are utilized as nitrogen and chlorine sources, respectively. This new method tolerates a range of synthetically valuable internal olefins that are all incompatible with existing asymmetric olefin aminochlorination methods.

### Introduction


Enantioselective olefin halo-functionalization reactions constitute a range of synthetically valuable yet challenging transformations.<sup>1</sup> Although a variety of excellent asymmetric olefin halo-oxygenation reactions have been discovered,<sup>2</sup> there are much fewer asymmetric olefin aminohalogenation methods available.<sup>3</sup> In particular, there have been just a few reported catalytic asymmetric olefin aminochlorination reactions.<sup>4</sup> In one instance, Feng discovered the chiral Lewis acid-catalyzed aminochlorination of chalconic and other  $\alpha,\beta$ -unsaturated olefins.<sup>4a,c</sup> Also, Chemler reported copper-catalyzed aminochlorination of terminal olefins with chlorine radical donors in the presence of  $MnO_2$  (Scheme 1A).<sup>4b</sup> Despite these and other important discoveries, catalytic asymmetric aminochlorination methods for internal, non-chalconic olefins have yet to be developed. These methods would be synthetically valuable because they would readily provide vicinal amino chlorides, a class of important chiral building blocks. Moreover, asymmetric olefin aminochlorination that proceeds through an iron-nitrenoid intermediate has not yet been reported.<sup>5</sup>

We previously discovered  $Fe(BF_4)_2$ -based catalysts for both diastereoselective and enantioselective intramolecular olefin aminofluorination reactions.<sup>6</sup> Our initial attempts to apply these catalysts to olefin aminochlorination reactions led to either low diastereoselectivity or low yield, presumably due to the reason that chlorine and fluorine atom-transfer may proceed through


distinct mechanisms. Therefore, we explored a range of activating group-ligand combinations and discovered entirely new catalytic conditions for asymmetric olefin aminochlorination. Herein, we describe iron-catalyzed enantioselective and diastereoselective intramolecular aminochlorination for a range of internal, non-chalconic olefins (ee up to 92%, dr up to 15 : 1). In these reactions, a functionalized hydroxylamine and chloride ion were utilized as nitrogen and chlorine sources, respectively. This method tolerates a range of synthetically valuable internal olefins that are all incompatible with existing asymmetric olefin aminochlorination approaches; it also provides a new approach that is complementary to known methods for the asymmetric synthesis of amino chlorides with contiguous stereogenic centers.

Prior to this research, Bach reported an  $FeCl_2$ -catalyzed racemic intramolecular olefin aminochlorination method using acyl azides,  $TMSCl$ , and  $EtOH$  under ligand-free conditions.<sup>7</sup>

A) previous work: asymmetric aminochlorination of chalconic and terminal olefins



B) current work: iron-catalyzed asymmetric aminochlorination of internal olefins with chloride ion



Scheme 1 Catalytic asymmetric olefin aminochlorination: summary of this work and other existing asymmetric methods.

<sup>a</sup>Department of Chemistry, Georgia State University, 100 Piedmont Avenue SE, Atlanta, Georgia, 30303, USA. E-mail: [hxu@gsu.edu](mailto:hxu@gsu.edu); Fax: +1-404-413-5505; Tel: +1-404-413-5553


<sup>b</sup>Department of Chemistry, Beijing Normal University, Beijing, 100875, China

<sup>†</sup> Electronic supplementary information (ESI) available: Experimental procedure, characterization data for all new compounds, selected NMR spectra and HPLC traces. CCDC 1041826. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c5sc00221d

<sup>‡</sup> These authors contributed equally.



Table 1 Catalyst discovery for the iron-catalyzed diastereoselective olefin aminochlorination reaction




Entry <sup>a</sup>	Fe(X) <sub>2</sub>	Ligand (mol%)	Conversion <sup>b</sup>	Yield <sup>c</sup>	dr <sup>b</sup> (anti : syn)
1	FeCl <sub>2</sub>	None	62%	45%	2 : 1
2	FeCl <sub>2</sub>	L1 (20)	>95%	80%	>20 : 1
3	Fe(NTf <sub>2</sub> ) <sub>2</sub>	L1 (20)	>95%	86%	>20 : 1
4	Fe(NTf <sub>2</sub> ) <sub>2</sub>	L2 (10)	>95%	82%	0.83 : 1
5	Fe(NTf <sub>2</sub> ) <sub>2</sub>	L3 (10)	61%	34%	0.25 : 1
6	Fe(NTf <sub>2</sub> ) <sub>2</sub>	L4 (20)	>95%	75%	1.8 : 1

<sup>a</sup> Unless stated otherwise, the reactions were carried out under a nitrogen atmosphere. TBAC: tetra-*n*-butylammonium chloride. <sup>b</sup> Conversion and dr were determined by <sup>1</sup>H NMR. <sup>c</sup> Isolated yield.

Excellent syn-selectivity was observed with styrenyl olefins (dr up to > 20 : 1). However, poor diastereoselectivity was recorded with non-styrenyl acyclic olefins (dr: 1 : 1). The new method presented here has a few unique features which complement the existing iron-catalyzed olefin aminochlorination method. First, excellent anti-selectivity has been observed across a wide range of styrenyl and non-styrenyl olefins. Second, good to excellent enantioselectivity has been achieved with a variety of internal, non-chalconic olefins (ee up to 92%). Finally, acyl azides are non-reactive under the described reaction conditions (*vide infra*), which suggests that iron-nitrenoid generation may proceed *via* different pathways compared with the known azide activation pathway.

## Results and discussion

A cinnamyl alcohol-derived acyloxyl carbamate **1** was selected as the model substrate for catalyst discovery (Table 1).<sup>8</sup> In the presence of tetra-*n*-butylammonium chloride (TBAC), we observed that FeCl<sub>2</sub> alone catalyzed a sluggish reaction under ligand-free conditions (entry 1, 45% yield, dr: 2 : 1).<sup>9</sup> However, the FeCl<sub>2</sub>-phenanthroline (**L1**) complex catalyzed the anti-



**Scheme 2** Iron-catalyzed aminochlorination with a *cis*-olefin and an acyl azide. <sup>a</sup>Reaction conditions: Fe(NTf<sub>2</sub>)<sub>2</sub> (10 mol%), **L1** (20 mol%), TBAC (2.5 equiv.), CH<sub>2</sub>Cl<sub>2</sub>, 0 °C, 2 h. <sup>b</sup>Reaction conditions: Fe(NTf<sub>2</sub>)<sub>2</sub> (10 mol%), **L4** (20 mol%), TBAC (2.5 equiv.), CH<sub>2</sub>Cl<sub>2</sub>, 0 °C, 2 h.

Table 2 Substrate scope of the iron-catalyzed diastereoselective olefin aminochlorination reaction

olefinic substrates	Fe(NTf <sub>2</sub> ) <sub>2</sub> (10 mol %) <b>L1</b> (20 mol %) TBAC (2.5 equiv) CH <sub>2</sub> Cl <sub>2</sub> , 0 °C, 2 h	olefin aminochlorination products			
1	86% yield, dr >20:1 from <i>E</i> olefin 83% yield, dr: 0.46:1 from <i>Z</i> olefin	2	86% yield, dr >20:1 <sup>a</sup>	3	70% yield, dr: 7:1
4	67% yield, dr: 10:1	5	76% yield, dr: 10:1	6	76% yield, dr: 12:1
7	61% yield, dr >20:1 <sup>a</sup>	8	59%, dr >20:1	9	93% yield, dr: 4.7:1 from <i>E</i> olefin 84% yield, dr: 7:1 from <i>Z</i> olefin <sup>a</sup>
10	50% yield, dr >20:1	11	76% yield	12	69% yield, dr >20:1 <sup>b</sup>
13	77% yield <sup>c</sup>	14	88% yield, dr: 1.7:1	15	64% yield, dr >20:1

<sup>a</sup> Reaction conditions: -15 °C, 2 h. <sup>b</sup> Reaction conditions: 0 °C, 5 h.

<sup>c</sup> Reaction conditions: 0 °C, 12 h.



aminochlorination with significantly improved yield and dr (entry 2, 80% yield, dr > 20 : 1). We also noted that the  $\text{Fe}(\text{NTf}_2)_2\text{-L1}$  complex provided essentially the same reactivity and diastereoselectivity (entry 3, 86% yield, dr > 20 : 1). Interestingly, the  $\text{Fe}(\text{NTf}_2)_2$ -bisoxazoline (**L2**) complex resulted in a loss of diastereoselectivity (entry 4, 82% yield, dr: 0.83 : 1). Furthermore, the  $\text{Fe}(\text{NTf}_2)_2\text{-L3}$  complex promoted the syn-aminochlorination with moderate yield and dr (entry 5, 34% yield, dr: 0.25 : 1). We also observed that the  $\text{Fe}(\text{NTf}_2)_2\text{-L4}$  complex catalyzed the anti-aminochlorination with a modest dr (entry 6, 75% yield, dr: 1.8 : 1). Notably, an iron-**L4** complex resulted in high dr and reaction rate in the previously reported olefin aminofluorination reaction.<sup>6</sup> These observations suggest that ligands are involved in the diastereoselectivity-determining step and provide excellent opportunities for diastereo-control.


The observed ligand-enabled diastereo-control with *trans*-olefin **1** prompted us to evaluate *cis*-olefin **1'** (Scheme 2). To our surprise, the  $\text{Fe}(\text{NTf}_2)_2\text{-L1}$  complex catalyzed syn-aminochlorination, while the  $\text{Fe}(\text{NTf}_2)_2\text{-L4}$  complex promoted anti-aminochlorination with essentially the same dr (Scheme 2). The different reaction profiles for isomeric olefins **1** and **1'** suggest that the aminochlorination reaction is neither stereospecific nor fully stereo-convergent, which is significantly different from the iron-catalyzed olefin aminofluorination reaction.<sup>6</sup>

Furthermore, an acyl azide **3** was evaluated under the reaction conditions as a control experiment. Interestingly, the acyl azide **3** was fully recovered and no aminochlorination product

was detected. These results suggest that the activation of acyloxyl carbamates (**1** and **1'**) may proceed *via* different pathways compared with the known azide activation pathway.<sup>7</sup>

We subsequently explored a range of olefins under the optimized conditions to evaluate the scope and limitations of this anti-aminochlorination method (Table 2). We discovered that di-substituted styrenyl olefins are generally good substrates; both electron-donating and electron-withdrawing substituents are compatible with this method (entries 1–4). Importantly, *ortho*-substituents and pyridyl groups are both tolerated (entries 5–6). Furthermore, extended aromatics, including naphthyl olefins, are reasonable substrates (entries 7–8). Moreover, isomeric ene-ynes are both excellent substrates for the stereo-convergent and anti-selective method (entry 9). Additionally, we observed that both styrenyl and non-styrenyl tri-substituted olefins undergo aminochlorination smoothly with excellent dr (entries 10–11).<sup>10</sup> We also discovered that a cyclohexyl-substituted olefin was an excellent substrate (entry 12, dr > 20 : 1). Further exploration revealed that both 1,1-disubstituted olefins and dienes are viable substrates with excellent regioselectivity (entries 13–14). Most notably, a cyclic olefin could also undergo highly diastereoselective anti-aminochlorination (entry 15, dr > 20 : 1), yielding a product which is difficult to obtain with known methods.<sup>11</sup> Since the  $\text{FeCl}_2\text{-L1}$  complex provides essentially the same dr and yield in these diastereoselective reactions,  $\text{FeCl}_2$  can be a convenient substitute for  $\text{Fe}(\text{NTf}_2)_2$  in racemic reactions.

Table 3 Catalyst discovery for the iron-catalyzed asymmetric olefin aminochlorination reaction



Entry <sup>a</sup>	R	Ligand	Conversion <sup>c</sup>	Yield <sup>d</sup>	dr <sup>c</sup> (anti : syn)	ee <sup>e</sup> (anti)	ee <sup>e</sup> (syn)
1	$3,5-(\text{CF}_3)_2\text{-Ph}$	<b>L5</b>	>95%	53%	9.9 : 1	84%	<5%
2	$3,5-(\text{CF}_3)_2\text{-Ph}$	<b>L6</b>	>95%	68%	0.5 : 1	24%	79%
3	$3,5-(\text{CF}_3)_2\text{-Ph}$	<b>L7</b>	88%	61%	1.7 : 1	<5%	<5%
4	$3,5-(\text{CF}_3)_2\text{-Ph}$	<b>L8</b>	>95%	32%	2.5 : 1	47%	30%
5	$3,5-(\text{CF}_3)_2\text{-Ph}$	<b>L9</b>	>95%	82%	0.5 : 1	8%	24%
6 <sup>f</sup>	$3,5-(\text{CF}_3)_2\text{-Ph}$	<b>L5</b>	>95%	51%	11.0 : 1	90%	<5%
7 <sup>f</sup>	$\text{CH}_3$	<b>L5</b>	>95%	42%	1.1 : 1	97%	<5%
8 <sup>f</sup>	$\text{CH}_2\text{Cl}$	<b>L5</b>	>95%	67%	9.6 : 1	89%	<5%
9 <sup>f,g</sup>	$\text{CH}_2\text{Cl}$	<b>L5</b>	>95%	58%	9.0 : 1	83%	<5%

<sup>a</sup> Unless stated otherwise, the reactions were carried out under a nitrogen atmosphere with 4 Å molecular sieves. <sup>b</sup> Reaction conditions:  $\text{Boc}_2\text{O}$ ,  $\text{Et}_3\text{N}$ , DMAP; then  $\text{Cs}_2\text{CO}_3$ , MeOH, 85% over two steps; see ESI for details. <sup>c</sup> Conversion and dr were determined by  $^1\text{H}$  NMR. <sup>d</sup> Isolated yield. <sup>e</sup> Enantiomeric excess (ee) was measured by HPLC with chiral columns; the absolute stereochemistry was determined by X-ray crystallographic analysis of an analog of **2a**. <sup>f</sup> The reaction was carried out at  $-60\text{ }^\circ\text{C}$  for 12 h. <sup>g</sup> The  $\text{FeCl}_2\text{-L5}$  complex was used.



In order to fulfil the need for catalytic asymmetric olefin aminochlorination, we further explored asymmetric induction for internal, non-chalconic olefins with a variety of iron-chiral ligand complexes (Table 3).<sup>12</sup> First, we discovered that the iron-L5 complex induced diastereoselective and enantioselective anti-aminochlorination, albeit with a low yield, mostly due to the competing aminohydroxylation reaction (entry 1, 53% yield, dr: 9.9 : 1). Interestingly, the anti-addition product **2a** was obtained with excellent ee (84% ee), while the syn-addition product **2b** was obtained essentially as a racemate (<5% ee).<sup>13</sup> Additionally, a two-step procedure can convert **2a** to a chlorinated amino alcohol triad **4** without ee erosion.<sup>14</sup> Next, we observed that the iron-L6 complex induced moderately diastereoselective syn-aminochlorination (entry 2, 68% yield, dr: 0.48 : 1). To our surprise, the anti-addition product **2a** was obtained with moderate ee (24% ee), while the syn-addition product **2b** was isolated with significant ee (79% ee). Furthermore, we evaluated chiral ligands L7 and L8 and determined that they are less effective for asymmetric induction (entries 3–4). Additionally, chiral ligand L9 induced fast yet non-selective aminochlorination with a high overall yield (entry 5).<sup>15</sup> With the iron-L5 complex in hand, we subsequently explored other reaction parameters. First, a decreased reaction temperature was found to benefit both dr and ee (entry 6, dr: 11 : 1 and 90% ee for **2a** at  $-60^{\circ}\text{C}$ ). Next, replacing the 3,5-bis(trifluoromethyl)benzoyl activating group with a smaller acetyl group further enhanced the ee (entry 7, 97% ee for **2a**); however, much lower dr and yield were obtained (entry 7, dr: 1.1 : 1, 42% yield). Finally, a chloroacetyl activating group induced an effective balance between overall yield and stereoselectivity (entry 8, 67% yield, dr: 9.6 : 1 and 89% ee for **2a**). We also observed that the  $\text{FeCl}_2$ -L5 complex induced a slightly less selective reaction with a lower yield (entry 9, 58% yield, dr: 9.0 : 1 and 83% ee for **2a**).


In order to evaluate the scope of this asymmetric method, we explored the asymmetric induction with a range of internal olefins (Table 4). The chiral catalyst provides excellent asymmetric induction with styrenyl olefins. A range of *para*-substituted styrenyl olefins with different electronic properties were converted to the corresponding aminochlorination products with high dr and ee (entries 1–6, dr: 9.6–15 : 1, ee: 86–91%). Additionally, *meta*-substituted styrenyl olefins are also good substrates but with slightly decreased ee (entries 7–9, dr: 10–15 : 1, ee: 80–87%). However, we discovered that *ortho*-substitution of styrenes has a deleterious effect on ee (entries 10–11, dr: 4.5–12 : 1, ee: 77–79%). Interestingly, both  $\alpha$ - and  $\beta$ -naphthyl olefins are excellent substrates (entries 12–13, dr: 4.5–10 : 1, ee: 89–92%). To our delight, a 3-pyridyl olefin with a basic nitrogen atom is a reasonable substrate for the asymmetric aminochlorination (entry 14, dr: 1.8 : 1, ee: 70% for the anti-diastereomer). Moreover, we observed that the iron-L5 complex can induce significant ee in the aminochlorination with non-styrenyl olefins (entry 15, dr: 2 : 1, ee: 54% for the anti-diastereomer). To our surprise, the iron-L6 complex proved to be uniquely effective for the asymmetric induction with tri-substituted olefins, while the iron-L5 complex was less effective (entry 16, dr: 2.3 : 1, ee: 86% for the anti-diastereomer).<sup>16</sup>

Table 4 Substrate scope for the iron-catalyzed asymmetric olefin aminochlorination reaction

olefinic substrates <sup>a</sup>	$\text{Fe}(\text{NTf}_2)_2$ (15 mol %) L5 (15 mol %) TBAC (2.5 equiv)	CHCl <sub>3</sub> , $-60^{\circ}\text{C}$ , 12 h	olefin aminochlorination products <sup>a</sup>
1			
2			
3			
4		67% yield, dr: 9.6:1 89% ee	
5		65% yield, dr: 15:1 91% ee	
6		69% yield, dr: 5.2:1 87% ee	
7		84% yield, dr: 12:1 90% ee	
8		62% yield, dr: 11:1 88% ee	
9		71% yield, dr: 11:1 86% ee	
10		75% yield, dr: 12:1 87% ee	
11		63% yield, dr: 10:1 80% ee	
12		71% yield, dr: 15:1 80% ee	
13		78% yield, dr: 4.5:1 77% ee <sup>b</sup>	
14		55% yield, dr: 12:1 79% ee	
15		63% yield, dr: 10:1 92% ee <sup>b</sup>	
16		66% yield, dr: 2:1 45% yield, dr: 2.3:1 54% ee <sup>b,c</sup> 86% ee <sup>b,d</sup>	
		53% yield, dr: 4.5:1 89% ee <sup>b</sup>	
		51% yield, dr: 1.8:1 70% ee	

<sup>a</sup> Unless stated otherwise, mono-chloroacetyl was selected as the activating group for asymmetric catalysis; the ee for all syn-aminochlorination products was less than 5%. <sup>b</sup> Bis(trifluoromethyl)-benzoyl was selected as the activating group. <sup>c</sup> The ee for the syn-addition product was 12%. <sup>d</sup> L6 was used as the ligand for asymmetric induction; the ee for the syn-addition product was 50%.

During the exploration of substrate scope, it was surprising to observe completely different ee values for anti- and syn-diastereomers (e.g. **2a** and **2b**). In contrast, exactly the same ee for both diastereomeric products was observed in the iron-catalyzed aminofluorination of **1**.<sup>6</sup> In order to obtain greater mechanistic insights, we carried out ee analysis for all isolable products using several control experiments (Scheme 3). First, in an  $\text{Fe}(\text{NTf}_2)_2$ -catalyzed reaction with *trans*-olefin **1**, two amino-chlorination products were obtained (Scheme 3A, 90% ee for **2a**, <5% ee for **2b**, dr: 11 : 1).<sup>17</sup> Simultaneously, diastereomers **5a** and **5b** were also isolated with the same ee as two competing olefin aminohydroxylation products (Scheme 3A, 88% ee for **5a** and **5b**, dr: 4 : 1). However, completely different selectivity (both dr and ee) was observed in an  $\text{Fe}(\text{NTf}_2)_2$ -catalyzed reaction with *cis*-olefin **1'** (Scheme 3A, 85% ee for **2a** and 31% ee for **2b**, dr:



Scheme 3 Control experiments to probe the mechanism. <sup>a</sup>Reaction conditions:  $\text{Fe}(\text{NTf}_2)_2$  (15 mol%), L1 (15 mol%), TBAC (2.5 equiv.),  $\text{CHCl}_3$ ,  $-60^\circ\text{C}$ , 12 h. <sup>b</sup>Reaction conditions:  $\text{Fe}(\text{NTf}_2)_2$  (15 mol%), L1 (15 mol%),  $\text{CHCl}_3$ ,  $-60^\circ\text{C}$ , 12 h.


6 : 1; 93% ee for 5a and 83% ee for 5b, dr: 7 : 1). In both cases, 5a and 5b cannot be converted to 2a under the reaction conditions.

These observations provide several important mechanistic insights. First, the non-stereospecificity observed in the iron-catalyzed olefin aminochlorination suggests that the formation of C–N and C–Cl bonds occurs in a stepwise fashion.<sup>18</sup> Second, the lack of complete stereo-convergence between the reaction profiles of isomeric olefins (1 and 1') suggests that C–N bond formation may be the rate- and ee-determining step.<sup>18</sup> Furthermore, since essentially the same ee was observed for 2a, 5a, and 5b from the reaction with *trans*-olefin 1, it is likely that these products are derived from the same intermediate after the ee-determining step. Additionally, the fact that the syn-aminochlorination product 2b was isolated as a racemate suggests that 2b may be derived from non-stereoselective pathways

which are distinct from the one leading to the formation of 2a, 5a, and 5b.

The product divergence (2a vs. 5a/b) after the ee-determining step is mechanistically interesting. Therefore, we studied the effect of external chloride ion. To our surprise, in the absence of TBAC, the  $\text{Fe}(\text{NTf}_2)_2$ –L5 complex alone was ineffective for the nitrogen atom-transfer at  $-60^\circ\text{C}$ ; 1 and 1' were both fully recovered (Scheme 3B). However, aminochlorination occurred as soon as a stoichiometric amount of TBAC was introduced. This observation suggests that the  $\text{Fe}(\text{NTf}_2)_2$ –L5 complex may serve as a pre-catalyst and it may be activated by chloride ion *in situ*.

In order to test this hypothesis, we further carried out the  $\text{FeCl}_2$ -catalyzed reaction in the presence of TBAC (Scheme 3C). Notably, 2a was isolated with essentially the same ee as that obtained under the standard conditions (88% ee for 2a and <5%



Scheme 4 Proposed mechanistic working hypothesis for the iron-catalyzed asymmetric aminochlorination of *trans*-olefin 1.

ee for **2b**). This result suggests that the catalytically relevant species may also be generated from the  $\text{FeCl}_2\text{-L5}$  complex.

To probe for more mechanistic details, we subsequently carried out the  $\text{FeCl}_2$ -promoted olefin aminochlorination in the absence of TBAC (100 mol%  $\text{FeCl}_2$ , 100 mol% **L5**, Scheme 3C). Under these conditions,  $\text{FeCl}_2$  is the only available chlorine source. Surprisingly, we discovered that **2a** was obtained with essentially the same ee compared with the two previous control experiments (88% ee for **2a**). Furthermore, a syn-amino-hydroxylation product **5a** was isolated with excellent dr and ee (dr > 20 : 1, 88% ee). These observations suggest that Fe–Cl bond cleavage may be relevant for the chlorine atom-transfer step during the enantioselective anti-aminochlorination.<sup>19</sup> In addition, we also identified a small amount of aziridine **6** (15% yield, 82% ee), and further discovered that it could not be converted to either **2a** or **5a** under the reaction conditions.

With the accumulated mechanistic evidence, we propose a plausible mechanistic working hypothesis for the iron-catalyzed asymmetric aminochlorination of *trans*-olefin **1** (Scheme 4). First, the iron catalyst reversibly cleaves the N–O bond in the acyloxy carbamate **1**, generating iron-nitrenoid **A** with chloride as a counter ion. From there, **A** may participate in enantioselective and diastereoselective aminochlorination and amino-hydroxylation to afford **2a** and **5a**, respectively. Since the aminochlorination–aminohydroxylation competition occurs after the ee-determining step, **2a** is obtained with essentially the same ee as **5a**. At the same time, **1** may be converted to **2b** via a non-stereoselective pathway which is distinct from the one leading to the formation of **2a** and **5a**. Further mechanistic studies are required to elucidate the details.

## Conclusions

In conclusion, we have described an iron-catalyzed enantioselective and diastereoselective aminochlorination method for internal, non-chalconic olefins. This method tolerates a range of synthetically valuable olefins that are all incompatible with existing asymmetric olefin aminochlorination methods. It also provides a complementary approach for the asymmetric synthesis of amino chlorides with contiguous stereogenic centers. Our preliminary mechanistic studies revealed that an  $\text{FeCl}_2$ -derived nitrenoid may be a feasible reactive intermediate and that Fe–Cl bond cleavage may be relevant for stereoselective chlorine atom-transfer. Our current efforts are focused on the mechanistic investigation of this new reaction and method development for the enantioselective intermolecular olefin aminochlorination.

## Acknowledgements

This work was supported by the National Institutes of Health (GM110382) and Georgia State University. Z.-Y. G. was supported by NSFC (21272027) and a fellowship from China Scholarship Council.

## Notes and references

- For selected reviews of asymmetric olefin halofunctionalization, see: (a) S. E. Denmark, W. E. Kuester and M. T. Burk, *Angew. Chem., Int. Ed.*, 2012, **51**, 10938; (b) S. R. Chemler and M. T. Bovino, *ACS Catal.*, 2013, **3**, 1076; (c) S. A. Snyder, D. S. Treitler and A. P. Brucks, *Aldrichimica Acta*, 2011, **44**, 27.
- For selected reports on catalytic asymmetric olefin halo-oxidation, see: (a) S. H. Kang, S. B. Lee and C. M. Park, *J. Am. Chem. Soc.*, 2003, **125**, 15748; (b) G. E. Veitch and E. N. Jacobsen, *Angew. Chem., Int. Ed.*, 2010, **49**, 7332; (c) W. Zhang, S. Zheng, N. Liu, J. B. Werness, I. A. Guzei and W. Tang, *J. Am. Chem. Soc.*, 2010, **132**, 3664; (d) L. Zhou, C. K. Tan, X. Jiang, F. Chen and Y.-Y. Yeung, *J. Am. Chem. Soc.*, 2010, **132**, 15474; (e) K. Murai, T. Matsushita, A. Nakamura, S. Fukushima, M. Shimura and H. Fujioka, *Angew. Chem., Int. Ed.*, 2010, **49**, 9174; (f) S. E. Denmark and M. T. Burk, *Org. Lett.*, 2012, **14**, 256; (g) D. Huang, H. Wang, F. Xue, H. Guan, L. Li, X. Peng and Y. Shi, *Org. Lett.*, 2011, **13**, 6350; (h) R. Yousefi, D. C. Whitehead, J. M. Mueller, R. J. Staples and B. Borhan, *Org. Lett.*, 2011, **13**, 608; (i) R. Yousefi, K. D. Ashtekar, D. C. Whitehead, J. E. Jackson and B. Borhan, *J. Am. Chem. Soc.*, 2013, **135**, 14524; (j) D. H. Paull, C. Fang, J. R. Donald, A. D. Pansick and S. F. Martin, *J. Am. Chem. Soc.*, 2012, **134**, 11128; (k) M. C. Dobish and J. N. Johnston, *J. Am. Chem. Soc.*, 2012, **134**, 6068; (l) Y.-M. Wang, J. Wu, C. Hoong, V. Rauniar and F. D. Toste, *J. Am. Chem. Soc.*, 2012, **134**, 12928; (m) V. Rauniar, A. D. Lackner, G. L. Hamilton and F. D. Toste, *Science*, 2011, **334**, 1681; (n) T. Honjo, R. J. Phipps, V. Rauniar and F. D. Toste, *Angew. Chem., Int. Ed.*, 2012, **51**, 9684; (o) J. Wu, Y.-M. Wang, A. Driljevic, V. Rauniar, R. J. Phipps and F. D. Toste, *Proc. Natl. Acad. Sci. U. S. A.*, 2013, **110**, 13729; (p) H. Nakatsuji, Y. Sawamura, A. Sakakura and K. Ishihara, *Angew. Chem., Int. Ed.*, 2014, **53**, 6974; (q) L. Li, C. Su, X. Liu, H. Tian and Y. Shi, *Org. Lett.*, 2014, **16**, 3728.
- For selected reports on catalytic asymmetric olefin aminobromination and aminoiodination, see: (a) Y. Cai, X. Liu, Y. Hui, J. Jiang, W. Wang, W. Chen, L. Lin and X. Feng, *Angew. Chem., Int. Ed.*, 2010, **49**, 6160; (b) L. Zhou, J. Chen, C. K. Tan and Y.-Y. Yeung, *J. Am. Chem. Soc.*, 2011, **133**, 9164; (c) Y. F. Cai, X. H. Liu, J. Li, W. L. Chen, W. T. Wang, L. L. Lin and X. M. Feng, *Chem.-Eur. J.*, 2011, **17**, 14916; (d) A. Alix, C. Lalli, P. Retailleau and G. Masson, *J. Am. Chem. Soc.*, 2012, **134**, 10389; (e) D. Huang, X. Liu, L. Li, Y. Cai, W. Liu and Y. Shi, *J. Am. Chem. Soc.*, 2013, **135**, 8101; (f) C. S. Brindle, C. S. Yeung and E. N. Jacobsen, *Chem. Sci.*, 2013, **4**, 2100; (g) F. Chen, C. K. Tan and Y.-Y. Yeung, *J. Am. Chem. Soc.*, 2013, **135**, 1232. For mechanistically related asymmetric olefin sulfenofunctionalization, see: (h) S. E. Denmark and H. M. Chi, *J. Am. Chem. Soc.*, 2014, **136**, 8915; (i) S. E. Denmark, E. Hartmann, D. J. P. Kornfilt and H. Wang, *Nat. Chem.*, 2014, **6**, 1056.



4 For existing asymmetric olefin aminochlorination methods, see: (a) Y. F. Cai, X. H. Liu, J. Jiang, W. L. Chen, L. L. Lin and X. M. Feng, *J. Am. Chem. Soc.*, 2011, **133**, 5636; (b) M. T. Bovino and S. R. Chemler, *Angew. Chem., Int. Ed.*, 2012, **51**, 3923; (c) Y. Cai, X. Liu, P. Zhou, Y. Kuang, L. Lin and X. M. Feng, *Chem. Commun.*, 2013, **49**, 8054.

5 For catalytic olefin aminoxylation that proceeds through an iron-nitrenoid intermediate, see: (a) G.-S. Liu, Y.-Q. Zhang, Y.-A. Yuan and H. Xu, *J. Am. Chem. Soc.*, 2013, **135**, 3343; (b) Y.-Q. Zhang, Y.-A. Yuan, G.-S. Liu and H. Xu, *Org. Lett.*, 2013, **15**, 3910; (c) D.-F. Lu, C.-L. Zhu, Z.-X. Jia and H. Xu, *J. Am. Chem. Soc.*, 2014, **136**, 13186.

6 D.-F. Lu, G.-S. Liu, C.-L. Zhu, B. Yuan and H. Xu, *Org. Lett.*, 2014, **16**, 2912.

7 (a) T. Bach, B. Schlummer and K. Harms, *Chem. Commun.*, 2000, 287; (b) T. Bach, B. Schlummer and K. Harms, *Chem.-Eur. J.*, 2001, **7**, 2581; (c) H. Danielec, J. Klügge, B. Schlummer and T. Bach, *Synthesis*, 2006, 551.

8 See ESI† for details of substrate synthesis. Acyloxy carbamates are reactive, while tosyloxyl and alkoxy carbamates are non-reactive and fully recovered under the reaction conditions.

9 The relative stereochemistry of **2a** was determined by comparison of the experimental NMR data with those reported in ref. 7. It was further corroborated by <sup>1</sup>H NMR and X-ray crystallographic analysis of a structural analog of **2a**. See ESI† for details.

10 The relative stereochemistry was assigned based on the <sup>1</sup>H NMR and X-ray crystallographic analysis of a structural analog described in ref. 6; see ESI† for details.

11 Complementary stereochemistry was achieved (in entry 15 of Table 2) compared with the known method reported in ref. 7, where the syn-aminochlorination product was isolated. This substrate did not undergo kinetic resolution with a chiral catalyst, the complex Fe(NTf<sub>2</sub>)<sub>2</sub>-**L5**. Both the starting material and product were isolated as racemates.

12 For leading references on chiral BOX and related ligands, see: (a) D. A. Evans, K. A. Woerpel, M. M. Hinman and M. M. Faul, *J. Am. Chem. Soc.*, 1991, **113**, 726; (b) H. Nishiyama, Y. Itoh, H. Matsumoto, S.-B. Park and K. Itoh, *J. Am. Chem. Soc.*, 1994, **116**, 2223; (c) Y. Nishikawa and H. Yamamoto, *J. Am. Chem. Soc.*, 2011, **133**, 8432.

13 The absolute stereochemistry of **2a** was determined by X-ray crystallographic analysis of a structural analog of **2a**. See ESI† for details.

14 For detailed procedure and HPLC traces of **4**, see ESI†.

15 For the synthesis of **L9**, see ref. 6.

16 The iron-**L5** complex catalyzed the reaction favoring the syn-addition product (dr (anti/syn): 0.47 : 1); ee for the anti-addition product was 60% and ee for the syn-addition product was <5%. The relative stereochemistry was assigned based on the <sup>1</sup>H NMR and X-ray crystallographic analysis of a structural analog described in ref. 6; see ESI† for details.

17 When a chloroacetyl group was used as the activating group, a different result was obtained. For details, see entry 8 of Table 3.

18 For an example of stepwise atom transfer reactions with different reaction profiles for *cis/trans* isomeric olefins, see: N. H. Lee and E. N. Jacobsen, *Tetrahedron Lett.*, 1991, **32**, 6533.

19 For the oxidation of a radical species by a high-valent metal through ligand transfer or electron transfer, see: (a) M. S. Kharasch and G. Sosnovsky, *J. Am. Chem. Soc.*, 1958, **80**, 756; (b) J. K. Kochi, *Science*, 1967, **155**, 415. For a relevant enzymatic C–H chlorination reaction of hydrocarbons catalyzed by iron-containing metalloenzymes, see: (c) F. H. Vaillancourt, J. Yin and C. T. Walsh, *Proc. Natl. Acad. Sci. U. S. A.*, 2005, **102**, 10111.