

Cite this: *Chem. Sci.*, 2015, **6**, 4479Received 29th April 2015
Accepted 29th May 2015DOI: 10.1039/c5sc01553g
www.rsc.org/chemicalscience

Introduction

Desymmetrization has emerged as a way to access chiral quaternary-carbon motifs, which are among the most challenging stereocenters to generate with enantiocontrol.^{1–4} Strategies involving C–H bond activation are especially promising yet rare.^{5,6} Given this challenge, we propose that prochiral aldehyde **1** could isomerize to scaffolds bearing quaternary centers *via* two possible pathways triggered by aldehyde C–H bond activation (Fig. 1). Herein, we communicate Rh-catalyzed olefin functionalizations, including hydroacylation and carboacylation from a common aldehyde. This initial report focuses on hydroacylation of bis(allyl)aldehydes to generate α -vinylcyclopentanones **2** bearing quaternary stereocenters.⁷ Mechanistic studies reveal a cascade process featuring an enantioselective olefin-isomerization followed by olefin-hydroacylation.

The use of oxygen, nitrogen, and sulfur-based functional groups has allowed breakthroughs in enantioselective Rh-catalyzed hydroacylation.⁸ These heteroatoms act as directing groups by binding to rhodium and favoring C–H bond activation while accelerating hydroacylation over competitive pathways, such as decarbonylation or catalyst decomposition.⁹ Fu demonstrated intramolecular hydroacylation of alkynals bearing β -methoxy groups (Fig. 2a).¹⁰ Our laboratory reported intermolecular hydroacylation of cyclopropenes using chelating aldehydes, specifically salicylaldehyde derivatives (Fig. 2b).⁵ Given their ability to bind Rh, we reasoned that olefins could be used as directing groups for hydroacylation.¹¹ We were encouraged that Tanaka and Suemune reported

desymmetrization of β -bis(alkenyl) aldehydes (Fig. 2c).¹² Although not proposed, we reason that the pendant olefin in their substrate could be acting as a directing group. These previous desymmetrizations by hydroacylation generate ketones bearing β -quaternary stereocenters. Given this limitation, we chose to develop a complementary desymmetrization of α -trisubstituted aldehydes, which represents a sterically hindered and thus, challenging substrate class.^{13b–c,14} If successful, our strategy would allow access to cyclopentanones bearing α -quaternary centers, whereby the pendant olefin serves as both a directing group and versatile handle for further elaboration.




Fig. 1 Two pathways to quaternary-carbon motifs from desymmetrization of **1**.




Fig. 2 Previous desymmetrizations by hydroacylation result in ketones bearing β -quaternary stereocenters.

Department of Chemistry, University of California-Irvine, 4403 Natural Sciences 1, Irvine, California 92697, USA. E-mail: dongv@uci.edu

† Electronic supplementary information (ESI) available: Materials and methods, reaction procedures, characterization data. CCDC 1056687, 1062118. For ESI and crystallographic data in CIF or other electronic format see DOI: [10.1039/c5sc01553g](https://doi.org/10.1039/c5sc01553g)



Results and discussion

To test our proposal, we studied the desymmetrization of model **1a**, which can be prepared in one-step from commercially available phenylacetaldehyde.¹⁵ Aldehyde **1a** was subjected to cationic Rh(i)-catalysts and various bidentate phosphine ligands that are known to promote formyl C–H bond activation.¹⁶ We imagined that oxidative addition followed by alkene insertion would generate metallacycle **5**, which could diverge into various scaffolds (Table 1). The choice of phosphine ligand had a dramatic impact on product outcome and enabled chemoselective formation of two major products, cyclopentanone **2a** and bicyclo[2.2.1]heptanone **3a**.

With a BINAP-ligated rhodium catalyst, we observed formation of both **2a** and **3a** in 61% and 19% yields, respectively (entry 1). We discovered that the hydroacylation product, cyclopentanone **2a**, bears an internal olefin, which presumably results from isomerization of the terminal olefin. Carbometallation of the pendant olefin from **5** results in intermediate **7**,

which undergoes reductive elimination to form bicycloheptanone **3a** as a minor product. Use of BzDPPB ligand, however, favors the carboacetylation pathway to generate **3a** as the major product in 56% yield with high diastereoselectivity (>20 : 1, entry 2). This unique olefin functionalization takes advantage of C–H activation rather than strained C–C activation to achieve carboacetylation.¹⁷ While our study was in progress, Aïssa reported a related carbocyclization using pyridyl directing groups.¹⁸ We also observed bicyclo[3.2.1]octadione **4a** as a minor product in 10% yield (entry 2). The molecular structure of this homologated ketone **4a** was confirmed by X-ray crystallography (see ESI†). We believe that the second carbonyl arises from a disproportionation process where a second equivalent of aldehyde **1a** undergoes decarbonylation to generate CO.

With these promising leads in hand, we plan to further study each pathway and develop enantioselective variants. Towards this goal, we realized that electron-donating aromatic groups on phosphines enhance selectivity for **2a**. Among the ligands evaluated, (*R*)-DTBM-MeOBIPHEP provided the best reactivity

Table 1 Divergent pathways for desymmetrization of **1a** based on ligand-choice^a

Entry	Ligand	Major product, yield ^c	Minor product(s), yield ^d
1	(<i>R</i>)-BINAP	(<i>S</i>)- 2a 61%, 31% ee	3a 19%, 33% ee
2	BzDPPB	(\pm)- 3a 56%	(\pm)- 4a , 10% ^{e,f} (\pm)- 2a , 7%
3 ^b	(<i>R</i>)-DTBM-MeOBIPHEP	(<i>S</i>)- 2a 91%, 97% ee	3a and 4a not observed

^a Reaction conditions: $[(\text{coe})_2\text{RhCl}]_2$ (5 mol%), ligand (10 mol%), AgBF_4 (10 mol%), DCE (0.2 M), 40 °C, 36 h. ^b Reaction conditions: $[(\text{coe})_2\text{RhCl}]_2$ (2.5 mol%), ligand (5 mol%), AgBF_4 (5 mol%), 40 °C, 4 h. ^c Isolated yield. ^d Determined by GC-FID. ^e >20 : 1 *dr*, determined by ^1H NMR. ^f One equivalent of **1a** was used as a CO donor. DTBM: 3,5-di(*tert*-butyl)-4-methoxyphenyl.



and enantioselectivity for **2a** (entry 3). The absolute configuration of cyclopentanone **2a** was determined by elaboration with 2,4-dinitrophenylhydrazine to hydrazone **9**, in which the molecular structure was established by X-ray crystallographic analysis (Fig. 3).

With this protocol, we prepared eleven cyclopentanones bearing various α -quaternary stereocenters (Table 2). Aldehydes with aromatic substituents (**1a**–**1g**) undergo desymmetrization in 83–91% yields and high enantioselectivities (95–99% *ee*). Ether, aryl halide, and acetal functional groups are well-tolerated. Heteroaromatic aldehyde (**1h**) as well as aldehydes bearing aliphatic substituents (**1i**–**1k**) rearrange to the corresponding cyclopentanones in excellent enantioselectivities albeit using increased catalyst loading at lower temperature.^{19,20}

To understand the mechanism, we performed a deuterium-labelling study with **d-1a**. Desymmetrization of **d-1a**, under standard reaction conditions, led to exclusive formation of **d-2a** where the deuterium label was incorporated into the methyl group of the α -propenyl substituent (eqn (1)). This result indicates that isomerization of one allyl group takes place first through an endocyclic β -hydride elimination of a 5-membered rhodacycle **d-5a**.²¹ Our observations corroborate Aïssa's recent report on the isomerisation of 4-pentenals.²² Although β -hydride eliminations of this type are uncommon, it has been predicted that binding of a pendant alkene to the metal center significantly lowers the barrier to this process.²³

When the reaction of **d-1a** was quenched at an early stage (40% conversion to **d-2a**), we recovered three deuterated aldehydes, **d-1a**, **d-1a'** and **d-1a''** (eqn (2)). The observation of **d-1a'** suggests that olefin-insertion is reversible. Yet, the deuterium is incorporated into only the methyl group of the α -propenyl unit in product **d-2a**. This lack of deuterium scrambling on the cyclopentanone ring suggests that Rh-D insertion occurs with high enantioselectivity (with the olefin shown in red). Thus, the insertion step is both reversible and highly enantioselective.²⁴

Further experiments support the notion that the α -vinyl group (formed from initial isomerization) directs hydroacylation. For example, α -trisubstituted aldehyde **10** (with only one allyl group) does not undergo hydroacylation. Instead, this aldehyde undergoes isomerization to generate α -vinyl aldehyde **11** (eqn (3)).²² In addition, subjecting α -allylcyclopentanone **12a** to the optimized reaction conditions results in trace formation of α -vinylcyclopentanone **2a** (eqn (4)). Thus, the cyclopentanones obtained in Table 2 must arise from an

isomerization that occurs prior to hydroacylation. In contrast, we discovered that aldehyde **1l**, containing an acetal group, yields α -allylcyclopentanone **12l** as the major product (eqn (5)). In this case, we reason that the acetal acts as an oxygen-directing group which promotes hydroacylation over olefin isomerization.







Table 2 Desymmetrization of α -quaternary aldehyde **1** by isomerization-hydroacylations^a



^a Reaction conditions: 0.1 mmol **1**, *x* = 2.5, DCE (0.2 M), 40 °C, 4 h.

^b Reaction conditions: 1 mmol **1e** used. ^c Reaction conditions: *x* = 5, DCE (0.33 M), 30 °C, 2 h. ^d Reaction conditions: *x* = 6, DCE (0.33 M), 30 °C, 2 h.






Fig. 3 Determination of the absolute configuration of **2a** and X-ray crystal structure of **(2S)-9**.





Scheme 1 Proposed mechanism for Rh-catalyzed cascade.



On the basis of literature reports and our own observations, we propose a mechanism starting with cationic Rh(I)-complex activating the aldehyde C-H bond of **1** to form acyl-Rh(III)-hydride **13** (Scheme 1). Insertion of the olefin into Rh(III)-hydride **13** leads to formation of the more thermodynamically stable 5-membered metallacycle **5**.²⁵ A rare endocyclic β-hydride elimination takes place to produce isomerized acyl-Rh(III)-hydride **14**. The allyl olefin inserts into the Rh(III)-hydride to form a 6-membered rhodacycle **6**. Finally, reductive elimination affords the cyclopentanone product **2** and regenerates the Rh(I)-catalyst.

Conclusions

We have demonstrated a Rh-catalyzed enantioselective synthesis of α-quaternary cyclopentanones. Studies on the scope and mechanism support an olefin-assisted isomerization²³ and olefin-directed hydroacylation cascade. While endocyclic β-hydride elimination has been proposed in the literature on the basis of theoretical studies,²¹ our results provide experimental evidence for this elementary step. The use of a BIPHEP ligand enables high selectivity for one out of three possible rearrangements, all initiated by the activation of an aldehyde C-H bond. Insights from these studies will guide efforts to understand and expand the power of the related carboacylation

and bisacylation as routes to scaffolds containing chiral all-carbon stereocenters.

Acknowledgements

Funding provided by UC Irvine and the National Institutes of Health (GM105938). We are grateful to Eli Lilly for a Grantee Award. K.G.M.K. thanks NSERC of Canada for a CGS-D fellowship. We thank Van Nguyen for initial ligand studies, Kaitlyn Kung for help with substrate synthesis, and Dr Joseph W. Ziller for X-ray crystallographic analysis.

Notes and references

- For reviews on enantioselective construction of quaternary stereocenters, including desymmetrization strategies, see: (a) K. W. Quasdorf and L. E. Overman, *Nature*, 2014, **516**, 181; (b) C. J. Douglas and L. E. Overman, *Proc. Natl. Acad. Sci. U. S. A.*, 2004, **101**, 5363; (c) Y. Liu, S.-J. Han, W.-B. Liu and B. M. Stoltz, *Acc. Chem. Res.*, 2015, **48**, 740; (d) H. Lebel, J.-F. Marcoux, C. Molinaro and A. B. Charette, *Chem. Rev.*, 2003, **103**, 977.
- For recent alkene functionalizations to establish enantioenriched quaternary stereocenters, see: T.-S. Mei, H. H. Patel and M. S. Sigman, *Nature*, 2014, **508**, 340.
- For recent selected desymmetrizations to quaternary stereocenters, see: (a) F. Zhou, G.-J. Cheng, W. Yang, Y. Long, S. Zhang, Y.-D. Wu, X. Zhang and Q. Cai, *Angew. Chem., Int. Ed.*, 2014, **53**, 9555; (b) C. Roux, M. Candy, J.-M. Pons, O. Chuzel and C. Bressy, *Angew. Chem., Int. Ed.*, 2014, **53**, 766; (c) L. Yao, K. Liu, H.-Y. Tao, G.-F. Qiu, X. Zhou and C.-J. Wang, *Chem. Commun.*, 2013, **49**, 6078; (d) J. Y. Lee, Y. S. You and S. H. Kang, *J. Am. Chem. Soc.*, 2011, **133**, 1772.
- For selected examples of using carbenoids to set quaternary stereocenters, see: (a) J. F. Briones and H. M. L. Davies, *J. Am. Chem. Soc.*, 2013, **135**, 13314; (b) J. E. Spangler and H. M. L. Davies, *J. Am. Chem. Soc.*, 2013, **135**, 6802; (c) T. K. Hyster, K. E. Ruhl and T. Rovis, *J. Am. Chem. Soc.*, 2013, **135**, 5364.
- D. H. T. Phan, K. G. M. Kou and V. M. Dong, *J. Am. Chem. Soc.*, 2010, **132**, 16354.
- (a) K.-J. Xiao, D. W. Lin, M. Miura, R.-Y. Zhu, W. Gong, M. Wasa and J.-Q. Yu, *J. Am. Chem. Soc.*, 2014, **136**, 8138; (b) B.-F. Shi, Y.-H. Zhang, J. K. Lam, D.-H. Wang and J.-Q. Yu, *J. Am. Chem. Soc.*, 2010, **132**, 460.
- Glorius recently reported cyclopentanones bearing α-quaternary centers by an organocatalyzed NHC hydroacylation: D. Janssen-Müller, M. Schedler, M. Fleige, C. G. Daniliuc and F. Glorius, *Angew. Chem., Int. Ed.*, 2015, DOI: 10.1002/anie.201412302.
- (a) R. T. Stemmler and C. Bolm, *Adv. Synth. Catal.*, 2007, **349**, 1185; (b) M. M. Coulter, K. G. M. Kou, B. Galligan and V. M. Dong, *J. Am. Chem. Soc.*, 2010, **132**, 16330; (c) M. M. Coulter, P. K. Dornan and V. M. Dong, *J. Am. Chem. Soc.*, 2009, **131**, 6932; (d) H. A. Khan, K. G. M. Kou and V. M. Dong, *Chem. Sci.*, 2011, **2**, 407; (e) J. D. Osborne,



H. E. Randell-Sly, G. S. Currie, A. R. Cowley and M. C. Willis, *J. Am. Chem. Soc.*, 2008, **130**, 17232; For recent enantioselective intramolecular hydroacylations without chelation-assistance, see: (f) X.-W. Du, A. Ghosh and L. M. Stanley, *Org. Lett.*, 2014, **16**, 4036; (g) A. Ghosh and L. M. Stanley, *Chem. Commun.*, 2014, **50**, 2765.

9 M. C. Willis, *Chem. Rev.*, 2010, **110**, 725.

10 K. Tanaka and G. C. Fu, *J. Am. Chem. Soc.*, 2002, **124**, 10296.

11 (a) M. Tobisu, I. Hyodo, M. Onoe and N. Chatani, *Chem. Commun.*, 2008, 6013; Olefin-assisted transformations using other transition-metal catalysts, see: (b) S. GÜLAK and A. J. von Wangelin, *Angew. Chem., Int. Ed.*, 2012, **51**, 1357; (c) P. Gandeepan and C.-H. Cheng, *J. Am. Chem. Soc.*, 2012, **134**, 5738.

12 (a) M. Tanaka, M. Imai, M. Fujio, E. Sakamoto, M. Takahashi, Y. Eto-Kato, X. M. Wu, K. Funakoshi, K. Sakai and H. Suemune, *J. Org. Chem.*, 2000, **65**, 5806; (b) M. Tanaka, M. Takahashi, E. Sakamoto, M. Imai, A. Matsui, M. Fujio, K. Funakoshi, K. Sakai and H. Suemune, *Tetrahedron*, 2001, **57**, 1197.

13 (a) T. J. Hoffman and E. M. Carreira, *Angew. Chem., Int. Ed.*, 2011, **50**, 10670; (b) R. E. Campbell Jr, C. F. Lochow, K. P. Vora and R. G. Miller, *J. Am. Chem. Soc.*, 1980, **102**, 5824; (c) R. C. Larock, K. Oertle and G. F. Potter, *J. Am. Chem. Soc.*, 1980, **102**, 190; There is some evidence for alkynes promoting C–H activation: (d) K. Tanaka and G. C. Fu, *Org. Lett.*, 2002, **4**, 933.

14 (a) D. P. Fairlie and B. Bosnich, *Organometallics*, 1988, **7**, 936; (b) S. Mukherjee and B. List, *J. Am. Chem. Soc.*, 2007, **129**, 11336; (c) M. Rolandsgård, S. Baldawi, D. Sirbu, V. Bjørnstad, C. Rømming and K. Undheim, *Tetrahedron*, 2005, **61**, 4129; (d) R. Okamoto and K. Tanaka, *Org. Lett.*, 2013, **15**, 2112.

15 G. Jiang and B. List, *Adv. Synth. Catal.*, 2011, **353**, 1667.

16 (a) Y. Shibata and K. Tanaka, *J. Am. Chem. Soc.*, 2009, **131**, 12552; (b) K. Tanaka, Y. Shibata, T. Suda, Y. Hagiwara and M. Hirano, *Org. Lett.*, 2007, **9**, 1215; (c) K. G. M. Kou, D. N. Le and V. M. Dong, *J. Am. Chem. Soc.*, 2014, **136**, 9471; (d) S. K. Murphy, A. Bruch and V. M. Dong, *Angew. Chem., Int. Ed.*, 2014, **53**, 2455.

17 For examples of carboacylation by C–C bond activation, see: (a) H. M. Ko and G. Dong, *Nat. Chem.*, 2014, **6**, 739; (b) L. Souillart and N. Cramer, *Angew. Chem., Int. Ed.*, 2014, **53**, 9640; (c) L. Souillart, E. Parker and N. Cramer, *Angew. Chem., Int. Ed.*, 2014, **53**, 3001; (d) T. Xu, H. M. Ko, N. A. Savage and G. Dong, *J. Am. Chem. Soc.*, 2012, **134**, 20005; (e) A. M. Dreis and C. J. Douglas, *J. Am. Chem. Soc.*, 2009, **131**, 412.

18 C. Aïssa, K. Y. T. Ho, D. J. Tetlow and M. Pin-Nó, *Angew. Chem., Int. Ed.*, 2014, **53**, 4209.

19 Experiments with 5 mol% catalyst at 40 °C gave a mixture of 2 and 12 (2 : 12 = 1 : 2).

20 Preliminary experiments using phenylbis(methallyl) aldehyde resulted in 55% isolated yield for the non-isomerized hydroacylation product 12, and no hydroacylation product obtained using phenylbis(crotyl) aldehyde.

21 An endocyclic β-hydride elimination within a metallacyclopentanone structure has been computed by DFT in a study on the mechanism of rhodium-catalyzed decarbonylation: I. F. D. Hyatt, H. K. Anderson, A. T. Morehead Jr and A. L. Sargent, *Organometallics*, 2008, **27**, 135.

22 S. Y. Y. Yip and C. Aïssa, *Angew. Chem., Int. Ed.*, 2015, **54**, 6870. In this work, isomerization by endocyclic β-hydride elimination was supported by a deuterium labelling study using phenyl(d₂-allyl)aldehydes.

23 X. Huang, J. Zhu and Z. Lin, *Organometallics*, 2004, **23**, 4154.

24 Further studies are required to identify the turn-over limiting step. On the basis of the IUPAC Gold book (<http://old.iupac.org/goldbook/P04862.pdf>), the selectivity-determining step is identical to, or occurs later than, the rate-controlling step on the reaction coordinate. See also: E. M. Simmons and J. F. Hartwig, *Angew. Chem., Int. Ed.*, 2012, **51**, 3066.

25 I. Omae, *J. Organomet. Chem.*, 2007, **692**, 2608.

