CrossMark
click for updatesCite this: *Chem. Sci.*, 2015, 6, 5595Received 12th May 2015
Accepted 24th June 2015

DOI: 10.1039/c5sc01737h

www.rsc.org/chemicalscience

Pd(II)-catalyzed remote regiodivergent *ortho*- and *meta*-C–H functionalizations of phenylethylamines†

Shangda Li,^{‡,a} Huafang Ji,^{‡,a} Lei Cai^a and Gang Li^{*ab}

Site selectivity control is of vital importance in the direct functionalization of inert C–H bonds. Reported here is a novel example of remote regiodivergent *ortho*- and *meta*-C–H bond functionalizations of phenylethylamine derivatives by using a novel 2-cyanobenzoyl group as the original directing functionality, where the regioselectivity was adjusted by a methylation. The potential of the method was exemplified by sequential functionalizations of both *ortho*- and *meta*-C–H bonds of a phenylethylamine derivative in a streamlined manner.

Introduction

Controlling site selectivity is an outstanding challenge in the direct functionalization of inert C–H bonds that are ubiquitous in organic molecules.¹ The increasing applications of these type of transformations in organic synthesis also demand accessibility to diverse site selectivities.² While numerous directing groups have been introduced to assist the cleavage of proximal *ortho*-C–H bonds in most cases with transition metals,^{1,3–10} directing group assisted *meta*-selective C–H functionalization of arenes has proved especially challenging and is still very rare.^{5,6,8,9} In 2009, a remarkable breakthrough was reported by Gaunt *et al.*, who developed a carbonyl group directed unprecedented *meta*-selective C–H arylation of anilides by using a Cu(II) catalyst and diaryliodonium salts.^{5a} This method was later extended to α -aryl carbonyl compounds by the same group.^{5b} Another impressive breakthrough came from the Frost group, who introduced an ingenious method of *meta*-selective C–H sulfonation of 2-phenylpyridines *via* cyclometalated ruthenium intermediates.^{6a,b} A similar strategy was then employed by Ackermann to realize a *meta*-selective C–H alkylation with secondary alkyl halides.^{6c} Recently, a small number of ground-breaking examples of Pd(II) catalyzed directed *meta*-selective C–H functionalizations of arenes that were attached with elegantly devised nitrile-based templates were disclosed, pioneered by Yu and then further

studied by Tan and Maiti.⁸ By using the above directing group assisted *meta*-selective C–H functionalization of arenes, elegant regiodivergent functionalizations of *ortho*- and *meta*-C–H bonds have been reported by Gaunt,^{4b,5b} Frost^{6b} and Yu,^{8b} and examples of reactions reported by Gaunt^{4b,5b} and Frost^{6b} could even be performed sequentially.^{8i,11,12} However, the use of analogous directing groups to achieve *remote-selective* regiodivergent activation of *ortho*- and *meta*-C–H bonds has not been examined and remains a significant challenge.^{13,14} We envision that such methodology is highly desirable for drug discovery and material sciences, since it only requires a single operation to achieve a different remote regioselectivity.^{2f} Herein, we report a novel strategy for regiodivergent *ortho*- and *meta*-C–H functionalizations of phenylethylamine derivatives.

To test our hypothesis of a regiodivergent C–H functionalization strategy by using analogous directing groups, we selected phenylethylamines as the testing compounds, since they are a class of aromatic compounds that are important core structures of numerous drug molecules (Fig. 1). Moreover, although *ortho*-C–H functionalizations have been reported for phenylethylamine derivatives, their *meta*-selective C–H functionalization remains elusive.¹⁵ Inspired by recent studies on directed *meta*-selective C–H functionalizations of arenes,⁸ we proposed that a 2-cyanobenzoyl group could act as the key directing functionality for both *ortho*- and *meta*-C–H functionalizations of phenylethylamines with a Pd(II) catalyst by taking advantage of the σ and π coordination ability of the nitrile group (Scheme 1).¹⁶ However, during our study we found that our proposed mode of *ortho*-selective C–H bond cleavage was not feasible and a novel remote-selective *ortho*-C–H bond cleavage was observed instead (*vide infra*).^{13,14}

Results and discussion

To examine our original hypothesis (Scheme 2), we chose olefination as the model reaction.^{10,17} After extensive condition

^aKey Laboratory of Coal to Ethylene Glycol and Its Related Technology, Fujian Institute of Research on the Structure of Matter, Chinese Academy of Sciences, 155 West Yang-Qiao Road, Fuzhou, Fujian, 350002, P. R. China. E-mail: gangli@fjirsm.ac.cn

^bState Key Laboratory of Structural Chemistry, Chinese Academy of Sciences, Fuzhou, Fujian, 350002, P. R. China

† Electronic supplementary information (ESI) available: Experimental procedures and data for new compounds. CCDC 1056709 and 1406595. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c5sc01737h

‡ These authors contributed equally.

Fig. 1 Representative drugs containing a phenylethylamine core.

Scheme 1 Hypothesis of regioselectivity changed by a methylation.

screening with $\text{Pd}(\text{OAc})_2$ as the catalyst (see ESI†), we were able to produce a high combined yield of *ortho*-olefinated products by treating **1a** with ethyl acrylate under oxygen with hexafluoroisopropanol (HFIP) as an additive and *N*-acetyl-glycine (Ac-Gly-OH) ligand.^{8a,18} Interestingly, the 2-cyanobenzoyl motif cyclized to an imidamide derivative in the products. To ascertain the mechanism of this olefination, **1a** was subjected to the above reaction conditions without adding ethyl acrylate, affording **1a'** that was believed to be the reactive substrate for the olefination. Indeed, after **1a'** was treated with the same olefination conditions, the desired products were generated in similar yields (see ESI†). Although this reaction pathway is not desired from our original hypothesis, the site selectivity of this reaction is surprisingly uncommon since the imino group of **1a'**, the most likely directing group on **1a'**, directed the cleavage of a remote *ortho*-C–H bond rather than a proximal *ortho*-C–H bond on the arene attached to the imidamide, which is in marked contrast to the *ortho*-C–H functionalizations of arylimine derivatives.¹⁹ The exact origin of the selectivity is unclear at present, and the study of the mechanism is under way.¹⁴

Several representative substrates were then surveyed briefly (Table 1). It was found that electron-withdrawing groups like chloride and fluoride were tolerated (**2b–c**), giving good yields of desired products. Good to excellent yields of products were also generated with substrates containing electron-donating groups such as methyl at the *meta*-position (**2d**) and methoxy at the *ortho*- (**2e**) and *para*-position (**2f**).

Having established the remote-selective *ortho*-C–H olefination of the secondary phenylethylamide, we were eager to test whether the selectivity could be switched to a remote-selective *meta*-C–H olefination after the secondary amide is methylated into a tertiary one (see the ESI† for methylation with MeI). Starting with the above *ortho*-olefination reaction conditions, we were very delighted to find that *N*-methyl amide **3a** could lead to a 10% yield of the desired product with a trace of other regioisomers (Table 2, entry 1). Inspired by the previous discovery that HFIP was a compatible solvent with nitrile-based templates,⁸ we switched the solvent to HFIP and found that the combined yield of desired products was increased dramatically

Scheme 2 A novel remote-selective *ortho*-C–H olefination.

to 58% with silver acetate as the sole oxidant (entry 2). Since when using weakly acidic HFIP as the sole solvent some substrate might decompose, DCE was added as the co-solvent, resulting in an increased yield of 73% (entry 3). To optimize the solvent system, we decreased the volume of HFIP to 15% and found that the combined yield was only slightly improved (entry 4). However, a further decreased volume of HFIP led to a much diminished yield (entry 5). Other solvents were also screened, but the combination of DCE and HFIP proved to be the best. The addition of a weak base, such as KHCO_3 , to tune the acidity of the reaction system was not effective either (entry 6). Since a higher catalytic turnover of the Pd catalyst was observed with 50% volume of HFIP, we repeated the reaction with this solvent system at 80 °C and found that the combined yield was improved to 90% in 32 hours under nitrogen (entry 7), representing the highest catalytic turnover of the Pd catalyst. Finally, by adding 5 equivalents of DMF we were able to get more mono-olefinated product in 28 hours while maintaining the overall efficiency (entry 8, see the ESI† for more condition screenings). However, further screening of reaction conditions could not result in better mono- vs. di-olefination selectivity at present,

Table 1 Representative substrates of remote *ortho*-C–H olefination^a

^a Reaction conditions: **1** (0.2 mmol), ethyl acrylate (0.4 mmol), $\text{Pd}(\text{OAc})_2$ (10 mol%), Ac-Gly-OH (20 mol%), HFIP (0.6 mmol), Ag_2CO_3 (0.06 mmol), O_2 (1 atm), *t*-amyl-OH (2 mL), 24–48 h, 90 °C. Isolated yields are reported, see the ESI for details. ^b 80 °C. ^c 70 °C.

Table 2 Screening of reaction conditions for *meta*-C–H olefination^a

Entry	Solvents [v/v]	T (°C)	Yield (%) [4a _{mono} , 4a _{(m,m')di}]	3a (%)
1 ^b	<i>t</i> -Amyl-OH	90	10 [10, 0]	90
2	HFIP	90	58 [13, 45]	Trace
3	DCE/HFIP [50/50]	90	71 [32, 39]	Trace
4	DCE/HFIP [85/15]	90	73 [48, 25]	10
5	DCE/HFIP [95/5]	90	39 [32, 7]	44
6 ^c	DCE/HFIP [85/15]	90	26 [26, 0]	55
7 ^{d,e}	DCE/HFIP [50/50]	80	90 [46, 44]	Trace
8 ^{d,f}	DCE/HFIP [50/50]	80	90 [58, 32]	Trace

^a Reaction conditions: **3a** (0.2 mmol), ethyl acrylate (0.4 mmol), Pd(OAc)₂ (10 mol%), Ac-Gly-OH (20 mol%), AgOAc (0.6 mmol), solvent (2 mL), 24 h, 80–90 °C. Yield was determined by ¹H NMR analysis using CH₂Br₂ as the internal standard. ^b Using the same conditions as in Scheme 2. ^c KHCO₃ (2 equiv.) was added. ^d Under N₂. ^e 32 h. Isolated yields were 45% of **4a_{mono}** and 37% of **4a_{(m,m')di}**. ^f 28 h, DMF (5 equiv.) was added.

and a study on this issue is actively being carried out in our laboratory. The *meta*-selectivity was unambiguously verified by X-ray crystallographic analysis of a derivative obtained by hydrolyzing the ester group of **4a_{mono}** (see the ESI†).

With the optimized conditions at hand, we examined the scope of this remote *meta*-selective olefination protocol (Table 3). *Ortho*-substituted substrates with both electron-donating methyl and methoxy and electron-withdrawing bromo and chloro groups proved to be suitable substrates, producing good combined yields of *meta*-olefinated products (**4b–4e**). It is worth noting that arenes bearing bromo or chloro substituents (**4d** and **4e**) were compatible substrates, enabling further elaboration at the halogenated positions. Moreover, although we could not circumvent di-olefination (**4b**, **4d–e**), the fact that both *meta*-positions of 2-substituted substrates could be functionalized provides a great opportunity for synthesis of diversely substituted arenes, which is particularly beneficial for the drug discovery industry. The remaining *meta*-position of *meta*-substituted substrates was also selectively olefinated in high yields (**4f–4i**). *Para*-substituted compounds carrying methoxy or halide groups were also viable substrates for obtaining high yields of the desired products (**4j–4l**). Notably, despite the steric hindrance, the olefin partner could also be installed selectively at the *meta*-position with poly-substituted substrates (**4m–4n**), displaying an uncommon procedure for constructing new penta-substituted phenylethylamines. It is interesting to note that the reaction was not sensitive to the difference of steric hindrance and both *meta*-positions of **3n** could be olefinated. Finally, substituents at the benzylic position were also tolerated (**4o** and **4p**), presenting the potential utility of this protocol with a drug molecule (**4p**). The *meta*-selectivity of various substrates

was generally excellent with only minor amounts of other isomers whose amounts were hard to determine due to the presence of rotamers in the ¹H NMR spectra of the crude olefinated products. The exceptional substrate is **3g**, which also generated around 10% of other isomeric products owing to the electron-donating methoxy substituent. However, it is notable that the intrinsic electronic biases of the molecules were overall successfully overridden (**4d–4f**, **4k–4n**). Moreover, removal of the directing group was smoothly realized by hydrolysis with HCl to afford high yields of new *meta*-substituted phenylethylamines (see the ESI†).

To further expand the scope of this reaction, we examined various olefin coupling partners and found olefination of **3f** with α,β -unsaturated ketone, amide and phosphonate afforded desired products in good yields (Table 4, **6a–6c**). We were also pleased to find *trans*-2-butenate reacted stereoselectively with **3f** to give **6d** in moderate yield. It is noteworthy that this reaction was also compatible with cyclic tri-substituted olefin to give high yield of allylated product (**6e**). Finally, electron deficient

Table 3 *meta*-Olefination of phenylethylamine derivatives^a

3a-p	4a-p
4a_{mono} : 45%	4b_{mono} : 59% ^{b,c}
4a_{(m,m')di} : 37%	4b_{(m,m')di} : 15%
4c : 61% ^c	4d_{mono} (m) : 11% ^{b,d}
4d_{mono} (m') : 39%	4d_{(m,m')di} : 29%
4e_{mono} (m) : 13% ^{b,d}	4f : 76% ^c
4e_{mono} (m') : 28%	4g : 63% ^{c,e}
4e_{(m,m')di} : 41%	4h : 78% ^c
4i : 78% ^{b,d}	4j : 74% ^c
4k_{mono} : 61% ^b	4l_{mono} : 59% ^b
4k_{(m,m')di} : 19%	4l_{(m,m')di} : 27%
4m_{mono} : 50% ^{b,d}	4n_{mono} (m) : 24% ^{b,d}
4m_{(m,m')di} : 24%	4n_{mono} (m') : 20%
4o_{mono} : 59% ^c	4p_{mono} : 37% ^b
4o_{(m,m')di} : 19%	4p_{(m,m')di} : 35%
Baclofen Derivatives	

^a Reaction conditions: **3** (0.2 mmol), ethyl acrylate (0.4 mmol), Pd(OAc)₂ (10 mol%), Ac-Gly-OH (20 mol%), AgOAc (0.6 mmol), DCE (1 mL), HFIP (1 mL), 24–48 h, 80 °C, N₂. Isolated yields are reported, see the ESI† for details. ^b 90 °C. ^c DMF (1 mmol) was added. ^d DCE (0 mL)/HFIP (2 mL).

^e Around 10% of other isomers detected by ¹H NMR.

Table 4 Scope of olefin coupling partners^a

^a Reaction conditions: **3f** (0.1 mmol), **5** (0.2 mmol), Pd(OAc)₂ (10 mol%), Ac-Gly-OH (20 mol%), AgOAc (0.3 mmol), DMF (0.5 mmol), DCE (0.5 mL), HFIP (0.5 mL), 28 h, 80 °C, N₂. Isolated yields are reported.

styrene such as pentafluorostyrene **5f** was also effective with this method to produce excellent yield of product (**6f**), albeit electron-rich styrenes were not applicable coupling partners.

Finally, to demonstrate the potential of our method for streamlined synthesis of highly substituted arenes, we first subjected **1c** to our standard *ortho*-olefination conditions with **5f** and obtained *ortho*-olefinated **7** in 86% yield (Scheme 3). Then, much to our delight, we were able to convert **7** to the desired amide **8** with the required nitrile group which was reconstructed simultaneously with methylation by using LiHMDS, followed by hydrogenation of the double bond.²⁰ Lastly, the *meta*-selective allylation proceeded efficiently with tri-substituted olefin **5e** to afford tetrasubstituted arene **9** in good yield, enabling the building of complexity in a concise manner.

Scheme 3 Sequential *ortho*- and *meta*-C–H functionalizations. (a) **5f** (2 equiv.), Pd(OAc)₂ (10 mol%), Ac-Gly-OH (20 mol%), HFIP (3 equiv.), Ag₂CO₃ (30 mol%), O₂ (1 atm), *t*-amyl-OH, 24 h, 90 °C, 86% yield; (b) LiHMDS (2.5 equiv.), MeI (3 equiv.), THF, –15 °C, 58% yield (85% yield based on recovered starting material [brsm]); (c) Pd/C (12 mol%), H₂ (1 atm), MeOH, 98%; (d) **5e** (2 equiv.), Pd(OAc)₂ (10 mol%), Ac-Gly-OH (20 mol%), AgOAc (3 equiv.), DCE (1 mL)/HFIP (1 mL), 48 h, 90 °C, N₂, 73%.

Conclusions

In summary, a novel example of remote regiodivergent *ortho*- and *meta*-C–H functionalizations has been developed with phenylethylamine derivatives by introducing a novel 2-cyano-benzoyl group as the original directing functionality. A single methylation was sufficient to switch the remote regioselectivity. This method also enabled the novel sequential functionalizations of *ortho*- and *meta*-C–H bonds of a phenylethylamine derivative. Further development of this strategy will improve C–H functionalization to become a more versatile synthetic tool.

Acknowledgements

This work was supported by the Fujian Institute of Research on the Structure of Matter, NSFC (21402198) and “The 1000 Youth Talents Program”. We gratefully appreciate Prof. Jin-Quan Yu at The Scripps Research Institute, Prof. Weiping Su at FJIRSM and Prof. Richard P. Hsung at the University of Wisconsin-Madison for helpful comments. We also thank Prof. Daqiang Yuan at FJIRSM for crystallographic analysis.

Notes and references

- Selected recent reviews: (a) D. Alberico, M. E. Scott and M. Lautens, *Chem. Rev.*, 2007, **107**, 174; (b) I. V. Seregin and V. Gevorgyan, *Chem. Soc. Rev.*, 2007, **36**, 1173; (c) F. Kakiuchi and T. Kochi, *Synthesis*, 2008, **2008**, 3013; (d) X. Chen, K. M. Engle, D.-H. Wang and J.-Q. Yu, *Angew. Chem., Int. Ed.*, 2009, **48**, 5094; (e) O. Daugulis, H.-Q. Do and D. Shabashov, *Acc. Chem. Res.*, 2009, **42**, 1074; (f) T. W. Lyons and M. S. Sanford, *Chem. Rev.*, 2010, **110**, 1147; (g) D. A. Colby, R. G. Bergman and J. A. Ellman, *Chem. Rev.*, 2010, **110**, 624; (h) C.-L. Sun, B.-J. Li and Z.-J. Shi, *Chem. Rev.*, 2010, **111**, 1293; (i) T. Satoh and M. Miura, *Synthesis*, 2010, **2010**, 3395; (j) O. Baudoin, *Chem. Soc. Rev.*, 2011, **40**, 4902; (k) J. F. Hartwig, *Chem. Soc. Rev.*, 2011, **40**, 1992; (l) S. H. Cho, J. Y. Kim, J. Kwak and S. Chang, *Chem. Soc. Rev.*, 2011, **40**, 5068; (m) M. S. Sigman and E. W. Werner, *Acc. Chem. Res.*, 2012, **45**, 874; (n) C. Liu, H. Zhang, W. Shi and A. Lei, *Chem. Rev.*, 2011, **111**, 1780; (o) C. S. Yeung and V. M. Dong, *Chem. Rev.*, 2011, **111**, 1215; (p) S. R. Neufeldt and M. S. Sanford, *Acc. Chem. Res.*, 2012, **45**, 936; (q) N. Kuhl, M. N. Hopkinson, J. Wencel-Delord and F. Glorius, *Angew. Chem., Int. Ed.*, 2012, **51**, 10236; (r) P. B. Arockiam, C. Bruneau and P. H. Dixneuf, *Chem. Rev.*, 2012, **112**, 5879; (s) M. C. White, *Science*, 2012, **335**, 807; (t) G. Rouquet and N. Chatani, *Angew. Chem., Int. Ed.*, 2013, **52**, 11726; (u) L. Ackermann, *Acc. Chem. Res.*, 2014, **47**, 281; (v) M. Zhang, Y. Zhang, X. Jie, H. Zhao, G. Li and W. Su, *Org. Chem. Front.*, 2014, **1**, 843; (w) J. J. Topczewski and M. S. Sanford, *Chem. Sci.*, 2015, **6**, 70.
- (a) H. M. L. Davies and J. R. Manning, *Nature*, 2008, **451**, 417; (b) L. McMurray, F. O'Hara and M. J. Gaunt, *Chem. Soc. Rev.*, 2011, **40**, 1885; (c) W. R. Gutekunst and P. S. Baran, *Chem.*

- Soc. Rev.*, 2011, **40**, 1976; (d) D. Y. K. Chen and S. W. Youn, *Chem.-Eur. J.*, 2012, **18**, 9452; (e) J. Yamaguchi, A. D. Yamaguchi and K. Itami, *Angew. Chem., Int. Ed.*, 2012, **51**, 8960; (f) W.-D. Joanna and G. Frank, *Nat. Chem.*, 2013, **5**, 369.
- 3 Selected examples of Pd-catalyzed regioselective C-H functionalizations: (a) N. R. Deprez, D. Kalyani, A. Krause and M. S. Sanford, *J. Am. Chem. Soc.*, 2006, **128**, 4972; (b) D. R. Stuart and K. Fagnou, *Science*, 2007, **316**, 1172; (c) C. S. Yeung, X. Zhao, N. Borduas and V. M. Dong, *Chem. Sci.*, 2010, **1**, 331; (d) K. Ueda, S. Yanagisawa, J. Yamaguchi and K. Itami, *Angew. Chem., Int. Ed.*, 2010, **49**, 8946; (e) A. J. Hickman and M. S. Sanford, *ACS Catal.*, 2011, **1**, 170; (f) J. Karthikeyan and C.-H. Cheng, *Angew. Chem., Int. Ed.*, 2011, **50**, 9880; (g) A. N. Campbell, E. B. Meyer and S. S. Stahl, *Chem. Commun.*, 2011, **47**, 10257; (h) X. Wang, D. Leow and J.-Q. Yu, *J. Am. Chem. Soc.*, 2011, **133**, 13864; (i) K. Sun, Y. Li, T. Xiong, J. Zhang and Q. Zhang, *J. Am. Chem. Soc.*, 2011, **133**, 1694; (j) L. Jiao and T. Bach, *J. Am. Chem. Soc.*, 2011, **133**, 12990; (k) Z. Wu, F. Luo, S. Chen, Z. Li, H. Xiang and X. Zhou, *Chem. Commun.*, 2013, **49**, 7653; (l) R. Shrestha, P. Mukherjee, Y. Tan, Z. C. Litman and J. F. Hartwig, *J. Am. Chem. Soc.*, 2013, **135**, 8480; (m) D.-T. D. Tang, K. D. Collins, J. B. Ernst and F. Glorius, *Angew. Chem., Int. Ed.*, 2014, **53**, 1809.
- 4 Selected examples of non-Pd-catalyzed regioselective C-H functionalizations: (a) H. Chen, S. Schlecht, T. C. Semple and J. F. Hartwig, *Science*, 2000, **287**, 1995; (b) C.-L. Ciana, R. J. Phipps, J. R. Brandt, F.-M. Meyer and M. J. Gaunt, *Angew. Chem., Int. Ed.*, 2011, **50**, 458; (c) S. Mochida, K. Hirano, T. Satoh and M. Miura, *J. Org. Chem.*, 2011, **76**, 3024; (d) J. Kwak, M. Kim and S. Chang, *J. Am. Chem. Soc.*, 2011, **133**, 3780; (e) B. Li, Z.-H. Wu, Y.-F. Gu, C.-L. Sun, B.-Q. Wang and Z.-J. Shi, *Angew. Chem., Int. Ed.*, 2011, **50**, 1109; (f) J. Wencel-Delord, C. Nimphius, F. W. Patureau and F. Glorius, *Chem.-Asian J.*, 2012, **7**, 1208; (g) T. Andou, Y. Saga, H. Komai, S. Matsunaga and M. Kanai, *Angew. Chem., Int. Ed.*, 2013, **52**, 3213; (h) K. Ueda, K. Amaike, R. M. Maceiczky, K. Itami and J. Yamaguchi, *J. Am. Chem. Soc.*, 2014, **136**, 13226.
- 5 (a) R. J. Phipps and M. J. Gaunt, *Science*, 2009, **323**, 1593; (b) H. A. Duong, R. E. Gilligan, M. L. Cooke, R. J. Phipps and M. J. Gaunt, *Angew. Chem., Int. Ed.*, 2011, **50**, 463.
- 6 (a) O. Saidi, J. Marafie, A. E. W. Ledger, P. M. Liu, M. F. Mahon, G. Kociok-Köhn, M. K. Whittlesey and C. G. Frost, *J. Am. Chem. Soc.*, 2011, **133**, 19298; (b) C. Frost, W. Reynolds, P. Liu and G. Kociok-Köhn, *Synlett*, 2013, **24**, 2687; (c) N. Hofmann and L. Ackermann, *J. Am. Chem. Soc.*, 2013, **135**, 5877.
- 7 For other examples of non-Pd-catalyzed *meta*-C-H functionalizations: (a) J.-Y. Cho, M. K. Tse, D. Holmes, R. E. Maleczka and M. R. Smith, *Science*, 2002, **295**, 305; (b) T. Ishiyama, J. Takagi, K. Ishida, N. Miyaara, N. R. Anastasi and J. F. Hartwig, *J. Am. Chem. Soc.*, 2002, **124**, 390; (c) H.-Q. Do, R. M. K. Khan and O. Daugulis, *J. Am. Chem. Soc.*, 2008, **130**, 15185; (d) B.-J. Li and Z.-J. Shi, *Chem. Sci.*, 2011, **2**, 488; (e) C. Cheng and J. F. Hartwig, *Science*, 2014, **343**, 853.
- 8 For Pd-catalyzed *meta*-C-H functionalizations using a chelating directing template: (a) D. Leow, G. Li, T.-S. Mei and J.-Q. Yu, *Nature*, 2012, **486**, 518; (b) H.-X. Dai, G. Li, X.-G. Zhang, A. F. Stepan and J.-Q. Yu, *J. Am. Chem. Soc.*, 2013, **135**, 7567; (c) L. Wan, N. Dastbaravardeh, G. Li and J.-Q. Yu, *J. Am. Chem. Soc.*, 2013, **135**, 18056; (d) R.-Y. Tang, G. Li and J.-Q. Yu, *Nature*, 2014, **507**, 215; (e) G. Yang, P. Lindovska, D. Zhu, J. Kim, P. Wang, R.-Y. Tang, M. Movassaghi and J.-Q. Yu, *J. Am. Chem. Soc.*, 2014, **136**, 10807; (f) Y. Deng and J.-Q. Yu, *Angew. Chem., Int. Ed.*, 2015, **54**, 888; (g) S. Lee, H. Lee and K. L. Tan, *J. Am. Chem. Soc.*, 2013, **135**, 18778; (h) M. Bera, A. Modak, T. Patra, A. Maji and D. Maiti, *Org. Lett.*, 2014, **16**, 5760; (i) M. Bera, A. Maji, S. K. Sahoo and D. Maiti, *Angew. Chem., Int. Ed.*, 2015, DOI: 10.1002/anie.201503112.
- 9 During the preparation of our manuscript, two impressive reports on *meta*-selective C-H functionalizations *via* Pd/norbornene catalysis appeared: (a) X.-C. Wang, W. Gong, L.-Z. Fang, R.-Y. Zhu, S. Li, K. M. Engle and J.-Q. Yu, *Nature*, 2015, **519**, 334; (b) Z. Dong, J. Wang and G. Dong, *J. Am. Chem. Soc.*, 2015, **137**, 5887.
- 10 For other Pd-catalyzed *meta*-C-H functionalizations: (a) Y.-H. Zhang, B.-F. Shi and J.-Q. Yu, *J. Am. Chem. Soc.*, 2009, **131**, 5072; (b) J. Cornella, M. Righi and I. Larrosa, *Angew. Chem., Int. Ed.*, 2011, **50**, 9429; (c) M. Ye, G.-L. Gao and J.-Q. Yu, *J. Am. Chem. Soc.*, 2011, **133**, 6964; (d) M. Ye, G.-L. Gao, A. J. F. Edmunds, P. A. Worthington, J. A. Morris and J.-Q. Yu, *J. Am. Chem. Soc.*, 2011, **133**, 19090; (e) L. Zhou and W. Lu, *Organometallics*, 2012, **31**, 2124; (f) Y.-N. Wang, X.-Q. Guo, X.-H. Zhu, R. Zhong, L.-H. Cai and X.-F. Hou, *Chem. Commun.*, 2012, **48**, 10437; (g) F. Dai, Q. Gui, J. Liu, Z. Yang, X. Chen, R. Guo and Z. Tan, *Chem. Commun.*, 2013, **49**, 4634; (h) X. Cong, H. Tang, C. Wu and X. Zeng, *Organometallics*, 2013, **32**, 6565; (i) J. Luo, S. Preciado and I. Larrosa, *J. Am. Chem. Soc.*, 2014, **136**, 4109.
- 11 For other types of metal-catalyzed regiodivergent and/or sequential C-H functionalizations: (a) N. P. Grimster, C. Gauntlett, C. R. A. Godfrey and M. J. Gaunt, *Angew. Chem., Int. Ed.*, 2005, **44**, 3125; (b) E. M. Beck, N. P. Grimster, R. Hatley and M. J. Gaunt, *J. Am. Chem. Soc.*, 2006, **128**, 2528; (c) D. R. Stuart, E. Villemure and K. Fagnou, *J. Am. Chem. Soc.*, 2007, **129**, 12072; (d) E. M. Beck, R. Hatley and M. J. Gaunt, *Angew. Chem., Int. Ed.*, 2008, **47**, 3004; (e) T. W. Lyons, K. L. Hull and M. S. Sanford, *J. Am. Chem. Soc.*, 2011, **133**, 4455; (f) D. Katayev, M. Nakanishi, T. Burgi and E. P. Kundig, *Chem. Sci.*, 2012, **3**, 1422; (g) A. M. Wagner, A. J. Hickman and M. S. Sanford, *J. Am. Chem. Soc.*, 2013, **135**, 15710; (h) J. D. Dooley, S. Reddy Chidipudi and H. W. Lam, *J. Am. Chem. Soc.*, 2013, **135**, 10829; (i) Y. Su, H. Zhou, J. Chen, J. Xu, X. Wu, A. Lin and H. Yao, *Org. Lett.*, 2014, **16**, 4884; (j) N. Schroder, F. Lied and F. Glorius, *J. Am. Chem. Soc.*, 2015, **137**, 1448.
- 12 For more selected examples of sequential C-H functionalizations: (a) S. Yanagisawa, K. Ueda, H. Sekizawa

- and K. Itami, *J. Am. Chem. Soc.*, 2009, **131**, 14622; (b) K. M. Engle, D.-H. Wang and J.-Q. Yu, *Angew. Chem., Int. Ed.*, 2010, **49**, 6169; (c) W. R. Gutekunst and P. S. Baran, *J. Am. Chem. Soc.*, 2011, **133**, 19076; (d) H. Wang, G. Li, K. M. Engle, J.-Q. Yu and H. M. L. Davies, *J. Am. Chem. Soc.*, 2013, **135**, 6774; (e) H. J. Kim, M. J. Ajitha, Y. Lee, J. Ryu, J. Kim, Y. Lee, Y. Jung and S. Chang, *J. Am. Chem. Soc.*, 2014, **136**, 1132; (f) A. D. Yamaguchi, K. M. Chepiga, J. Yamaguchi, K. Itami and H. M. Davies, *J. Am. Chem. Soc.*, 2015, **137**, 644.
- 13 For remote C–H functionalizations: (a) R. Breslow, *Acc. Chem. Res.*, 1980, **13**, 170; (b) H. Schwarz, *Acc. Chem. Res.*, 1989, **22**, 282; (c) S. Das, C. D. Incarvito, R. H. Crabtree and G. W. Brudvig, *Science*, 2006, **312**, 1941; (d) J.-J. Li, R. Giri and J.-Q. Yu, *Tetrahedron*, 2008, **64**, 6979; (e) D. G. Pintori and M. F. Greaney, *J. Am. Chem. Soc.*, 2010, **133**, 1209; (f) E. E. Stache, C. A. Seizert and E. M. Ferreira, *Chem. Sci.*, 2012, **3**, 1623; (g) W. A. Nack, G. He, S.-Y. Zhang, C. Lu and G. Chen, *Org. Lett.*, 2013, **15**, 3440; (h) J. Schranck, A. Tlili and M. Beller, *Angew. Chem., Int. Ed.*, 2014, **53**, 9426.
- 14 For remote-selective *ortho*-C–H bond functionalization: (a) T. Saget and N. Cramer, *Angew. Chem., Int. Ed.*, 2013, **52**, 7865; *Angew. Chem.*, 2013, **125**, 8019; (b) W. B. Cross, E. G. Hope, Y.-H. Lin, S. A. Macgregor, K. Singh, G. A. Solan and N. Yahya, *Chem. Commun.*, 2013, **49**, 1918; (c) For a reverse selectivity, see: M. Lafrance, S. I. Gorelsky and K. Fagnou, *J. Am. Chem. Soc.*, 2007, **129**, 14570.
- 15 (a) J.-J. Li, T.-S. Mei and J.-Q. Yu, *Angew. Chem., Int. Ed.*, 2008, **47**, 6452; (b) G. He, C. Lu, Y. Zhao, W. A. Nack and G. Chen, *Org. Lett.*, 2012, **14**, 2944; (c) T.-S. Mei, D. Leow, H. Xiao, B. N. Laforteza and J.-Q. Yu, *Org. Lett.*, 2013, **15**, 3058; (d) A. Rodriguez, J. Albert, X. Ariza, J. Garcia, J. Granell, J. Farras, A. La Mela and E. Nicolas, *J. Org. Chem.*, 2014, **79**, 9578; (e) X. Ye and X. Shi, *Org. Lett.*, 2014, **16**, 4448; (f) J. Han, P. Liu, C. Wang, Q. Wang, J. Zhang, Y. Zhao, D. Shi, Z. Huang and Y. Zhao, *Org. Lett.*, 2014, **16**, 5682.
- 16 For π -coordination of nitrile group in C–H functionalizations: (a) P. Gandeepan and C. H. Cheng, *Chem.-Asian J.*, 2015, **10**, 824; (b) F. Kakiuchi, M. Sonoda, T. Tsujimoto, N. Chatani and S. Murai, *Chem. Lett.*, 1999, **28**, 1083; (c) W. Li, Z. Xu, P. Sun, X. Jiang and M. Fang, *Org. Lett.*, 2011, **13**, 1286; (d) J.-C. Wan, J.-M. Huang, Y.-H. Jhan and J.-C. Hsieh, *Org. Lett.*, 2013, **15**, 2742.
- 17 For reviews and selected recent examples of Pd-catalyzed C–H olefinations: (a) C. Jia, T. Kitamura and Y. Fujiwara, *Acc. Chem. Res.*, 2001, **34**, 633; (b) J. Le Bras and J. Muzart, *Chem. Rev.*, 2011, **111**, 1170; (c) L. Zhou and W. Lu, *Chem.-Eur. J.*, 2014, **20**, 634; (d) K. J. Stowers, K. C. Fortner and M. S. Sanford, *J. Am. Chem. Soc.*, 2011, **133**, 6541; (e) C. Huang, B. Chattopadhyay and V. Gevorgyan, *J. Am. Chem. Soc.*, 2011, **133**, 12406; (f) P. Gandeepan and C. H. Cheng, *J. Am. Chem. Soc.*, 2012, **134**, 5738; (g) R. Shi, L. Lu, H. Zhang, B. Chen, Y. Sha, C. Liu and A. Lei, *Angew. Chem., Int. Ed.*, 2013, **52**, 10582; (h) Y. Shang, X. Jie, J. Zhou, P. Hu, S. Huang and W. Su, *Angew. Chem., Int. Ed.*, 2013, **52**, 1299; (i) A. Deb, S. Bag, R. Kancherla and D. Maiti, *J. Am. Chem. Soc.*, 2014, **136**, 13602.
- 18 (a) K. M. Engle, D.-H. Wang and J.-Q. Yu, *J. Am. Chem. Soc.*, 2010, **132**, 14137; (b) G.-J. Cheng, Y.-F. Yang, P. Liu, P. Chen, T.-Y. Sun, G. Li, X.-H. Zhang, K. N. Houk, J.-Q. Yu and Y.-D. Wu, *J. Am. Chem. Soc.*, 2014, **136**, 894.
- 19 Selected examples: (a) A. R. Dick, K. L. Hull and M. S. Sanford, *J. Am. Chem. Soc.*, 2004, **126**, 2300; (b) H.-Y. Thu, W.-Y. Yu and C.-M. Che, *J. Am. Chem. Soc.*, 2006, **128**, 9048; (c) V. S. Thirunavukkarasu, K. Parthasarathy and C.-H. Cheng, *Angew. Chem., Int. Ed.*, 2008, **47**, 9462; (d) Y. Tan and J. F. Hartwig, *J. Am. Chem. Soc.*, 2010, **132**, 3676; (e) S.-J. Lou, D.-Q. Xu and Z.-Y. Xu, *Angew. Chem., Int. Ed.*, 2014, **53**, 10330.
- 20 The hydrogenation after the methylation was necessary to avoid the resulting substrate with a double bond acting as an olefin coupling partner in the following *meta*-selective olefination step.