

Cite this: *Chem. Sci.*, 2015, **6**, 6392

A T-shaped $\text{Ni}[\kappa^2-(\text{CF}_2)_4]$ NHC complex: unusual $\text{C}_{\text{sp}}^3-\text{F}$ and $\text{M}-\text{C}^{\text{F}}$ bond functionalization reactions[†]

Nicholas O. Andrella, Alexandre J. Sicard, Serge I. Gorelsky, Ilia Korobkov and R. Tom Baker*

Received 25th May 2015
Accepted 27th July 2015DOI: 10.1039/c5sc01886b
www.rsc.org/chemicalscience

Introduction

Fluorocarbons and their derivatives are valuable as refrigerants, agrochemicals, unique solvents/surfactants and fluoropharmaceuticals, with annual sales of the latter alone in the billions of dollars.¹ As the demand for fluorinated chemicals has increased, so too have synthetic methods for introducing fluorine and fluorocarbon groups.^{2–4} Despite recent advances, transition metal-mediated/-catalyzed routes are rare in comparison to the well-developed organometallic chemistry of hydrocarbons.⁵ The challenge rests in the stability of metal–perfluoroalkyl ($\text{M}-\text{R}^{\text{F}}$) bonds, relative to metal–alkyl bonds.⁶ $\text{M}-\text{C}^{\text{F}}$ bonds are typically inert to processes such as insertion/alkyl migration reactions, vital to metal-mediated catalytic cycles.⁷ Moreover, $\text{C}-\text{F}$ bonds are stronger than $\text{C}-\text{H}$ bonds,^{1a,b} posing another obstacle to metal-based approaches.


We are investigating perfluoronickelacyclopentane complexes (PNCPs) as platforms for functionalized fluorocarbons with an initial focus on fundamental stoichiometric reactions. PNCPs have been synthesized previously by reaction of tetrafluoroethylene (TFE, $\text{CF}_2=\text{CF}_2$) with Ni^0 complexes. The displacement of P ligands by bidentate ligands has also been reported (Scheme 1).⁸

To date, reports concerning the reactivity of PNCPs are sparse: Burch and co-workers found that Lewis acidic BF_3 effects fluoride abstraction from $\text{C}\alpha$ and phosphine migration to the activated carbon (Scheme 2a).⁹ Extending this reaction to the unsymmetrical $\text{P}^{\text{S}}\text{S}$ chelate, we showed that treatment with

excess isonitrile effected cleavage of the $\text{Ni}-\text{C}^{\text{F}}$ bond (Scheme 2b).¹⁰ With phosphite co-ligands, a remarkable hydrogenolysis reaction enables the selective catalytic hydrodimerization of TFE (Scheme 2c).¹¹ As far as we know, this reaction is the only example of a perfluoro-metallacyclopentane participating in a catalytic cycle.¹²

Our approach to metallacycle functionalization hinges on the reactivity of metal-activated $\text{C}\alpha-\text{F}$ bonds¹³ wherein we replace a $\text{C}-\text{F}$ bond by $\text{C}-\text{Nu}$ vs. the current paradigm $\text{C}-\text{L}$ (Nu = nucleophile, L = ligand). Using N-heterocyclic carbenes (NHCs),¹⁴ we aimed to access low-coordinate PNCPs wherein the strong $\text{M}-\text{C}_{\text{NHC}}$ bond may also prevent ligand migration to $\text{C}\alpha$. There is considerable precedent for such an approach to low-coordinate metal complexes.¹⁵ Hillhouse and coworkers prepared a two-coordinate nickel-imido complex bearing the exceptionally bulky IPr^* ligand (analog of IPr with 2,6-bis(diphenyl-methyl)phenyl groups instead of 2,6-diisopropylphenyl).¹⁶ Similarly, Miyazaki and coworkers synthesized a T-shaped three-coordinate nickel(i) chloride species $[\text{Ni}(\text{IPr})_2\text{Cl}]$ by treatment of two-coordinate $[\text{Ni}(\text{IPr})_2]$ with aryl chlorides.¹⁷ Also, Hartwig *et al.* synthesized a low-valent, three-coordinate palladium(ii) norbornyl species $[\text{Pd}(\text{SIPr})(\text{NAr})(\text{Nor})]$, which underwent facile $\text{C}-\text{N}$ bond reductive elimination when heated.¹⁸

In this report we show that low-coordinate NHC Ni perfluorometallacycles undergo facile $\text{C}_{\text{sp}}^3-\text{F}$ and $\text{M}-\text{C}^{\text{F}}$ bond cleavage as well as $\text{C}\alpha$ -functionalization.⁹ We also demonstrate






Scheme 1 Synthesis of perfluoronickelacyclopentanes.

Department of Chemistry and Centre for Catalysis Research and Innovation (CCRI), University of Ottawa, 30 Marie Curie, Ottawa, ON K1N 6N5 Canada

† Electronic supplementary information (ESI) available: Experimental details, NMR spectra and X-ray crystallographic information. CCDC files 968465 (2), 968466 (3), 968467 (4a), 1028645 (5c) and 1412522 (3· H_2O); For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c5sc01886b



(a) Ca-F activation / phosphine migration (Burch *et al.*)(b) Ca-F activation / phosphine migration and ring-opening (Baker *et al.*)(c) Catalytic hydrodimerization of tetrafluoroethylene (Baker *et al.*)


Scheme 2 Previously reported reactivity of perfluoronickelacyclopentanes.

the first migration of a fluoroalkyl to a reactive carbon center. This is significantly different from the reactivity previously observed for phosphine Ni perflurometallacycle complexes.⁹

Results and discussion

Starting from bis(phosphite) PNCPs¹⁰ (**1a** and **1b**) we were able to cleanly synthesize coordinatively-saturated or -unsaturated nickel perflurometallacycles. Thus, **1a** reacts smoothly with 1 equiv. of *i*tBu (*i*tBu = 1,3-di-*tert*-butylimidazol-2-ylidene) to afford the NHC/phosphite product **2** (Scheme 3, top; X-ray structural characterization presented in ESI†).¹⁹ Significantly, the reaction between the larger SIPr ligand [SIPr = 1,3-bis(2,6-diisopropylphenyl)imidazolin-2-ylidene] and a nickel metallacycle with sterically-demanding co-ligands (**1b**) results in displacement of both phosphite ligands, yielding the pseudo-three-coordinate/14e-Ni(II) metallacycle **3** (Scheme 3, bottom).


The molecular structure of complex **3**, as determined by single crystal X-ray diffraction, exhibits a T-shaped coordination about the Ni and features a weak agostic interaction with the isopropyl methyl group (Ni–C = 2.757(1) Å; compare Ni–C^F bond distance trans to the NHC (1.934(1) Å) with that trans to the agostic interaction (1.875(1) Å) (Fig. 1a). The ¹⁹F NMR



Scheme 3 Synthesis of NHC perfluoronickelacyclopentane complexes.

spectrum of **3** in C₆D₆ is consistent with ideal C_{2v} symmetry at room temperature, with only two distinct singlet resonances at –101.9 (F α) and –138.6 ppm (F β). While these resonances both broaden significantly at 213 K, it is apparent that the T-flip interconversion encounters only a small energy barrier.²⁰ To confirm this, we carried out DFT calculations (at the B3LYP/TZVP level with and without the empirical dispersion correction of Grimme).²¹ Intriguingly, the calculations reveal two spin-singlet structures with a very small energy difference ($\Delta G_{298\text{ K}} = 0.0\text{--}1.5\text{ kcal mol}^{-1}$). The first computed structure coincides well with the observed solid-state structure of **3**; the 3-center bond order index between the Ni and the corresponding C–H bond of 0.05 is much less than 8/27 (~0.3), the maximum possible value for a 3-center 2-electron bond. As a result, the Mayer valence index for Ni in structure **3** is only 3.09. The second structure (**3'**) features a weak η^3 interaction between the aryl group of the NHC ligand at the 4th coordination site of the Ni atom (Fig. 1b). Mayer bond orders for the corresponding three Ni–C interactions are in the 0.02–0.05 range, with a total bond order of 0.09. This suggests a semi-bidentate binding mode for the class of NHC ligands possessing pendant aryl groups. From calculations with the dispersion correction, structure **3'** has the same Gibbs free energy as **3**. Without the dispersion correction, structure **3'** is actually 1.5 kcal mol⁻¹ lower in energy than **3**. The 3-coordinate structure with trigonal coordination around Ni and symmetric binding of the C₄F₈ ligand is a transition state with a low energy ($\Delta G^\ddagger_{298\text{ K}} = 2.1\text{ kcal mol}^{-1}$ relative to **3**). Thus, it is clear that cleavage of the weak agostic and/or η^3 -aryl bond is facile and allows for rapid reorientation of the ligands around the Ni center. Attempts to obtain evidence for structure **3'** by low temperature NMR were frustrated by dynamic processes associated with the T-flip and hindered rotations about the M–C and perhaps N–C bonds.

The HOMO of **3** ($\varepsilon = -6.01\text{ eV}$; Fig. 2, left) is localized on the Ni (87%), primarily from a d_{z²} orbital contribution (71%). Lower-lying orbitals display interactions between metal d_{xz}, d_{yz} orbitals and the π -system of the aryl group.²² The LUMO ($\varepsilon = -1.96\text{ eV}$; Fig. 3, right) is an anti-bonding combination of the metal d_{x²-y²} orbital (total Ni character of 45%) with the π -donor

Fig. 1 (A) ORTEP representation of the molecular structure of **3**. Thermal-ellipsoid probabilities are set to 35% with hydrogen atoms omitted for clarity. The Ni–C(1) distance is 1.854(2) Å. (B) Optimized structure of low energy Ni–aryl isomer **3'**; Ni–C_{aryl} distances are = 2.818, 3.329, 3.379, 4.166, 4.204, 4.543 Å. The Ni–C(17) distance is 1.989 Å.




Fig. 2 The HOMO (left) and LUMO (right) of 3. Isosurface values of 0.04 au are used.

orbitals of the NHC and C_4F_8 ligands. Thus, reactivity of the M–C bond is likely under orbital control and arises from an interaction with the HOMO of 3. In contrast, C–F bond activation is likely a combination of orbital and charge control with a slant towards the latter as the hardness of the Lewis acid increases.²³

Initial studies on the C–F bond activation reactions of 3 are promising in the context of synthesizing functionalized fluorocarbons by metal-mediated approaches. Firstly, when 3 is treated with the Lewis acid TMSOTf (TMS = Me_3Si , OTf = SO_3CF_3), α -fluoride-abstraction, accompanied by Ni–C^F bond cleavage and C^F–C^F bond formation, furnishes a rare perfluorocyclobutyl complex 4a (Scheme 4, 75% isolated yield).²⁴ The driving force behind this transformation is likely related to the triflate leaving group ability and the formation of a strong C–C bond.²⁵ Importantly, the NHC remains bound to the nickel atom (*i.e.*, does not migrate to $C\alpha$), potentially opening new pathways to functionalized fluorocarbon derivatives. Upon heating complex 4a (80 °C in C_6D_6 , 24 h), perfluorocyclobutene is produced, presumably *via* a β -fluoride elimination mechanism, although the metal-containing co-product(s) have not yet been identified.²⁶ Interestingly, a single OTf containing product can be discerned by ¹⁹F NMR (−93.37 ppm) but a Ni–F signal could not be located. The ¹H NMR shows that the NHC remains intact. Upon addition of PPh_3 to the reaction mixture, PPh_3F_2

was identified as a major product, suggesting formation of a Ni–F thermolysis co-product.

The distorted square planar structure of complex 4a (Fig. 3) features a bidentate triflate ligand which can also likely access the κ^1 -mode in solution as evidenced by the simple ¹⁹F NMR spectrum²⁷ and observed tendency to eliminate. The perfluorocyclobutyl ring is nearly planar, the Ni–C bond is short [1.890(2) Å] due to the weak σ -trans influence ligand, and the $C\alpha$ –F bond distance (1.384(3) Å) is considerably longer than the other C–F bonds (average of 1.33 Å).

The reactivity enhancement offered by low-coordinate 3 is evidenced by the sluggish reaction of 4-coordinate complex 2 with TMSOTf to give a mixture of unidentified products. Indeed, monitoring the reaction of 3 and TMSOTf at −25 °C allowed for the identification of a Ni–C₄F₇ intermediate 5a apparently containing a $C\alpha$ –OTf linkage (triflate CF_3 ¹⁹F NMR resonance is coupled to $C\alpha$ –F: $J_{FF} = 11$ Hz). This is in contrast to previous suggestions of a metal fluorocarbene intermediate (Scheme 5).⁹

Having established that the NHC ligand remains bound to the metal upon fluoride-abstraction from 3, we shifted our focus to C–F bond functionalization using Brønsted acids. Treatment of 3 with trifluoroacetic acid [TFA; $pK_a = 3.4$ (DMSO)]^{28a} gives HF and the more stable (*vs.* 5a) trifluoroacetate-substituted metallacycle 5b that could be characterized spectroscopically at room temperature (Scheme 6). Nonetheless, accompanying formation of perfluoro-cyclobutene, presumably formed *via* an analogous structure to 4a, led us to move to weaker Brønsted acids. Remarkably, reaction of 3 with acetic acid [$pK_a = 12$ (DMSO)]^{28b} (Scheme 6, bottom) yielded the stable ester metallacycle 5c (30% isolated yield) as well as the Ni–C^F bond cleavage product 6a in a 1 : 1 ratio. At a similar acidity [$pK_a = 11$ (DMSO)]^{28c} but increased steric bulk, 2,4,6-trimethyl-benzoic acid gave a 10 : 1 mixture favouring the ring cleavage product, 6b.

The molecular structure of 5c features similar Ni–C bond distances (1.895(7) *vs.* 1.896(6) Å) and a distorted square planar coordination (Fig. 4). The functionalized heptafluoro-metallacyclopentane ring is puckered with the smallest C–C bond distance being $C(32)\alpha$ – $C(39)\beta$ [1.49(1) Å]. The carbonyl oxygen completes the nickel coordination sphere [Ni–O1 = 1.969(4) Å].

The ¹⁹F NMR spectra of 5a–d are very similar and support our original proposal for the low temperature intermediate 5a in the





Fig. 3 ORTEP representation of the molecular structure of 4a with thermal ellipsoid probabilities set to 30% and hydrogen atoms omitted for clarity. The Ni–C(1) distance is 1.854(2) Å.



Scheme 4 Synthesis and decomposition of 4a.



Scheme 5 Possible intermediates in the reaction of 3 with TMSOTf.







Fig. 4 ORTEP representation of the molecular structure of **5c** with thermal ellipsoid probabilities set to 30% and hydrogen atoms omitted for clarity. The Ni-C(1) distance is 1.928(2) Å.

reaction of **3** with TMSOTf. The $\text{C}\alpha\text{-F}$ ^{19}F chemical shifts of the functionalized carbon, (-117.2 and -119.2 ppm) can be compared with those of the phosphonium analogs (-115.6 and -117.8 ppm) shown in Scheme 2.


As expected, the ring-opened products **6a** and **b** display nearly identical ^{19}F NMR chemical shift patterns with the $\text{C}\gamma\text{-}$ and $\text{C}\delta\text{-F}$ resonances distinguished by F-H coupling of **6** and 52 Hz, respectively. These unique complexes have been identified as their potassium cation adducts using ESI-MS (747.2 g mol^{-1} and 851.4 g mol^{-1} respectively) and are surprisingly inert to thermolysis at 80°C in C_6D_6 for 20 h.

Considering the importance of esters as synthons in organic transformations²⁹ this C–O bond-forming reaction **3** \rightarrow **5** is very appealing from the standpoint of synthesizing functionalized fluorocarbons. As such, understanding competing pathways for M–C vs. C–F bond cleavage would be valuable.³⁰ Viable reaction pathways can be considered as proceeding *via* either 5- or 6-membered transition states (Scheme 7). The selective HF elimination observed for TFA, is eroded as Ni–C bond protonolysis (orbital control) competes using acids of intermediate acidity (e.g. $\text{p}K_a \sim 11$).³¹ With the bulkier trimethylbenzoic acid, kinetic acidity factors in the tighter 5-membered ring transition state could severely limit HF elimination.³²




Scheme 6 Reaction of **3** with trifluoroacetic, acetic and 2,4,6-trimethylbenzoic acids.

1) C–F bond protonolysis



5-membered concerted TS; favoured by stronger acid (TFA)

2) Metal-alkyl protonolysis



6-membered concerted TS (S_E2); favoured with bulky weaker acid

Scheme 7 Proposed reaction pathways for C–F activation vs. Ni–R^F protonolysis.

Conclusions

In summary, we have prepared the first NHC-perfluorometallacyclopentane complexes and exploited the bulky SiPr ligand to stabilize a pseudo-three-coordinate nickelacycle, **3**. Importantly, **3** undergoes $\text{C}\alpha\text{-F}$ abstraction reactions without migration of the NHC ligand. Instead, we see an unprecedented migration of the fluoroalkyl to the reactive carbon center, giving rise to the novel perfluorocyclobutyl complex *via* Ni–C^F bond cleavage. More importantly, the low-coordinate nature of **3** allows for ring functionalization. Strong acids favour selective $\text{C}\alpha$ functionalization, but the resulting products are unstable with respect to competing metallacycle ring contraction and elimination of perfluorocyclobutene. With less acidic reagents stable ring-functionalized products are formed but a competing Ni–C^F bond cleavage pathway comes into play and dominates for bulkier carboxylic acids. These are the first examples of selective functionalization of a PNCP and synthesis of thermally stable Ni–C₄F₈H complexes. These results are encouraging in the context of developing metallacycle-based routes to functionalized fluorocarbons.

Ongoing work is focused on (a) expanding the scope of ring functionalization substrates suitable for reactions with **3** and (b) reductive (see Scheme 2, above) and oxidative approaches for removing the functionalized fluorocarbon fragments from the metal. Preliminary results of the hydrogenolysis of compound **3** indicate enhanced reactivity towards H_2 (*i.e.*, at 7 psig and 25°C) *vs.* reported 4-coordinate phosphite variants.^{11a} However, loss of selectivity³³ is observed with the synthesis of two distinct products. Full details of these results will be published in due time.

Acknowledgements

We thank the NSERC and the Canada Research Chairs program for generous financial support and the University of Ottawa, Foundation for Innovation and Ontario Ministry of Economic Development and Innovation for essential infrastructure. N. O. A. gratefully acknowledges support from the province of Ontario, NSERC and the University of Ottawa (OGS and CGS-M/D).



Notes and references

1 (a) R. D. Chambers, *Fluorine in Organic Chemistry*, Blackwell, Oxford, 2004; (b) P. Kirsch, *Modern Fluoroorganic Chemistry: Synthesis, Reactivity, Applications*, Wiley-VCH, Weinheim, 2 edn, 2013; (c) S. Purser, P. R. Moore, S. Swallow and V. Gouverneur, *Chem. Soc. Rev.*, 2008, **37**, 320; (d) I. Ojima, *Fluorine in Medicinal Chemistry and Chemical Biology*, John Wiley & Sons, Ltd, Chichester, 2009.

2 Stoichiometric $[\text{Cu}]$ - CF_3 reagents for trifluoromethylation of organic substrates, particularly aryl halides: (a) H. Morimoto, T. Tsubogo, N. D. Litvinas and J. F. Hartwig, *Angew. Chem., Int. Ed.*, 2011, **50**, 3793; (b) G. G. Dubinina, H. Furutachi and D. A. Vicic, *J. Am. Chem. Soc.*, 2008, **130**, 8600; (c) D. M. Wiemers and D. J. Burton, *J. Am. Chem. Soc.*, 1986, **108**, 832.

3 Metal-catalyzed C–F and C– CF_3 bond formation: (a) T. Furuya, A. S. Kamlet and T. Ritter, *Nature*, 2011, **473**, 470; (b) E. J. Cho, T. D. Senecal, T. Kinzel, Y. Zhang, D. A. Watson and S. L. Buchwald, *Science*, 2010, **328**, 1678; (c) V. V. Grushin, *Acc. Chem. Res.*, 2010, **43**, 160; (d) J. M. Brown and V. Gouverneur, *Angew. Chem., Int. Ed.*, 2009, **48**, 8610; (e) K. L. Hull, W. Q. Anani and M. S. Sanford, *J. Am. Chem. Soc.*, 2006, **128**, 7134.

4 Non-metal based approaches to introducing fluorocarbon fragments to organic substrates: (a) F. Wang, T. Luo, J.-B. Hu, Y. Wang, H. S. Krishnan, P. V. Jog, S. K. Ganesh, G. K. S. Prakash and G. A. Olah, *Angew. Chem., Int. Ed.*, 2011, **50**, 7153 (CF_2 addition to alkenes); (b) R. P. Singh, G. Cao, R. L. Kirchmeier and J. M. Shreeve, *J. Org. Chem.*, 1999, **64**, 2873. (Me_3Si - CF_3 addition to carbonyl groups).

5 (a) H.-J. Arpe, *Industrial Organic Chemistry*, Wiley-VCH, Weinheim, 5th edn, 2010; (b) *Applied Homogeneous Catalysis with Organometallic Compounds*, ed. B. Cornils and W. A. Herrmann, Wiley-VCH, New York, 1996.

6 (a) F. L. Taw, A. E. Clark, A. H. Mueller, M. T. Janicke, T. Cantat, B. L. Scott, P. J. Hay, R. P. Hughes and J. L. Kiplinger, *Organometallics*, 2012, **31**, 1484; (b) R. P. Hughes, *Adv. Organomet. Chem.*, 1990, **31**, 183; (c) J. A. Morrison, *Advances in Organometallic Chemistry: Trifluoromethyl-Containing Transition Metal Complexes*, Academic Press, 1993, vol. 35; (d) P. J. Brothers and W. R. Roper, *Chem. Rev.*, 1988, **88**, 1293.

7 $[\text{Mn}]$ - CF_3 inert to CO migratory-insertion: S. K. Shin and J. L. Beauchamps, *J. Am. Chem. Soc.*, 1990, **112**, 2057.

8 (a) M. Ohashi, M. Shibata, H. Saito, T. Kambara and S. Ogoshi, *Organometallics*, 2013, **32**, 3631; (b) W. Gasafit-Martin, G. Oberendfellner and K. von Werner, *Can. J. Chem.*, 1996, **74**, 1922; (c) M. Green, A. Laguna, J. L. Spencer and F. G. A. Stone, *J. Chem. Soc., Dalton Trans.*, 1977, 1010; (d) J. Browning, H. D. Empsall, M. Green and F. G. A. Stone, *J. Chem. Soc., Dalton Trans.*, 1973, 381; (e) C. S. Cundy, M. Green and F. G. A. Stone, *J. Chem. Soc. A*, 1970, 1647; (f) M. Green, R. B. L. Osborn, A. J. Rest and F. G. A. Stone, *J. Chem. Soc. A*, 1968, 2525; (g) P. M. Treichel and F. G. A. Stone, *Adv. Organomet. Chem.*, 1964, **1**, 143.

9 R. R. Burch, J. C. Calabrese and S. D. Ittel, *Organometallics*, 1988, **7**, 1642.

10 K. A. Giffin, D. J. Harrison, I. Korobkov and R. T. Baker, *Organometallics*, 2013, **32**, 7424.

11 (a) R. T. Baker, R. P. Beatty, W. B. Farnham and R. L. Wallace Jr, *US Pat.*, 5670679, 1997; (b) Generation of perfluorocyclobutene or perfluorocyclobutane from an iron perfluorometallacyclopentane complex, under thermal or thermal/oxidizing(Br_2) conditions, respectively, T. A. Manuel, S. L. Stafford and F. G. A. Stone, *J. Am. Chem. Soc.*, 1961, **83**, 249.

12 M. Ohashi and S. Ogoshi, Catalytic Transformations of Fluorinated Olefins. in *Topics in Organometallic Chemistry*, Springer, Switzerland, 2014.

13 Activation of C–F bonds in $[\text{M}]$ - R^F complexes: (a) R. P. Hughes, *J. Fluorine Chem.*, 2010, **131**, 1059; (b) J. Goodman, V. V. Grushin, R. B. Larichev, S. A. Macgregor, W. J. Marshall and D. C. Roe, *J. Am. Chem. Soc.*, 2009, **131**, 4236; (c) S. A. Garratt, R. P. Hughes, I. Kovacik, A. J. Ward, S. Willemsen and D. Zhang, *J. Am. Chem. Soc.*, 2005, **127**, 15585; (d) H. Torrens, *Coord. Chem. Rev.*, 2005, **249**, 1957; (e) D. Huang, P. R. Koren, K. Folting, E. R. Davidson and K. G. Caulton, *J. Am. Chem. Soc.*, 2000, **122**, 8916; (f) T. G. Richmond and D. F. Shriver, *Organometallics*, 1984, **3**, 314; (g) T. G. Richmond and D. F. Shriver, *Organometallics*, 1983, **2**, 1061, see also ref. 8a.

14 (a) R. H. Crabtree, *J. Organomet. Chem.*, 2005, **690**, 5451; (b) D. J. D. Wilson, S. A. Couchman and J. L. Dutton, *Inorg. Chem.*, 2012, **51**, 7657.

15 D. J. Nelson and S. P. Nolan, *Chem. Soc. Rev.*, 2013, **42**, 6723.

16 C. A. Laskowsky, A. J. Miller, G. L. Hillhouse and T. R. Cundari, *J. Am. Chem. Soc.*, 2011, **133**, 771.

17 S. Miyazaki, Y. Koga, T. Matsumoto and K. Matsubara, *Chem. Commun.*, 2010, **46**, 1932.

18 P. S. Hanley, S. L. Marquard, T. R. Cundari and J. F. Hartwig, *J. Am. Chem. Soc.*, 2012, **134**, 15281.

19 Ni–RF bond lengths for structure 2 (see ESI†): 1.956(1) Å (trans to P); 1.945(1) Å (trans to C). Ni–C(NHC) bond length for 2: 1.945(1) Å.

20 See ESI page S14 for low temperature NMR spectra.†

21 S. Grimme, *J. Comput. Chem.*, 2006, **27**, 1787.

22 See ESI page S29.†

23 N. Islam and D. C. Gosh, *Theoretical and Computational Research in the 21st century*, ed. N. Islam, CRC Press, Boca Raton, 2014, ch. 1, pp. 1–58.

24 To the best of our knowledge, the only previously reported metal perfluorocyclobutyl complexes are: (a) B. L. Dyatkin, B. I. Martynov, L. G. Martynova, N. G. Kizim, S. R. Sterlin, Z. A. Stumbrevichute and L. A. Federov, *J. Organomet. Chem.*, 1973, **57**, 423; (b) B. L. Dyatkin, S. R. Sterlin, B. I. Martynov, E. I. Mysov and I. L. Knunyants, *Tetrahedron*, 1971, **27**, 2843.

25 (a) The leaving group ability of OTf, R. K. Crossland, W. E. Wells and V. J. Shiner, *J. Am. Chem. Soc.*, 1971, **93**, 4217.



26 (a) T. Miura, Y. Ito and M. Murakami, *Chem. Lett.*, 2008, **37**, 1006; (b) C. J. Bourgeois, R. P. Hughes, J. Yuan, A. G. DiPasquale and A. L. Rheingold, *Organometallics*, 2006, **25**, 2908.

27 For NMR spectra of compound **4a** see pages S21–S22.†

28 (a) M. H. Haindl, M. B. Schmid, K. Zeitler and R. M. Gschwind, *RSC Adv.*, 2012, **2**, 5941; (b) T. Cohen, D. A. Bennett and A. J. Mura Jr, *J. Org. Chem.*, 1976, **41**, 2507; (c) J. Jover, R. Bosque and J. Sales, *QSAR Comb. Sci.*, 2008, **27**, 563.

29 W. Riemenschneider and H. M. Bolt, Esters, Organic, in *Ullman's Encyclopedia of Industrial Chemistry*, Wiley-VCH, Weinheim, 2005.

30 Previous work showed protonation of C–F bond instead of Ir–CH₃ using a strong acid: R. P. Hughes, D. Zhang, L. N. Zakharov and A. L. Rheingold, *Organometallics*, 2002, **21**, 4902.

31 E. W. Kalberer, J. F. Houlis and D. M. Roddick, *Organometallics*, 2004, **23**, 4112–4115.

32 Similar reactivity was previously observed for 3-membered perfluorinated metallacycles: W. Xu, H. Sun, Z. Xiong and X. Li, *Organometallics*, 2013, **32**, 7122.

33 R. P. Hughes and J. M. Smith, *J. Am. Chem. Soc.*, 1999, **121**, 6084.

