

Chemical Science

www.rsc.org/chemicalscience

The Royal Society of Chemistry is the world's leading chemistry community. Through our high impact journals and publications we connect the world with the chemical sciences and invest the profits back into the chemistry community.

IN THIS ISSUE

ISSN 2041-6539 CODEN CSHCBM 6(3) 1593–2124 (2015)



Cover

See Bo Wang, Ying-Wei Yang et al., pp. 1640–1644.
Image reproduced by permission of Ying-Wei Yang from *Chem. Sci.*, 2015, 6, 1640.



Inside cover

See Stuart L. James et al., pp. 1645–1649.
Image reproduced by permission of Stuart L. James from *Chem. Sci.*, 2015, 6, 1645.

PERSPECTIVES

1614

Experiences with a researcher-centric ELN

K. A. Badiola, C. Bird, W. S. Brocklesby, J. Casson, R. T. Chapman, S. J. Coles, J. R. Cronshaw, A. Fisher, J. G. Frey,* D. Gloria, M. C. Grossel, D. B. Hibbert, N. Knight, L. K. Mapp, L. Marazzi, B. Matthews, A. Milsted, R. S. Minns, K. T. Mueller, K. Murphy, T. Parkinson, R. Quinnell, J. S. Robinson, M. N. Robertson, M. Robins, E. Springate, G. Tizzard, M. H. Todd, A. E. Williamson, C. Willoughby, E. Yang and P. M. Ylioja

Chemical scientists' experiences with LabTrove.

LabTrove




1630

Ion-mediated conformational switches

Peter C. Knipe, Sam Thompson* and Andrew D. Hamilton*

The ability to control the conformation of a single molecule in a reversible and stimulus dependent manner holds great promise for fields as disparate as drug delivery and molecular electronics. Here we offer a perspective on recent developments in ion-mediated switching architectures and their ability to perform in a range of settings.



Editorial staff**Interim executive editor**

May Copsey

Deputy editor

Jeanne Andres

Editorial production manager

Philippa Ross

Development editors

Alessia Millermaggi

Cesar Palmero

Publishing editors

Matthew Bown, Sage Bowser, Hugh Cowley,
 Ruth Dilleen, Cally Haynes, Alan Holder,
 Samantha Ivel, James Moore, Liisa Niitsoo,
 Victoria Richards, Susan Weatherby, Rachel Wood

Publishing assistants

Natalie Ford, Bethany Johnson, Rebecca Wojturska

Publisher

Jamie Humphrey

For queries about submitted articles please contact
 Philippa Ross, Editorial production manager, in the
 first instance. E-mail chemicalscience@rsc.org

For pre-submission queries please contact
 May Copsey, Interim executive editor.
 E-mail chemicalscience-rsc@rsc.org

Chemical Science (electronic: ISSN 2041-6539)
 is published monthly by the Royal Society of
 Chemistry, Thomas Graham House, Science Park,
 Milton Road, Cambridge, CB4 0WF, UK.

Chemical Science is a Gold Open Access journal
 and all articles from 2015 onwards are free to
 read. Please email orders@rsc.org to register
 your interest or contact RSC Order Department,
 Royal Society of Chemistry, Thomas Graham
 House, Science Park, Milton Road, Cambridge,
 CB4 0WF, UK

Tel +44 (0)1223 432398; E-mail orders@rsc.org

Advertisement sales: Tel +44 (0) 1223 432246;
 Fax +44 (0) 1223 426017; E-mail advertising@rsc.org

For marketing opportunities relating to this journal,
 contact marketing@rsc.org

Chemical Science

www.rsc.org/chemicalscience**Editorial board****Editor-in-chief**

Daniel G. Nocera, Harvard University

Associate editors

Zhenan Bao, Stanford University

Christopher C. Cummins,
 Massachusetts Institute of
 TechnologyKazunari Domen, University of Tokyo
 Vy Dong, University of California,
 IrvineMatthew Gaunt, University of
 CambridgeHubert Girault, Federal Polytechnic
 School of LausanneChris Hunter, University of
 CambridgeDavid A. Leigh, University of
 ManchesterKopin Liu, Academia Sinica
 Wonwoo Nam, Ewha Womans
 UniversityF. Dean Toste, University of California,
 Berkeley

Haw Yang, Princeton University

Jihong Yu, Jilin University

Advisory board

Takuzo Aida, University of Tokyo

Markus Antonietti, Max Planck

Institute of Colloids and Interfaces

Polly Arnold, University of Edinburgh

Xinhe Bao, Dalian Institute of
 Chemical Physics

Guy Bertrand, University of California,

Los Angeles

Jeffrey Bode, Swiss Federal Institute of
 Technology Zurich

Christopher Chang, University of

California, Berkeley

Chi-Ming Che, University of Hong
 Kong

Jason Chin, Medical Research Council

Laboratory of Molecular Biology

Daniel Chiu, University of Washington

Graham Cooks, Purdue University

Eugenio Coronado, University of

Valencia

Lee Cronin, University of Glasgow

Gautam R. Desiraju, Indian Institute of

Science, Bangalore

James Durrant, Imperial College

London

Ben Feringa, University of Groningen

Cynthia Friend, Harvard University

Makoto Fujita, University of Tokyo

Philip Gale, University of Southampton

Song Gao, Peking University

Jinlong Gong, Tianjin University

Justin Gooding, University of New

South Wales

Michael Graetzel, Federal Polytechnic

School of Lausanne

Duncan Graham, University of

Strathclyde

Buxing Han, Chinese Academy of

Sciences

Jeremy Harvey, University of Bristol

Christy Haynes, University of

Minnesota

Johan Hofkens, Catholic University

of Leuven

Linda Hsieh-Wilson, California

Institute of Technology

Eric Jacobsen, Harvard University

Takashi Kato, University of Tokyo

Seong Keun Kim, Seoul National

University

Jerome Lacour, University of Geneva

James Leighton, Columbia University

Steve Ley, University of Cambridge

Chao-Jun Li, McGill University

Wenbin Lin, University of North

Carolina

Watson Loh, Instituto de Química

Julie Macpherson, University of

Warwick

Stephen Mann, University of Bristol

James McCusker, Michigan State

University

Bert Meijer, Eindhoven University of

Technology

Nils Metzler-Nolte, Ruhr University

Bochum

Scott Miller, Yale University

Daniel Mindiola, Indiana University

Mohammad Movassagh,

Massachusetts Institute of

Technology

Jonathan Nitschke, University of

Cambridge

Kyoko Nozaki, University of Tokyo

Takashi Ooi, Nagoya University

Rachel O'Reilly, University of Warwick

Michel Orrit, Leiden University

Oleg Ozerov, Texas A&M University

Hongkun Park, Harvard University

Rasmita Raval, University of Liverpool

Paul Reider, Princeton University

Stuart Rowan, Case Western Reserve

University

Richmond Sarpong, University of

California, Berkeley

Gregory Scholes, University of

Toronto

Oliver Seitz, Humboldt University of

Berlin

Kay Severin, Federal Polytechnic

School of Lausanne

Mikiko Sodeoka, RIKEN

Brian Stoltz, California Institute of

Technology

Weihong Tan, University of Florida

He Tian, East China University of

Science and Technology

Zhong-Qun Tian, Xiamen University

Andrei Tokmakoff, University of

Chicago

Jan Van Hest, Radboud University

Tom Welton, Imperial College London

Christina White, University of Illinois

Martin Wolf, Fritz Haber Institute of

the Max Planck Society

Omar Yaghi, University of California,

Los Angeles

Vivian Yann, University of Hong Kong

Yang Yang, University of California,

Los Angeles

Shu-Hong Yu, University of Science

and Technology of China

Qi-Lin Zhou, Nankai University

Information for authors

Full details on how to submit material for publication in Chemical Science are given in the Instructions for Authors (available from <http://www.rsc.org/authors>). Submissions should be made via the journal's homepage: <http://www.rsc.org/chemicalscience>.

Authors may reproduce/republish portions of their published contribution without seeking permission from the RSC, provided that any such republication is accompanied by an acknowledgement in the form: (Original Citation)–Reproduced by permission of The Royal Society of Chemistry.

This journal is ©The Royal Society of Chemistry 2015. Apart from fair dealing for the purposes of research or private study for non-commercial purposes, or criticism or review, as permitted under the Copyright, Designs and

Patents Act 1988 and the Copyright and Related Rights Regulation 2003, this publication may only be reproduced, stored or transmitted, in any form or by any means, with the prior permission in writing of the Publishers or in the case of reprographic reproduction in accordance with the terms of licences issued by the Copyright Licensing Agency in the UK. US copyright law is applicable to users in the USA.

The Royal Society of Chemistry takes reasonable care in the preparation of this publication but does not accept liability for the consequences of any errors or omissions.

© The paper used in this publication meets the requirements of ANSI/NISO Z3948–1992 (Permanence of Paper). Registered Charity No. 207890.




EDGE ARTICLES

1640

Stimuli-responsive metal–organic frameworks gated by pillar[5]arene supramolecular switches

Li-Li Tan, Haiwei Li, Yu-Chen Qiu, Dai-Xiong Chen, Xin Wang, Rui-Yi Pan, Yan Wang, Sean Xiao-An Zhang, Bo Wang* and Ying-Wei Yang*

Mechanized monodisperse nano metal–organic frameworks (NMOFs) gated by carboxylatopillar[5]arene (CP5) switches with bio-friendly pH-triggered release capabilities were constructed for the first time as a new stimuli-responsive theranostic hybrid platform.




1645

Synthesis by extrusion: continuous, large-scale preparation of MOFs using little or no solvent

Deborah Crawford, José Casaban, Robert Haydon, Nicola Giri, Tony McNally and Stuart L. James*

Continuous flow mechanochemical and melt-phase synthesis at kg h^{-1} rates from solid reagents and either no solvent, or only minimal solvent, is reported.




1650

Integration of an anti-tumor drug into nanocrystalline assemblies for sustained drug release

Xiangrui Yang, Shichao Wu, Yang Li, Yu Huang, Jinyan Lin, Di Chang, Shefang Ye, Liya Xie, Yuan Jiang* and Zhenqing Hou*

A bio-inspired approach was used to integrate an anti-tumor drug into nanocrystalline assemblies for sustained drug release.




1655

Critical analysis of the limitations of Bleaney's theory of magnetic anisotropy in paramagnetic lanthanide coordination complexes


Alexander M. Funk, Katie-Louise N. A. Finney, Peter Harvey, Alan M. Kenwright, Emily R. Neil, Nicola J. Rogers, P. Kanthi Senanayake and David Parker*

The origins of the breakdown of Bleaney's theory of magnetic anisotropy are described, based on an analysis of eleven different complexes of the second half of the 4f elements that form isostructural series.



EDGE ARTICLES

1663




An ambient stable core-substituted perylene bisimide dianion: isolation and single crystal structure analysis

Sabine Seifert, David Schmidt and Frank Würthner*

The reduction of a highly electron deficient PBI afforded the corresponding dianion disodium salt that was characterized by single crystal structure analysis.

1668




Mesoporous inorganic salts with crystal defects: unusual catalysts and catalyst supports

Xinchen Kang, Wenting Shang, Qinggong Zhu, Jianling Zhang, Tao Jiang,* Buxing Han,* Zhonghua Wu, Zhihong Li and Xueqing Xing

Mesoporous LaF_3 , NdF_3 , and YF_3 particles with crystal defects, which are excellent catalysts and catalyst supports, have been synthesized successfully.

1676




Metal oxidation states in biological water splitting

Vera Krewald, Marius Retegan, Nicholas Cox, Johannes Messinger, Wolfgang Lubitz, Serena DeBeer, Frank Neese and Dimitrios A. Pantazis*

A central question in biological water splitting concerns the oxidation states of the manganese ions that comprise the oxygen-evolving complex of photosystem II.

1696



5,20-Bis(α -oligothienyl)-substituted [26]hexaphyrins possessing electronic circuits strongly perturbed by meso-oligothienyl substituents

Hirotaka Mori, Masaaki Suzuki, Woojae Kim, Jong Min Lim, Dongho Kim* and Atsuhiro Osuka*

A series of [26]hexaphyrins(1.1.1.1.1) bearing two α -oligothienyl substituents at 5,20-positions have been synthesised and are shown to have a dumbbell hexaphyrin conformation, to which the α -oligothienyl groups are linked to form an acyclic helix-like conjugated network.




EDGE ARTICLES

1701

Photostick: a method for selective isolation of target cells from culture

Miao-Ping Chien, Christopher A. Werley, Samouil L. Farhi and Adam E. Cohen*


A new method enables optical selection of one or more cells in culture *via* photochemical crosslinking to the culture dish.

1706

Inherently chiral electrodes: the tool for chiral voltammetry

Serena Arnaboldi,* Tiziana Benincori, Roberto Cirilli, Włodzimierz Kutner, Mirko Magni, Patrizia Romana Mussini,* Krzysztof Noworyta and Francesco Sannicolò*


Artificial inherently chiral electrode surfaces perform clear enantiodiscrimination and enantiomeric ratio quantification, opening the way to chiral voltammetry.



1712

Emptying and filling a tunnel bronze


Peter M. Marley, Tesfaye A. Abtew, Katie E. Farley, Gregory A. Horrocks, Robert V. Dennis, Peihong Zhang and Sarbajit Banerjee*

We report the synthesis of a new tunnel-structured polymorph of V_2O_5 (ζ - V_2O_5) synthesized by topotactic ion extraction.

1719


Synergistic cascade catalysis by metal nanoparticles and Lewis acids in hydrogen autotransfer

Gerald C. Y. Choo, Hiroyuki Miyamura and Shū Kobayashi*

Synergistic cascade catalysis of Au/Pd nanoparticles/Lewis acids achieved *N*-alkylation of primary amides to secondary amides with alcohols *via* hydrogen autotransfer.

EDGE ARTICLES


1728

**CIT-7, a crystalline, molecular sieve with pores bounded by 8 and 10-membered rings**

Joel E. Schmidt, Dan Xie, Thomas Rea and Mark E. Davis*

CIT-7, a new microporous material that contains 8 and 10-membered rings prepared as a silicate, an aluminosilicate or titanosilicate.


1735

**Mechanism, reactivity, and selectivity of the iridium-catalyzed C(sp³)–H borylation of chlorosilanes**

Genping Huang, Marcin Kalek, Rong-Zhen Liao and Fahmi Himo*

DFT calculations are used to elucidate the reaction mechanism, the role of the chlorosilyl group, and primary vs. secondary and C(sp³)–H vs. C(sp²)–H selectivity of the iridium-catalyzed borylation of chlorosilanes.


1747

**Design of an intelligent sub-50 nm nuclear-targeting nanotheranostic system for imaging guided intranuclear radiosensitization**

Wenpei Fan, Bo Shen, Wenbo Bu,* Xiangpeng Zheng, Qianjun He, Zhaowen Cui, Kuaile Zhao, Shengjian Zhang and Jianlin Shi*

MR/UCL bimodal imaging guided intranuclear radiosensitization was firstly achieved by an intelligent nuclear-targeting nanotheranostic system.

1754

**A facile route to electronically conductive polyelectrolyte brushes as platforms of molecular wires**

Karol Wolski, Michał Szuwarzyński and Szczeban Zapotoczny*

Conjugated polyelectrolyte brushes grafted from a conductive surface and forming a 1D macromolecular pathway for charge transport are synthesized.




EDGE ARTICLES

1761

Catalysis of water oxidation in acetonitrile by iridium oxide nanoparticles

Jonnathan C. Hidalgo-Acosta, Manuel A. Méndez, Micheál D. Scanlon, Heron Vrubel, Véronique Amstutz, Wojciech Adamiak, Marcin Opallo and Hubert H. Girault*

Micro-domains of water molecules surrounded by organic solvent exhibit enhanced reactivity towards oxidation compared to highly hydrogen-bonded bulk water molecules.




1770

Protein ubiquitination and formation of polyubiquitin chains without ATP, E1 and E2 enzymes

Sungjin Park, David T. Krist and Alexander V. Statsyuk*

Protein ubiquitination without ATP. This paper reports a chemical strategy to ubiquitinate proteins without ATP, E1, and E2 enzymes, offering new insights on the biochemical mechanism of E3s.




1780

MM quadruply bonded complexes supported by vinylbenzoate ligands: synthesis, characterization, photophysical properties and application as synthons

Samantha E. Brown-Xu, Malcolm H. Chisholm,* Christopher B. Durr, Thomas F. Spilker and Philip J. Young

MM complexes as potential synthons for the development of higher order extended structures via Heck coupling reactions, exhibiting interesting photophysical properties.




1792

Discovery of potent inhibitors of human β -tryptase from pre-equilibrated dynamic combinatorial libraries


Qian-Qian Jiang, Wilhelm Sicking, Martin Ehlers and Carsten Schmuck*

Pre-equilibrated combinatorial libraries based on multivalent peptide acyl hydrazones were used to find potent inhibitors of β -tryptase. The best inhibitors bind to the protein surface, and inhibit β -tryptase with nanomolar affinity (K_i ca. 10 nM) and high selectivity in a reversible and non-competitive way.



EDGE ARTICLES

1801




A faux hawk fullerene with PCBM-like properties

Long K. San, Eric V. Bukovsky, Bryon W. Larson, James B. Whitaker, S. H. M. Deng, Nikos Kopidakis,* Garry Rumbles,* Alexey A. Popov,* Yu-Sheng Chen,* Xue-Bin Wang,* Olga V. Boltalina* and Steven H. Strauss*

A fluorinated faux hawk fullerene with comparable OPV-relevant TRMC performance and far greater thermal stability than PCBM is reported.

1816




C–H arylation of triphenylene, naphthalene and related arenes using Pd/C

Karl D. Collins, Roman Honeker, Suhelen Vásquez-Céspedes, Dan-Tam D. Tang and Frank Glorius*

A highly selective arylation of a number of polyaromatic hydrocarbons (PAHs) with aryliodonium salts and Pd/C as the only reagent is reported.

1825




Fluorescent/phosphorescent dual-emissive conjugated polymer dots for hypoxia bioimaging

Qiang Zhao,* Xiaobo Zhou, Tianye Cao, Kenneth Yin Zhang, Lijuan Yang, Shujuan Liu, Hua Liang, Huiran Yang, Fuyou Li* and Wei Huang*

Fluorescent/phosphorescent dual-emissive conjugated polymer dots were designed and synthesized, which were used for tumor hypoxia sensing via ratiometric imaging and photoluminescence lifetime imaging.

1832



The energy blocker inside the power house: mitochondria targeted delivery of 3-bromopyruvate

Sean Marrache and Shanta Dhar*

Mitochondrial delivery of 3-bromopyruvate for metabolic reprogramming of cancer cells.




EDGE ARTICLES

1846

Strain discrimination of *Yersinia pestis* using a SERS-based electrochemically driven melting curve analysis of variable number tandem repeat sequences

E. Papadopoulou, N. Gale, S. A. Goodchild, D. W. Cleary, S. A. Weller, T. Brown and P. N. Bartlett*

Variable number tandem repeats in DNA extracted from three bacterial *Y. pestis* strains have been differentiated using *E*-melting analysis monitored by SERS, combined with the use of a blocker oligonucleotide.




1853

Functionalized graphene-based biomimetic microsensor interfacing with living cells to sensitively monitor nitric oxide release

Yan-Ling Liu, Xue-Ying Wang, Jia-Quan Xu, Chong Xiao, Yan-Hong Liu, Xin-Wei Zhang, Jun-Tao Liu and Wei-Hua Huang*

We present a biomimetic and reusable microsensor with sub-nanomolar sensitivity by elaborate functionalizing graphene for monitoring NO release in real-time.




1859

Dehydrogenation, disproportionation and transfer hydrogenation reactions of formic acid catalyzed by molybdenum hydride compounds

Michelle C. Neary and Gerard Parkin*

Cyclopentadienyl molybdenum hydride compounds are catalysts for dehydrogenation, disproportionation and transfer hydrogenation reactions of formic acid, in which the latter provides a means to reduce aldehydes and ketones.




1866

Highly sensitive and multiplexed analysis of CpG methylation at single-base resolution with ligation-based exponential amplification


Fengxia Su, Limei Wang, Yueying Sun, Chenghui Liu, Xinrui Duan and Zhengping Li*

Multiple CpG methylation can be accurately detected in one-tube ligase chain reaction (LCR) amplification with high sensitivity and specificity.



EDGE ARTICLES

1873




Well-defined BiOCl colloidal ultrathin nanosheets: synthesis, characterization, and application in photocatalytic aerobic oxidation of secondary amines

Yihui Wu, Bo Yuan, Mingrun Li, Wen-Hua Zhang,* Yan Liu* and Can Li*

Well-defined BiOCl ultrathin nanosheets were prepared by a facile colloidal route, and exhibit high photocatalytic performance toward the oxidation of secondary amines to corresponding imines under visible irradiation.

1879




A rationally designed small molecule for identifying an *in vivo* link between metal-amyloid- β complexes and the pathogenesis of Alzheimer's disease

Michael W. Beck, Shin Bi Oh, Richard A. Kerr, Hyuck Jin Lee, So Hee Kim, Sujeong Kim, Milim Jang, Brandon T. Ruotolo,* Joo-Yong Lee* and Mi Hee Lim*

An *in vivo* chemical tool designed to target metal- $\text{A}\beta$ complexes and modulate their activity was applied to the 5XFAD mouse model of Alzheimer's disease (AD) demonstrating the involvement of metal- $\text{A}\beta$ in AD pathology.

1887




Principal factors that determine the extension of detection range in molecular beacon aptamer/conjugated polyelectrolyte bioassays

Ji-Eun Jeong, Boram Kim, Shinjae Woo, Sungu Hwang, Guillermo C. Bazan* and Han Young Woo*

A new bioassay strategy based on the molecular beacon aptamer/conjugated polyelectrolyte demonstrates a fine-tuning of the detection range and limit of detection for weakly-binding targets.

1895



What can NMR spectroscopy of selenoureas and phosphinidenes teach us about the π -accepting abilities of *N*-heterocyclic carbenes?

Sai V. C. Vummala, David J. Nelson, Albert Poater, Adrián Gómez-Suárez, David B. Cordes, Alexandra M. Z. Slawin, Steven P. Nolan* and Luigi Cavallo*

The relationship between the NMR chemical shifts of phosphinidene and selenourea compounds and the π -accepting ability of the related carbene ligands has been investigated.




EDGE ARTICLES

1905

Increased carrier mobility in end-functionalized oligosilanes

S. Surampudi, M.-L. Yeh, M. A. Siegler, J. F. Martinez Hardigree, T. A. Kasl, H. E. Katz and R. S. Klausen*

A class of hybrid oligosilane–arene materials outperforms the σ - and π -conjugated parent structures.




1910

Helically structured metal–organic frameworks fabricated by using supramolecular assemblies as templates

Hui Wang, Wei Zhu, Jian Li, Tian Tian, Yue Lan, Ning Gao, Chen Wang, Meng Zhang, Charl F. J. Faul and Guangtao Li*

Here we report on the use of a self-assembled organic object to template the first example of a nanoscale metal–organic framework (MOF) with a helical morphology.




1917

Mechanistic analysis of an asymmetric palladium-catalyzed conjugate addition of arylboronic acids to β -substituted cyclic enones

Cornelia L. Boeser, Jeffrey C. Holder, Buck L. H. Taylor, K. N. Houk, Brian M. Stoltz and Richard N. Zare*

Mechanistic insight into a Pd-catalyzed conjugate addition reaction was gained using desorption electrospray ionization coupled to mass spectrometry.




1923

Rh(III)-catalyzed C–H olefination of *N*-pentafluoroaryl benzamides using air as the sole oxidant


Yi Lu,* Huai-Wei Wang, Jillian E. Spangler, Kai Chen, Pei-Pei Cui, Yue Zhao, Wei-Yin Sun* and Jin-Quan Yu*

Rhodium(III)-catalyzed C–H olefination reaction using air as the sole oxidant.



EDGE ARTICLES


1928

This work:**Cu-catalyzed transannulation reaction of pyridotriazoles with terminal alkynes under aerobic conditions: efficient synthesis of indolizines**

V. Helan, A. V. Gulevich and V. Gevorgyan*

A Cu(i)-catalyzed denitrogenative transannulation reaction of pyridotriazoles with terminal alkynes en route to indolizines was developed.


1932

**Tetraphenylpyrazine-based AIEgens: facile preparation and tunable light emission**

Ming Chen, Lingzhi Li, Han Nie, Jiaqi Tong, Lulin Yan, Bin Xu, Jing Zhi Sun, Wenjing Tian, Zujin Zhao, Anjun Qin* and Ben Zhong Tang*

Structurally new AIEgens with tunable emission based on tetraphenylpyrazine were rationally designed and readily prepared.


1938

**Magnetic MOF microreactors for recyclable size-selective biocatalysis**

Jia Huo, Jordi Aguilera-Sigalat, Samir El-Hankari and Darren Bradshaw*

Highly porous magnetic MOF microreactors can be prepared around a Pickering-stabilized hydrogel core, providing a facile means for the encapsulation of enzymes for size-selective biocatalysis.

1944

**A far-red emitting probe for unambiguous detection of mobile zinc in acidic vesicles and deep tissue**

Pablo Rivera-Fuentes, Alexandra T. Wrobel, Melissa L. Zastrow, Mustafa Khan, John Georgiou, Thomas T. Luyben, John C. Roder, Kenichi Okamoto and Stephen J. Lippard*

A fluorescent sensor that is pH-insensitive and compatible with two-photon microscopy was developed and applied to live cell and deep tissue imaging.




EDGE ARTICLES

1949

Stimuli-responsive hybrid materials: breathing in magnetic layered double hydroxides induced by a thermoresponsive molecule

Gonzalo Abellán, Jose Luis Jordá, Pedro Atienzar, María Varela, Miriam Jaafar, Julio Gómez-Herrero, Félix Zamora, Antonio Ribera, Hermenegildo García* and Eugenio Coronado*

A hybrid magnetic multilayer material consisting of CoAl-LDH ferromagnetic layers intercalated with thermoresponsive molecules diluted with a flexible surfactant has been obtained.




1959

Self-assembly of tetrareduced corannulene with mixed Li–Rb clusters: dynamic transformations, unique structures and record ^{7}Li NMR shifts

Alexander S. Filatov, Sarah N. Spisak, Alexander V. Zabula, James McNeely, Andrey Yu. Rogachev and Marina A. Petrukhina*

The record NMR shifts for the centrally encapsulated lithium ions are correlated with molecular and electronic structures of novel mixed alkali metal sandwiches.




1967

Air-stable ambipolar field-effect transistor based on a solution-processed octanaphthoxy-substituted tris(phthalocyaninato) europium semiconductor with high and balanced carrier mobilities

Xia Kong, Xia Zhang, Dameng Gao, Dongdong Qi, Yanli Chen* and Jianzhuang Jiang*

Simple solvent vapor annealing over QLS film-based OFET devices fabricated from (Pc)Eu[Pc(ONh)₈]Eu[Pc(ONh)₈] led to a high and balanced ambipolar performance.




1973

DNA-based visual majority logic gate with one-vote veto function


Daoqing Fan, Kun Wang, Jinbo Zhu, Yong Xia, Yanchao Han, Yaqing Liu* and Erkang Wang*

A label-free and enzyme-free three-input majority logic gate with one-vote veto function was developed for the first time.



EDGE ARTICLES


1979

**Chemiluminescent probes for imaging H_2S in living animals**

J. Cao, R. Lopez, J. M. Thacker, J. Y. Moon, C. Jiang, S. N. S. Morris, J. H. Bauer, P. Tao, R. P. Mason and A. R. Lippert*

Responsive 1,2-dioxetane chemiluminescent probes have been developed that display instantaneous, sensitive, and selective responses to H_2S and are capable of imaging H_2S in living mice.


1986

**Mesoporous silica nanoparticles for ^{19}F magnetic resonance imaging, fluorescence imaging, and drug delivery**

Tatsuya Nakamura, Fuminori Sugihara, Hisashi Matsushita, Yoshichika Yoshioka, Shin Mizukami and Kazuya Kikuchi*

We described perfluorocarbon encapsulated in mesoporous silica nanoparticles which enabled dual modal imaging (NIR/ ^{19}F MRI) and drug delivery.


1991

**Transformable H-bonds and conformation in compressed glucose**

Ewa Patyk and Andrzej Katrusiak*

Upon pressurisation above 5.4 GPa, α -D-glucose transforms into a new polymorph, with an altered molecular conformation and with intermolecular hydrogen bonds reshuffled.

1996

**Dynamic tracking of pathogenic receptor expression of live cells using pyrenyl glycoanthraquinone-decorated graphene electrodes**

Xiao-Peng He,* Bi-Wen Zhu, Yi Zang, Jia Li,* Guo-Rong Chen, He Tian and Yi-Tao Long*

Dynamic tracking of pathogenic receptor expression with live cells is made possible by pyrenyl glycoanthraquinones decorated on graphene electrodes.




EDGE ARTICLES

2002

A carbohydrate-grafted nanovesicle with activatable optical and acoustic contrasts for dual modality high performance tumor imaging

Xuanjun Wu, Bijuan Lin, Mingzhu Yu, Liu Yang, Jiahui Han and Shoufa Han*

High-performance illumination of subcutaneous tumor and liver tumor foci was achieved with sialic acid-targeted acid-responsive nanovesicles which become fluorescent and photoacoustic upon internalization into tumor lysosomes.




2010

A family of N-heterocyclic carbene-stabilized boreonium ions for metal-free imine hydrogenation catalysis

Jeffrey M. Farrell, Roy T. Posaratnanathan and Douglas W. Stephan*

Room-temperature metal-free hydrogenation catalysis.




2016

Electrophilic bis-fluorophosphonium dications: Lewis acid catalysts from diphosphines

Michael H. Holthausen, Rashi R. Hiranandani and Douglas W. Stephan*

A series of electrophilic bis-fluorophosphonium dications derived from diphosphines with naphthalene- and (oligo)-methylene-linkers is presented. The resulting Lewis acidity is demonstrated to depend on the spatial proximity between the P moieties as evidenced in several Lewis acid catalyzed transformations.




2022

The gold(i)…lead(ii) interaction: a relativistic connection


Raquel Echeverría, José M. López-de-Luzuriaga,* Miguel Monge and M. Elena Olmos

The heteronuclear complex $[\text{Pb}(\text{HB}(\text{pz})_3)\text{Au}(\text{C}_6\text{Cl}_5)_2]$ is a blue emitter in the solid state and displays an unprecedented and unsupported $\text{Au}(\text{i})\cdots\text{Pb}(\text{ii})$ contact. *Ab initio* calculations show a very strong metallophilic interaction that is greatly influenced by relativistic effects.

Unsupported $\text{Au}(\text{i})\cdots\text{Pb}(\text{ii})$ closed-shell interaction

EDGE ARTICLES


2027


Visible light photocatalytic reduction of aldehydes by Rh(III)-H: a detailed mechanistic study

T. Ghosh, T. Slanina and B. König*

The slow visible light mediated generation of a rhodium hydride allows the chemoselective reduction of aldehydes in the presence of ketones. Electron transfer from the chromophore to the metal complex proceeds via a radical anion intermediate or a solvated electron as two competing reaction pathways.


2035


Excited-state hydrogen atom abstraction initiates the photochemistry of β -2'-deoxycytidine

Rafał Szabla,* Jesús Campos, Judit E. Šponer, Jiří Šponer, Robert W. Góra* and John D. Sutherland*

H-D exchange experiments and quantum-chemical calculations elucidate the mechanism of photoinduced anomerisation and nucleobase loss reactions observed in β -2'-deoxycytidine.


2044


A heterobimetallic complex featuring a Ti-Co multiple bond and its application to the reductive coupling of ketones to alkenes

Bing Wu, Mark W. Bezpalko, Bruce M. Foxman and Christine M. Thomas*

A Ti/Co heterobimetallic complex featuring a very short metal–metal triple bond has been synthesized. This complex promotes the reductive coupling reaction of aryl ketones into alkenes.

2050


A noble metal-free proton-exchange membrane fuel cell based on bio-inspired molecular catalysts

P. D. Tran, A. Morozan, S. Archambault, J. Heidkamp, P. Chenevier, H. Dau, M. Fontecave, A. Martinent,* B. Jousselme* and V. Artero*

Bio-inspired chemistry allowed for the development of the first noble metal-free polymer electrolyte membrane hydrogen fuel cell (PEMFC). The device proved operational under technologically relevant conditions.




EDGE ARTICLES

2054

Bimetallic lanthanide complexes that display a ratiometric response to oxygen concentrations

Thomas Just Sørensen,* Alan M. Kenwright*
and Stephen Faulkner*

Heterometallic lanthanide complexes can display a ratiometric response to oxygen concentrations.




2060

Formation of a C–C double bond from two aliphatic carbons. Multiple C–H activations in an iridium pincer complex

Alexey V. Polukeev, Rocío Marcos, Mårten S. G. Ahlquist
and Ola F. Wendt*

Iridium can mediate a reversible intramolecular coupling reaction involving up to four unactivated C_{sp^3} -H bonds, to give a carbon-carbon double bond.




2068

Brushing up from “anywhere” under sunlight: a universal surface-initiated polymerization from polydopamine-coated surfaces

Wenbo Sheng, Bin Li, Xiaolong Wang, Bin Dai, Bo Yu,^{*}
Xin Jia^{*} and Feng Zhou^{*}

A generalized surface-initiated photografting procedure, which utilizes polydopamine as a photosensitive initiating layer, allows functionalization of almost any substrate with thin polymer films under sunlight.




2074

HMGB1 bound to cisplatin–DNA adducts undergoes extensive acetylation and phosphorylation *in vivo*


Yafeng He, Yin Ding, Dan Wang, Wanjun Zhang,
Weizhong Chen, Xichun Liu, Weijie Qin, Xiaohong Qian,
Hao Chen* and Zijian Guo*

Here, an application of a biomolecular probe based on a peptide–oligonucleotide conjugate is presented as a novel method for investigating the recognition of cisplatin–DNA lesions by HMGB1 *in vivo*.



EDGE ARTICLES


2079


Adaptive binding and selection of compressed 1,ω-diammonium-alkanes via molecular encapsulation in water

Dan Dumitrescu, Yves-Marie Legrand, Eddy Petit, Arie van der Lee and Mihail Barboiu*

Alkane chains may be encapsulated inside rigid crystalline capsules, adopting specific conformations of different levels of compression that are sufficiently kinetically stable under the confined conditions, to allow a conventional structure determination by X-ray diffraction.


2087


Interfacial chemical oxidative synthesis of multifunctional polyfluoranthene

Xin-Gui Li,* Yaozu Liao, Mei-Rong Huang* and Richard B. Kaner*

Polyfluoranthene, synthesized by an interfacial cationic oxidative polymerization of fluoranthene, emits visible fluorescence that shows highly selective sensitivity to Fe³⁺.


2102


Light-induced crosslinkable semiconducting polymer dots

Yue Zhang, Fangmao Ye, Wei Sun, Jiangbo Yu, I-Che Wu, Yu Rong, Yong Zhang and Daniel T. Chiu*

This paper describes photocrosslinkable Pdots with enhanced colloidal, physical, and chemical stability, and excellent encapsulating ability of functional small molecules.

2110


Semi-synthesis of a HGF/SF kringle one (K1) domain scaffold generates a potent *in vivo* MET receptor agonist

Claire Simonneau, Bérénice Leclercq, Alexandra Mougel, Eric Adriaenssens, Charlotte Paquet, Laurent Raibaut, Nathalie Ollivier, Hervé Drobecq, Julien Marcoux, Sarah Cianfranì, David Tulasne, Hugo de Jonge, Oleg Melnyk* and Jérôme Vicogne*

Clustering of synthetic biotinylated K1 domain from HGF/SF by streptavidin is sufficient to generate a potent MET agonist.



CORRECTION

2122

Correction: Utilizing the bioorthogonal base-pairing system of L-DNA to design ideal DNA nanocarriers for enhanced delivery of nucleic acid cargos

Kyoung-Ran Kim, Taemin Lee, Byeong-Su Kim and Dae-Ro Ahn*