

CrossMark  
click for updatesCite this: *Anal. Methods*, 2016, 8, 5881

DOI: 10.1039/c6ay90103d

www.rsc.org/methods

## Correction: Enhanced luminol–O<sub>2</sub> chemiluminescence reaction by CuO nanoparticles as oxidase mimics and its application for determination of ceftazidime

Mortaza Iranifam,<sup>a\*</sup> Amin Imani-Nabiyyi,<sup>b</sup> Alireza Khataee<sup>c</sup> and Jalil Kalantari<sup>a</sup>Correction for 'Enhanced luminol–O<sub>2</sub> chemiluminescence reaction by CuO nanoparticles as oxidase mimics and its application for determination of ceftazidime' by Mortaza Iranifam *et al.*, *Anal. Methods*, 2016, 8, 3816–3823.

In the original manuscript, there were errors in the captions of Fig. 5–8. An incorrect figure was also shown as Fig. 8. The corrected figures and accompanying captions are shown below.




Fig. 5 CL spectra of luminol/DO and luminol/DO/CuO NPs CL systems. Concentrations: luminol,  $5 \times 10^{-4}$  mol L<sup>-1</sup>; NaOH,  $3 \times 10^{-1}$  mol L<sup>-1</sup>; CuO NPs, 10 ppm.

<sup>a</sup>Department of Chemistry, Faculty of Science, University of Maragheh, 55181-83111 Maragheh, Iran. E-mail: MortezaIranifam@yahoo.com; Fax: +98 41 37276060; Tel: +98 91 44075996

<sup>b</sup>Department of Chemistry, Payame Noor University (PNU), Iran

<sup>c</sup>Research Laboratory of Advanced Water and Wastewater Treatment Processes, Department of Applied Chemistry, Faculty of Chemistry, University of Tabriz, 51666-16471 Tabriz, Iran






Fig. 6 Effect of concentration of NaOH on the analytical signal intensity ( $\Delta I$ ). Concentrations: luminol,  $8 \times 10^{-6}$  mol L<sup>-1</sup>; CuO NPs, 10 ppm; CFZM,  $3 \times 10^{-6}$  mol L<sup>-1</sup>.




Fig. 7 Effect of concentration of luminol on the analytical signal intensity ( $\Delta I$ ). Concentrations: NaOH,  $3 \times 10^{-1}$  mol L<sup>-1</sup>; CuO NPs, 10 ppm; CFZM,  $3 \times 10^{-6}$  mol L<sup>-1</sup>.




Fig. 8 Effect of concentration of CuO NPs on the analytical signal intensity ( $\Delta I$ ). Concentrations: luminol,  $5 \times 10^{-4}$  mol L<sup>-1</sup>; NaOH,  $3 \times 10^{-1}$  mol L<sup>-1</sup>; CFZM,  $3 \times 10^{-6}$  mol L<sup>-1</sup>.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

