

CrossMark
click for updatesCite this: *Anal. Methods*, 2016, 8, 5883

DOI: 10.1039/c6ay90104b

www.rsc.org/methods

Correction: Self-validated detection of hydroquinone in medicated cosmetic products using a preanodized screen-printed ring disk carbon electrode

Hsueh-Hui Yang,^{*a} Hsiao-Han Ting^b and Ying Shih^{*b}

Correction for 'Self-validated detection of hydroquinone in medicated cosmetic products using a preanodized screen-printed ring disk carbon electrode' by Hsueh-Hui Yang *et al.*, *Anal. Methods*, 2016, DOI: 10.1039/c6ay01000h.

The original manuscript contained errors in the y-axis of Fig. 6c. The corrected figure is shown below.




Fig. 6 Calibration curve for the HQ detection at a flow rate of 0.9 mL min⁻¹ with an applied potential of 0.15 V at the disk electrode (a–c) and –0.1 V at the ring electrode (A–C).

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aDepartment of Medical Research, Buddhist Tzu Chi General Hospital, Hualien 970, Taiwan. E-mail: hhyang@tzuchi.com.tw; Fax: +886-3-8562019; Tel: +886-3-8561825 ext. 13052

^bDepartment of Cosmetic Science, Providence University, 200 Sec. 7, Taiwan Boulevard, Shalu Dist., Taichung City 433, Taiwan, Republic of China. E-mail: yingshih@pu.edu.tw; Fax: +886-4-26311167; Tel: +886-4-26328001 ext. 15470

