Cite this: *Chem. Commun.*, 2016, 52, 6178

Received 14th January 2016,
Accepted 30th March 2016

DOI: 10.1039/c6cc00359a

www.rsc.org/chemcomm

Photocatalytic oxidation of iron(II) complexes by dioxygen using 9-mesityl-10-methylacridinium ions†

Takeshi Tsudaka,^a Kei Ohkubo^{*ab} and Shunichi Fukuzumi^{*abc}

Photocatalytic oxidation of iron(II) complexes by dioxygen occurred using the organic photocatalysts, 9-mesityl-10-methylacridinium ions (Acr⁺-Mes) and 2-phenyl-4-(1-naphthyl) quinolinium ions (QuPh⁺-NA), in the presence of triflic acid in acetonitrile under visible light irradiation. The electron-transfer state of Acr⁺-Mes produced upon photo-excitation oxidized the iron(II) complexes, whereas it reduced dioxygen with protons to produce iron(III) complexes and H₂O₂.

Metal complexes are usually oxidized by inorganic oxidants such as cerium ammonium nitrate and lead dioxide.^{1–5} In such cases, stoichiometric amounts of inorganic oxidants are required to obtain oxidized metal complexes, producing inorganic wastes which cause environmental problems. The ideal oxidant, which is environmentally benign, is dioxygen (O₂), producing only hydrogen peroxide or water as the reduced product. However, the oxidation of metal complexes by O₂ is often endergonic even in the presence of an acid. Thus, an appropriate photocatalyst is required for the oxidation of metal complexes by O₂ in the presence of an acid. Ruthenium(II) complexes, such as [Ru(bpy)₃]²⁺ (bpy = 2,2'-bipyridine) are known to be oxidized by O₂ in the presence of an acid to yield the corresponding Ru(III) complexes.^{6,7} On the other hand, organic photocatalysts have merited increasing attention for a variety of oxidation reactions.^{8–14} However, there has been no report on photocatalytic oxidation of metal complexes by O₂ using organic photocatalysts.

We report herein the photocatalytic oxidation of iron(II) complexes by O₂ using 9-mesityl-10-methylacridinium ions (Acr⁺-Mes)¹⁵ and 2-phenyl-4-(1-naphthyl)quinolinium ions (QuPh⁺-NA)¹⁶ as

Fig. 1 Visible absorption change in photocatalytic oxidation of [Fe^{II}(bpy)₃]²⁺-(PF₆⁻)₂ (2.0 mM) by O₂ in O₂-saturated MeCN in the presence of [Acr⁺-Mes]-(ClO₄⁻) (0.20 mM) and HOTf (0.10 M) at 298 K under visible light irradiation using a xenon lamp with a cut filter (λ < 390 nm).

organic photocatalysts in the presence of triflic acid (HOTf) in acetonitrile (MeCN) under visible light irradiation [eqn (1)]. Visible light irradiation of O₂-saturated acetonitrile is shown in Fig. 1, where the absorption band at 520 nm due to [Fe^{II}(bpy)₃]²⁺ decreased, accompanied by the increase in absorption at 650 nm due to [Fe^{III}(bpy)₃]³⁺. [Fe^{II}(bpy)₃]²⁺ was not oxidized without Acr⁺-Mes under irradiation (Fig. S1 and S2 in the ESI†). The reduced product of O₂ was H₂O₂, which was detected by spectral titration with the use of the oxo[5,10,15,20-tetra(4-pyridyl)porphyrinato]titanium(IV) complex (see the Experimental section in the ESI†).¹⁷

(1)

^a Department of Material and Life Science, Graduate School of Engineering, Osaka University, ALCA and SENTAN, Japan Science and Technology Agency (JST), Suita, Osaka 565-0871, Japan. E-mail: ookubo@chem.eng.osaka-u.ac.jp; Fax: +81-6-6879-7370

^b Department of Chemistry and Nano Science, Ewha Womans University, Seoul 120-750, Korea. E-mail: fukuzumi@chem.eng.osaka-u.ac.jp

^c Faculty of Science and Technology, Meijo University, ALCA and SENTAN, Japan Science and Technology Agency (JST), Nagoya, Aichi 468-8502, Japan

† Electronic supplementary information (ESI) available: Experimental and kinetic details. See DOI: 10.1039/c6cc00359a

The photocatalytic oxidation of other iron(II) complexes by O_2 was examined in the presence of HOTf in O_2 -saturated MeCN (Fig. S3–S6 in the ESI†). The quantum yields of formation of iron(III) complexes were determined using a ferrioxalate actinometer (see Fig. S7–S9 and the Experimental section in the ESI†).¹⁸ The quantum yields (Φ) of photocatalytic oxidation of iron(II) complexes by O_2 in the presence of HOTf in MeCN are listed in Table 1 together with the concentrations of iron(II) complexes, HOTf and O_2 , the one-electron oxidation potentials of iron(II) complexes (E_{ox})¹⁹ and the free energy change of the oxidation (ΔG_{ox}). The ΔG_{ox} values were evaluated from the E_{ox} values and the E_{red} value of O_2 in the presence of an acid in MeCN (0.75 V vs. SCE) [eqn (2)].²⁰

$$\Delta G_{ox} = -2e(E_{red} - E_{ox}) \quad (2)$$

Judging from the ΔG_{ox} values, the photocatalytic oxidation of $[Fe^{II}(Clphen)_3]^{2+}$, $[Fe^{II}(bpy)_3]^{2+}$ and $[Fe^{II}(Me_2bpy)_3]^{2+}$ is endergonic, whereas that of $Fe^{II}(BrC_5H_4)_2$ and $Fe^{II}(BrC_5H_4)(C_5H_5)$ is exergonic. The Φ values increased with the decreasing the E_{ox} values of iron(II) complexes as the free energy change of the reaction in eqn (1) decreased to be thermodynamically more favourable. Dependence of Φ on concentrations of $[Fe^{II}(bpy)_3]^{2+}(PF_6^-)_2$, HOTf and O_2 is shown in Fig. 2 (parts a, b and c, respectively). The Φ value became constant with the increasing concentrations of $[Fe^{II}(bpy)_3]^{2+}(PF_6^-)_2$, HOTf and O_2 , respectively. The photocatalytic oxidation of iron(II) complexes is enhanced by using QuPh⁺-NA instead of Acr⁺-Mes as shown in Table 1. (Fig. S10–S14 in the ESI†).

Nanosecond laser flash photolysis measurements were performed in order to clarify the catalytic mechanism for photocatalytic oxidation of iron(II) complexes by O_2 using Acr⁺-Mes in the presence of HOTf in MeCN. Transient absorption spectra were taken after the nanosecond laser excitation at 355 nm of a deaerated MeCN solution of Acr⁺-Mes in the absence and presence of $[Fe^{II}(bpy)_3]^{2+}(PF_6^-)_2$ as shown in Fig. 3. The transient absorption band at 490 nm is due to the electron-transfer state of Acr⁺-Mes. In the presence of $[Fe^{II}(bpy)_3]^{2+}$ the absorption at 490 nm decayed more rapidly and the decay rate increased with the increasing concentration of $[Fe^{II}(bpy)_3]^{2+}$. The decay rate obeyed pseudo-first-order kinetics and the pseudo-first order rate constant increased linearly with the increasing concentration of $[Fe^{II}(bpy)_3]^{2+}(PF_6^-)_2$. From the slope the rate constant (k_{ox}) of electron transfer from $[Fe^{II}(bpy)_3]^{2+}$ to the electron-transfer state of Acr⁺-Mes was determined to be

Fig. 2 Dependence of Φ on concentrations of (a) $[Fe^{II}(bpy)_3]^{2+}$, (b) HOTf and (c) O_2 . Standard conditions: $[Acr^+-Mes](ClO_4^-) = 1.0$ mM; $[Fe^{II}(bpy)_3]^{2+}(PF_6^-)_2 = 2.0$ mM; $[HOTf] = 0.15$ M; $[O_2] = 2.6$ mM; dehydrated MeCN (0.40 mL); cell path length 0.1 cm; excitation wavelength: 420 nm.

$3.7 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ as shown in the inset of Fig. 3. Similarly the k_{ox} values of other iron(II) complexes were determined (see Fig. S15–S23 in the ESI†) as listed in Table 2 together with the E_{ox} values. The k_{ox} value increases with the decreasing E_{ox} values. In the presence of O_2 , electron transfer from the electron-transfer state of Acr⁺-Mes to O_2 is known to occur with a rate constant of $6.8 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$.²¹ The rate constant of electron transfer from the electron-transfer state of QuPh⁺-NA to O_2 was determined to be $6.3 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ (see Fig. S24 in the ESI†). Thus, the photocatalytic oxidation of iron(II) complexes (Fe^{II}) by O_2 in the presence of HOTf proceeds as shown in Scheme 1. Photoexcitation of Acr⁺-Mes results in the formation of the electron-transfer state of Acr⁺-Mes, which oxidizes iron(II) complexes to iron(III) complexes and reduces O_2 with protons to produce HO_2^* , which disproportionates to yield H_2O_2 .

In conclusion, iron(II) complexes are oxidized to iron(III) complexes by O_2 using Acr⁺-Mes as an organic photocatalyst in the presence of HOTf in MeCN under visible light irradiation *via* electron-transfer oxidation of iron(II) complexes and reduction of

Table 1 Product and quantum yields (Φ) of iron(III) complexes for photocatalytic oxidation of iron(II) complexes by O_2 in the presence of HOTf together with the one-electron oxidation potentials of iron(II) complexes (E_{ox}) and the free energy change of the oxidation (ΔG_{ox})

Entry	Fe^{II} complex	E_{ox} vs. SCE ^a , V	ΔG_{ox} , eV	Acr ⁺ -Mes yield ^d , %	TON	Φ^f , %	QuPh ⁺ -NA yield ^e , %	TON	Φ^f , %
1 ^b	$[Fe^{II}(Clphen)_3]^{2+}$	1.20	0.90	5	0.25	0.11	30	1.5	0.34
2 ^b	$[Fe^{II}(bpy)_3]^{2+}$	1.06	0.62	42	4.2	0.32	32	3.5	0.73
3 ^b	$[Fe^{II}(Me_2bpy)_3]^{2+}$	0.88	0.26	22	2.1	1.6	30	3.0	5.2
4 ^c	$Fe^{II}(BrC_5H_4)_2$	0.72	−0.06	81	4.1	7.2	87	4.2	13
5 ^c	$Fe^{II}(BrC_5H_4)(C_5H_5)$	0.53	−0.44	81	4.1	19	60	3.0	26

Clphen = 5-chloro-1,10-phenanthroline, bpy = 2,2'-bipyridine, Me₂bpy = 4,4'-dimethyl-2,2'-bipyridine, BrC₅H₄ = bromocyclopentadienyl, C₅H₅ = cyclopentadienyl, reaction conditions: [photocatalyst] = 0.20 mM; [HOTf] = 0.10 M; $[Fe^{II}(Clphen)_3]^{2+}(PF_6^-)_2 = 1.0$ mM, $[Fe^{II}(bpy)_3]^{2+}(PF_6^-)_2 = 2.0$ mM, $[Fe^{II}(Me_2bpy)_3]^{2+}(PF_6^-)_2 = 2.0$ mM, $[Fe^{II}(BrC_5H_4)_2] = 1.0$ mM, $[Fe^{II}(BrC_5H_4)(C_5H_5)] = 2.0$ mM. ^a Taken from ref. 19. ^b MeCN (0.40 mL), cell path length 0.1 cm. ^c MeCN (3.0 mL), cell path length 1 cm. ^d Photoirradiation ($\lambda > 390$ nm). ^e ($\lambda > 300$ nm). ^f See the Experimental section in the ESI.

Fig. 3 Transient absorption decay at 490 nm due to the electron-transfer state of $[\text{Acr}^+-\text{Mes}](\text{ClO}_4^-)$ with various concentrations of $[\text{Fe}^{\text{II}}(\text{bpy})_3]^{2+}(\text{PF}_6^-)_2$. Inset: Decay rate constant versus concentrations of $[\text{Fe}^{\text{II}}(\text{bpy})_3]^{2+}(\text{PF}_6^-)_2$.

Table 2 Rate constants of electron transfer from iron(II) complexes to the electron-transfer state of the organic photocatalyst and the one-electron oxidation potentials of iron(II) complexes (E_{ox})

Entry	Fe^{II} complex	E_{ox} vs. SCE, V	Acr^+-Mes k_{et} , $\text{M}^{-1} \text{s}^{-1}$	QuPh^+-NA k_{et} , $\text{M}^{-1} \text{s}^{-1}$
1	$[\text{Fe}^{\text{II}}(\text{Clphen})_3]^{2+}$	1.20	1.4×10^8	4.9×10^8
2	$[\text{Fe}^{\text{II}}(\text{bpy})_3]^{2+}$	1.06	3.7×10^8	4.5×10^8
3	$[\text{Fe}^{\text{II}}(\text{Me}_2\text{bpy})_3]^{2+}$	0.88	4.5×10^8	5.8×10^8
4	$\text{Fe}^{\text{II}}(\text{BrC}_5\text{H}_4)_2$	0.72	7.6×10^9	7.2×10^9
5	$\text{Fe}^{\text{II}}(\text{BrC}_5\text{H}_4)(\text{C}_5\text{H}_5)$	0.53	8.5×10^9	7.9×10^9

Scheme 1 Photocatalytic cycle for oxidation of iron(II) complexes to iron(III) by O_2 with Acr^+-Mes .

O_2 by the electron-transfer state of Acr^+-Mes produced upon photoexcitation of Acr^+-Mes , respectively. The present study provides an environmentally benign approach for oxidation of metal complexes by O_2 to obtain the oxidised metal complexes and hydrogen peroxide (H_2O_2). Because there are many synthetically useful oxidation reactions using H_2O_2 ,²² this study has paved a new

way for photocatalytic oxidation of substrates by O_2 with organic photocatalysts and iron(II) complexes.

This work was supported by Grants-in-Aid (no. 26620154 and 26288037 to K. O.) from the Ministry of Education, Culture, Sports, Science and Technology (MEXT); ALCA and SENTAN projects from JST, Japan (to S. F.).

Notes and references

- (a) L. Duan, A. Fischer, Y. Xu and L. Sun, *J. Am. Chem. Soc.*, 2009, **131**, 10397; (b) T. Ishizuka, S. Ohzu and T. Kojima, *Synlett*, 2014, 1667.
- (a) M. Murakami, D. Hong, T. Suenobu and S. Fukuzumi, *J. Am. Chem. Soc.*, 2011, **133**, 11605; (b) T. Kojima, K. Nakayama, K. Ikemura, T. Ogura and S. Fukuzumi, *J. Am. Chem. Soc.*, 2011, **133**, 11692.
- D. Hong, S. Mandal, Y. Yamada, Y.-M. Lee, W. Nam, A. Llobet and S. Fukuzumi, *Inorg. Chem.*, 2013, **52**, 9522–9531.
- (a) G. P. McDermott, P. Jones, N. W. Barnett, D. N. Donaldson and P. S. Francis, *Anal. Chem.*, 2011, **83**, 5453; (b) M. P. Patel, S. A. Varnum, D. Gandla, M. J. Zdilla and C. J. Martoff, *Anal. Methods*, 2014, **6**, 5818.
- (a) S. Fukuzumi, C. L. Wong and J. K. Kochi, *J. Am. Chem. Soc.*, 1980, **102**, 2928; (b) S. Fukuzumi, K. Miyamoto, T. Suenobu, E. Van Caemelbecke and K. M. Kadish, *J. Am. Chem. Soc.*, 1998, **120**, 2880–2889; (c) S. Fukuzumi, H. Miyao and T. Suenobu, *J. Phys. Chem. A*, 2005, **109**, 3285–3294.
- A. Das, V. Joshi, D. Kotkar, V. S. Pathak, V. Swayambunathan, P. V. Kamat and P. K. Ghosh, *J. Phys. Chem. A*, 2001, **105**, 6945–6954.
- S. Kato, J. Jung, T. Suenobu and S. Fukuzumi, *Energy Environ. Sci.*, 2013, **6**, 3756–3764.
- (a) M. Fagnoni, D. Dondi, D. Ravelli and A. Albini, *Chem. Rev.*, 2007, **107**, 2725; (b) D. Ravelli, M. Fagnoni and A. Albini, *Chem. Soc. Rev.*, 2013, **42**, 97.
- (a) D. A. Nicewicz and T. M. Nguyen, *ACS Catal.*, 2014, **4**, 355–360; (b) D. A. Nicewicz and D. S. Hamilton, *Synlett*, 2014, 1191–1196.
- D. Ravelli and M. Fagnoni, *ChemCatChem*, 2012, **4**, 169.
- (a) K. Ohkubo and S. Fukuzumi, *Bull. Chem. Soc. Jpn.*, 2009, **82**, 303; (b) S. Fukuzumi and K. Ohkubo, *Chem. Sci.*, 2013, **4**, 561.
- D. Ravelli, D. Dondi, M. Fagnoni and A. Albini, *Chem. Soc. Rev.*, 2009, **38**, 1999.
- (a) M. L. Marin, L. Santos-Juanes, A. Arques, A. M. Amat and M. A. Miranda, *Chem. Rev.*, 2012, **112**, 1710; (b) M. L. Marin, A. Miguel, L. Santos-Juanes, A. Arques, A. M. Amat and M. A. Miranda, *Photochem. Photobiol. Sci.*, 2007, **6**, 848.
- Y. Pan, S. Wang, C. W. Kee, E. Dubuisson, Y. Yang, K. P. Loh and C.-H. Tan, *Green Chem.*, 2011, **13**, 3341.
- (a) S. Fukuzumi, H. Kotani, K. Ohkubo, S. Ogo, N. V. Tkachenko and H. Lemmetyinen, *J. Am. Chem. Soc.*, 2004, **126**, 1600; (b) S. Fukuzumi, H. Kotani and K. Ohkubo, *Phys. Chem. Chem. Phys.*, 2008, **10**, 5159.
- (a) H. Kotani, K. Ohkubo and S. Fukuzumi, *Faraday Discuss.*, 2012, **155**, 89; (b) Y. Yamada, T. Miyahigashi, H. Kotani, K. Ohkubo and S. Fukuzumi, *J. Am. Chem. Soc.*, 2011, **133**, 16136.
- C. Matsubara, N. Kawamoto and K. Takamura, *Analyst*, 1992, **117**, 1781.
- C. G. Hatchard and C. A. Parker, *Proc. R. Soc. London, Ser. A*, 1956, **235**, 518–536.
- (a) J. Park, Y.-M. Lee, W. Nam and S. Fukuzumi, *J. Am. Chem. Soc.*, 2013, **135**, 5052; (b) H. Yoon, Y.-M. Lee, X. Wu, K.-B. Cho, R. Sarangi, W. Nam and S. Fukuzumi, *J. Am. Chem. Soc.*, 2013, **135**, 9186.
- R. Cofé and D. T. Sawyer, *Inorg. Chem.*, 1986, **25**, 2089.
- H. Kotani, K. Ohkubo and S. Fukuzumi, *J. Am. Chem. Soc.*, 2004, **126**, 15999.
- (a) A. Fingerhut, O. V. Serdyuk and S. B. Tsogoeva, *Green Chem.*, 2015, **17**, 2042; (b) Y. Zhu, Q. Wang, R. G. Cornwall and Y. Shi, *Chem. Rev.*, 2014, **114**, 8199; (c) S. Fukuzumi and K. Ohkubo, *Asian J. Org. Chem.*, 2015, **4**, 836.