


 Cite this: *Chem. Commun.*, 2016, 52, 6601

 Received 21st February 2016,  
 Accepted 11th April 2016


DOI: 10.1039/c6cc01572g

[www.rsc.org/chemcomm](http://www.rsc.org/chemcomm)

**A new reaction mode for bicyclo[1.1.0]tetraphosphabutanes is reported. The C=S and C=N bonds of phenyl isothiocyanate reversibly insert into a P–P bond of  $[(\text{CpNi}(\text{IMes}))_2(\mu\text{-}\eta^1\text{-}\eta^1\text{-P}_4)]$  ( $\text{1}^{\text{Mes}}$ , IMes = 1,3-bis(2,4,6-trimethylphenyl)imidazolin-2-ylidene), forming isomers **2a** and **2b**. X-ray crystallography and  $^{31}\text{P}^{\{1\text{H}\}}$  NMR spectroscopy revealed similar bicyclo[3.1.0]heterohexane structures for these compounds.**

Developing new, targeted and selective methods for the functionalisation of the  $\text{P}_4$  molecule remains a topical challenge despite the extensive research efforts carried out in the past.<sup>1,2</sup> Recent reports have focused on the use of nucleophilic carbanions and carbenes,<sup>3,4</sup> insertion reactions of p-block elements, *e.g.* phosphonium cations<sup>5</sup> and the use of main group element or transition metal-based radicals.<sup>6,7</sup> The latter approach often gives rise to bicyclo[1.1.0]tetraphosphabutanes **A**, which may be seen as potential intermediates on the way to a stepwise  $\text{P}_4$  degradation sequence (Scheme 1a). While various ‘ $\text{P}_4$  butterfly’ compounds of type **A** are known, it is interesting to note that their reactivity has only been explored to a small extent (Scheme 1b).<sup>1,6–9</sup> Previous studies mainly focused on iron complexes.<sup>1d,7a,c,9–11</sup> As reported by Scherer and Scheer, thermolysis or photolysis of  $[(\text{Cp}^{\text{R}}\text{Fe}(\text{CO}))_2(\mu\text{-}\eta^1\text{:}\eta^1\text{-P}_4)]$  ( $\text{Cp}^{\text{R}} = \text{C}_5\text{H}_2\text{-1,2,4-}t\text{Bu}_3$ ,  $\text{C}_5\text{H}_2\text{-1,2,4-}t\text{Bu}_3$ ,  $\text{C}_9\text{H}_5\text{-1,3-}t\text{Bu}_2$  and  $\text{C}_5\text{iPr}_5$ ) affords mixtures of polyphosphido complexes.<sup>7a,c</sup> Reactions with (phospha)alkynes evoked the  $\text{P}_3/\text{P}_1$  fragmentation of the bicyclo[1.1.0]tetraphosphabutane diyl fragment, forming phosphide, phospholide and diphosphacyclobutadiene components.<sup>9,10</sup> Further studies revealed that the ‘ $\text{P}_4$  butterfly’ may be protonated reversibly and coordinates as a chelate ligand to copper(i).<sup>11</sup> Here, we disclose a new reaction mode for metal-substituted bicyclo[1.1.0]tetraphosphabutanes. We have found

that phenyl isothiocyanate reversibly inserts into a P–P bond of the bicyclo[1.1.0]tetraphosphabutane scaffold of the dinuclear nickel complex  $[(\eta^5\text{-Cp})\text{Ni}(\text{IMes})_2(\mu\text{-}\eta^1\text{:}\eta^1\text{-P}_4)]$  ( $\text{1}^{\text{Mes}}$ , Scheme 1c).<sup>7b</sup> This unprecedented reaction affords the isomers **2a** and **2b**, which display a bicyclo[3.1.0]heterohexane skeleton. We describe the single-crystal X-ray structures and  $^{31}\text{P}^{\{1\text{H}\}}$  NMR data of these



**Scheme 1** (a) Formation of bicyclo[1.1.0]tetraphosphabutanes amenable for further transformations; (b) selected reactions of iron-substituted bicyclo[1.1.0]tetraphosphabutanes;  $\text{Cp}^{\text{R}} = \text{C}_5\text{H}_2\text{-1,2,4-}t\text{Bu}_3$ ,  $\text{C}_5\text{H}_2\text{-1,2,4-}t\text{Bu}_3$ ,  $\text{C}_9\text{H}_5\text{-1,3-}t\text{Bu}_2$ ,  $\text{C}_5\text{iPr}_5$ ,  $\text{R}^1 = \text{R}^2 = \text{Me, Ph}$ ;  $\text{R}^1 = \text{H}$ ,  $\text{R}^2 = \text{Ph, tBu, SiMe}_3$ ,  $\text{CO}_2\text{Me/Et}$ ,  $\text{R}^3 = \text{tBu, C}(\text{CH}_2)_5\text{Me}$ ; (c) synthesis of  $\text{1}^{\text{Dipp}}$  and  $\text{1}^{\text{Mes}}$  and reactivity toward phenyl isothiocyanate.<sup>6–10</sup>

<sup>a</sup> University of Regensburg, Institute of Inorganic Chemistry, 93040 Regensburg, Germany. E-mail: robert.wolf@ur.de

<sup>b</sup> Department of Chemistry and Pharmaceutical Sciences, VU University Amsterdam, De Boelelaan 1083, 1081 HV Amsterdam, The Netherlands

<sup>†</sup> Electronic supplementary information (ESI) available. CCDC 1446071–1446073. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c6cc01572g



new complexes and analyse the possible nature of additional reaction products using DFT calculations.

We recently synthesised the first nickel-substituted bicyclo[1.1.0]tetraphosphabutane,  $[(\eta^5\text{-Cp})\text{Ni}(\text{IDipp})_2(\mu\text{-}\eta^1\text{-}\eta^1\text{-P}_4)]$  (**1<sup>Dipp</sup>**, IDipp = 1,3-bis(2,6-diisopropylphenyl)imidazolin-2-ylidene).<sup>7b</sup> This complex is formed in a quantitative reaction from two equivalents  $[(\eta^5\text{-Cp})\text{Ni}(\text{IDipp})]$  and  $\text{P}_4$  (Scheme 1c). Subsequent work showed that the slightly less encumbered mesityl-substituted complex  $[(\eta^5\text{-Cp})\text{Ni}(\text{IMes})_2(\mu\text{-}\eta^1\text{-}\eta^1\text{-P}_4)]$  (**1<sup>Mes</sup>**) is obtained in an analogous fashion. **1<sup>Mes</sup>** was isolated as dark red air-sensitive crystals in 71% yield (Scheme 1c) and shows a better solubility than **1<sup>Dipp</sup>**, dissolving well in benzene, toluene, diethyl ether and tetrahydrofuran (ESI<sup>†</sup>).

In order to probe the reactivity of **1<sup>Dipp</sup>** and **1<sup>Mes</sup>**, we investigated reactions with heteroallenes. ADMX spin systems were observed by  $^{31}\text{P}$  NMR spectroscopy with  $\text{CS}_2$  (10 equiv.), suggesting an insertion into a P-P bond, but the products could not be isolated (ESI<sup>†</sup>). Isolable products were obtained with phenyl isothiocyanate, however. Monitoring the reaction of **1<sup>Mes</sup>** and PhNCS in  $[\text{D}_8]\text{THF}$  (Fig. 1) revealed that 7 equiv. PhNCS were necessary for full conversion of **1<sup>Mes</sup>** after four hours, while a large amount of **1<sup>Mes</sup>** (55%) remained in the reaction mixture with one equiv. PhNCS after one day (Fig. S10, ESI<sup>†</sup>). Two main products **2a** and **2b** (ADMX spin systems) and one minor species **2c** were detected (approximate ratio **2a** : **2b** : **2c** 75 : 20 : 5).<sup>‡</sup> The simultaneous formation of **2a**, **2b** and **2c** commences below 0 °C according to a VT NMR study ( $[\text{D}_8]\text{THF}$ , Fig. S8, ESI<sup>†</sup>). Prolonged reaction times and heating of the solution resulted in essentially the same product ratio, although the signal to noise ratio of the spectra decreased over time. In contrast, the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of the reaction of **1<sup>Dipp</sup>** with a large excess of PhNCS in  $[\text{D}_8]\text{THF}$  after two days at room temperature showed signals of a species similar to **2b** (15%, ADMX spin system), **1<sup>Dipp</sup>** (50%) and  $\text{P}_4$  (35%) (Fig. S9, ESI<sup>†</sup>).

Complex **2a** can be isolated as an analytically pure, dark brown solid in 31% yield by crystallising the crude product twice from toluene/n-hexane (ESI<sup>†</sup>). Crystallisation of the crude product from diethyl ether and recrystallization from toluene/n-hexane affords pure, crystalline **2b** in 16% isolated yield. Single-crystal XRD for **2a** (Fig. 2, top) revealed an unusual nickel-substituted bicyclo[3.1.0]-2-thia-1,4,5,6-tetraphosphahexane




Fig. 2 Solid-state molecular structures of **2a** (top) and **2b** (bottom). The hydrogen atoms are omitted for clarity. Thermal ellipsoids are drawn at 40% level. Selected bond lengths (Å) and angles (°) for **2a**: P1–C1 1.860(4), P4–S1 2.1257(13), P1–P2 2.1818(14), P2–P3 2.2182(15), P2–P4 2.2222(14), P3–P4 2.1935(15), C1–S1 1.794(4), C1–N1 1.278(6), Ni1–P1 2.2036(13), Ni2–P3 2.1906(11), P1–P2–P4 102.89(5), P2–P4–S1 102.86(6), P4–S1–C1 104.82(14), S1–C1–P1 122.3(2), C1–P1–P2 102.88(14), P3–P2–P4 59.20(5), P1–C1–N1 116.4(3), S1–C1–N1 121.3(2); for **2b**: P1–C1 1.828(3), P4–N1 1.785(3), P1–P2 2.2157(11), P2–P4 2.1969(10), P2–P3 2.2233(10), P3–P4 2.206(1), C1–N1 1.359(4), C1–S1 1.678(3), Ni1–P1 2.2188(9), Ni2–P3 2.2192(9), P1–P2–P4 95.28(4), P2–P4–N1 99.75(9), P4–N1–C1 124.9(2), N1–C1–P1 118.6(2), C1–P1–P2 100.48(3), P3–P2–P4 59.88(3), P1–C1–S1 117.68(18), N1–C1–S1 123.6(2).




Fig. 1 Synthesis of **2a** and **2b** (top), and  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum ( $[\text{D}_8]\text{THF}$ ) of the reaction of PhNCS and **1<sup>Mes</sup>** (7 : 1) at room temperature after four hours; ● = **2a**, ■ = **2b**, ▲ = **2c** (bottom).

moiety (P1–C1 1.860(4) Å, P4–S1 2.1257(13) Å) with an exocyclic imino function. The C1–N1 (1.278(6) Å) and S1–C1 (1.794(4) Å) bonds of **2a** are elongated compared to free aryl isothiocyanates.<sup>12</sup> The P–P distances (2.1818(14)–2.2222(14) Å) are in the range of single bonds.<sup>7</sup> The five-membered  $\text{CP}_3\text{S}$  heterocycle (P1–P2–P4–S1–C1) is almost flat ( $\sum_{\text{angles}} = 535.8^\circ$ ) and orthogonal ( $89.60(7)^\circ$ ) to the plane formed by P2, P3 and P4. The scaffold of **2a** is analogous to that of 2,3,4,6-tetra-*tert*-butylbicyclo[3.1.0]hexaphosphane synthesised by Baudler *et al.*<sup>13</sup>

The molecular structure of the regio isomer **2b** (Fig. 2, bottom) features a flat  $\text{CNP}_3$  heterocycle ( $\sum_{\text{angles}} = 539.0^\circ$ ) with a thioketone function (C1–S1 1.678(3) Å) and single bonds



between P1–C1 (1.828(3) Å) and P4–N1 (1.785(3) Å). The P–P distances in **2b** (2.1969(10)–2.2233(10) Å) are similar to those of **2a**. The CNP<sub>3</sub> ring forms an acute dihedral angle of 79.58(5)° with the P2–P3–P4 plane.

The <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of **2a** ([D<sub>8</sub>]THF, room temperature) features four broad multiplets at –150.1, –96.4, –75.0 and 32.1 ppm consistent with four chemically different P atoms. The signals are broad at room temperature (average half-width  $\tau_{\text{FWHM}} = 565$  Hz); they become sharper when the temperature is decreased to –80 °C (av.  $\tau_{\text{FWHM}} = 35$  Hz). Experimental and fitted <sup>31</sup>P{<sup>1</sup>H} NMR spectra in [D<sub>8</sub>]THF at –80 °C along with the assignment of the chemical shifts and coupling constants are shown in Fig. 3. The resonance at –151.8 ppm is assigned to P<sub>A</sub> connected to three P atoms based on the observation of three large <sup>1</sup>J(P,P)-coupling constants for this multiplet (<sup>1</sup>J(P<sub>APD</sub>) = –178 Hz, <sup>1</sup>J(P<sub>APM</sub>) = –185 Hz and <sup>1</sup>J(P<sub>APX</sub>) = –374 Hz). The P atoms coordinated to nickel ( $\delta(P_D) = –105.5$  ppm;  $\delta(P_X) = 27.8$  ppm) show a common large <sup>2</sup>J(P,P) coupling (<sup>2</sup>J(P<sub>D</sub>,P<sub>X</sub>) = 82 Hz), which may arise from an interaction of the lone pairs due to the conformational constraints of the bicyclo[3.1.0]heterohexane skeleton.<sup>13</sup>

Complex **2b** gives rise to four slightly broad <sup>31</sup>P{<sup>1</sup>H} NMR resonances at –182.1, –104.5, 4.5 and 50.1 ppm in [D<sub>8</sub>]THF at




Fig. 3 <sup>31</sup>P{<sup>1</sup>H} NMR spectra of **2a** (top) and **2b** (bottom, 161.98 MHz, [D<sub>8</sub>]THF, 193 K); **2a** (ADMX spin system):  $\delta_A = –151.8$  ppm,  $\delta_D = –105.5$  ppm,  $\delta_M = –78.0$  ppm,  $\delta_X = 27.8$  ppm, <sup>1</sup>J(P<sub>APD</sub>) = –178 Hz, <sup>1</sup>J(P<sub>APM</sub>) = –185 Hz, <sup>1</sup>J(P<sub>APX</sub>) = –374 Hz, <sup>1</sup>J(P<sub>D</sub>P<sub>M</sub>) = –238 Hz, <sup>2</sup>J(P<sub>D</sub>P<sub>X</sub>) = 82 Hz, <sup>2</sup>J(P<sub>M</sub>P<sub>X</sub>) = 9 Hz; **2b** (ADMX spin system):  $\delta_A = –180.5$  ppm,  $\delta_D = –109.1$  ppm,  $\delta_M = 12$  ppm,  $\delta_X = 48.2$  ppm, <sup>1</sup>J(P<sub>APD</sub>) = –193 Hz, <sup>1</sup>J(P<sub>APM</sub>) = –209 Hz, <sup>1</sup>J(P<sub>APX</sub>) = –282 Hz, <sup>1</sup>J(P<sub>D</sub>P<sub>M</sub>) = –237 Hz, <sup>2</sup>J(P<sub>D</sub>P<sub>X</sub>) = 57 Hz, <sup>2</sup>J(P<sub>M</sub>P<sub>X</sub>) = 10 Hz; expansions (inset) show the experimental (up) and fitted spectra (down). The signals assigned to **2b** are labeled with an asterisk.




Fig. 4 Calculated, relative Gibbs free energies (kcal mol<sup>–1</sup>) of **1<sup>Ph</sup>**, **2a<sup>Ph</sup>** and **2b<sup>Ph</sup>**. The relative Gibbs free energies refer to **1<sup>Ph</sup>** + PhNCS (kcal mol<sup>–1</sup>).

room temperature. The line width decreased from an average of  $\tau_{\text{FWHM}} = 33$  Hz at room temperature to  $\tau_{\text{FWHM}} = 23$  Hz upon cooling to –80 °C. The chemical shifts and coupling constants of **2b** lie in a similar range as observed for **2a** (Fig. 3, bottom) in agreement with the similar structure motif.

<sup>31</sup>P{<sup>1</sup>H} NMR studies indicate that the formation of **2a**, **2b** and **2c** is reversible; *i.e.* the products slowly equilibrate with the starting material **1<sup>Mes</sup>** in solution (ESI†). A mixture of **2a** (89%), **1<sup>Mes</sup>** (7%), **2c** (4%) and **2b** (traces) was detected upon storing a [D<sub>8</sub>]THF solution of pure **2a** in an NMR tube at room temperature for two days, while a 65:10:5:20 mixture (**2a**:**2b**:**2c**:**1<sup>Mes</sup>**) was present after one week.§ Additional multiplets of unidentified minor species can be observed upon prolonged storage (Fig. S11, ESI†). **2b** behaves similarly (Fig. S12, ESI†). IR monitoring of the decomposition of **2a** ([D<sub>8</sub>]THF, 60 °C, 13.5 hours) shows the formation of free PhNCS (Fig. S13, ESI†).

DFT calculations (ωB97X-D/6-311G(d,p) level)<sup>14</sup> were performed to gain additional insight into the thermodynamics of the reaction. The optimized structures of the truncated model complexes **1<sup>Ph</sup>**, **2a<sup>Ph</sup>** and **2b<sup>Ph</sup>**, where the Mes substituents were replaced by phenyl groups for computational efficiency, are in good agreement with the experimental structures (Fig. 4). The formation of **2a<sup>Ph</sup>** and **2b<sup>Ph</sup>** is exergonic, and the thermodynamic product of the reaction appears to be **2b<sup>Ph</sup>** (–15.2 kcal mol<sup>–1</sup> with respect to the starting materials), while **2a<sup>Ph</sup>** (–11.3 kcal mol<sup>–1</sup>) is a kinetic product.¶

In conclusion, the reaction of **1<sup>Mes</sup>** with PhNCS affords the novel complexes **2a** and **2b** with an unusual bicyclo[3.1.0]heterohexane skeleton. To our knowledge, this represents the first example of an insertion of a heteroallene into a P–P bond of a cyclopolypyrophane. In future work, it will be of interest to investigate whether similar reactions with polar multiple bonds offer a general route toward “functionalized” polyphosphanes.<sup>15</sup> Efficient preparative methods exist for a range of bicyclo[1.1.0]tetraphosphabutanes,<sup>1,6–9</sup> therefore, such transformations may provide a fruitful avenue to the stepwise and selective degradation of the P<sub>4</sub> molecule.

We thank B. Sc. Thomas Maier for experimental assistance. Funding by the Deutsche Forschungsgemeinschaft is gratefully acknowledged.

## Notes and references

‡ <sup>31</sup>P{<sup>1</sup>H} NMR data of **2c** ([D<sub>8</sub>]THF, A<sub>2</sub>MX spin system):  $\delta = –321.2$  (dd, 2P, P<sub>A</sub>, <sup>1</sup>J(P<sub>APM</sub>) = –178 Hz, <sup>1</sup>J(P<sub>APX</sub>) = –188 Hz, –141.5 (dt, 1P, P<sub>M</sub>, <sup>2</sup>J(P<sub>M</sub>P<sub>X</sub>) = 208 Hz), –85.8 (dt, 1P, P<sub>X</sub>) ppm.

§ The <sup>1</sup>H NMR spectrum of a freshly prepared [D<sub>8</sub>]THF solution of pure crystals of **2a** stored for one week at room temperature in an Ar-filled glove box also showed a mixture containing **2a**, **2b** and **1<sup>Mes</sup>** in a 94.5:0.5:5 ratio.

The structure and the mechanism of formation of the minor product **2c** ( $A_2MX$  spin system, *vide supra*) presently remains unclear. Five potential candidates were identified by our computations (Fig. S14, ESI†). These calculated isomers are adducts of the starting material with PhNCS ( $2c^{Add1}$  and  $2c^{Add2}$ ) or result from the insertion of the C=S or C≡N double bonds into the Ni–P bond ((E)- $2c^{Ins1}$ , (Z)- $2c^{Ins1}$  and  $2c^{Ins2}$ ).<sup>16</sup> Each of them has an energy significantly higher than that of **2a<sup>Ph</sup>** and **2b<sup>Ph</sup>**.

- 1 Reviews: (a) B. M. Cossairt, N. A. Piro and C. C. Cummins, *Chem. Rev.*, 2010, **110**, 4164; (b) M. Caporali, L. Gonsalvi, A. Rossin and M. Peruzzini, *Chem. Rev.*, 2010, **110**, 4178; (c) M. Peruzzini, I. de los Rios, A. Romerosa and F. Vizza, *Eur. J. Inorg. Chem.*, 2001, 593; (d) M. Peruzzini, R. R. Abdreimova, Y. Budnikova, A. Romerosa, O. J. Scherer and H. Sitzmann, *J. Organomet. Chem.*, 2004, **689**, 4319; (e) M. Peruzzini, L. Gonsalvi and A. Romerosa, *Chem. Soc. Rev.*, 2005, **34**, 1038.
- 2 Selected publications on transformations of transition metal phosphide complexes derived from  $P_4$  in one step: (a) P. Barbaro, C. Bazzicalupi, M. Peruzzini, S. Seniori Costantini and P. Stoppioni, *Angew. Chem., Int. Ed.*, 2012, **51**, 8628; (b) E. Mädl, M. V. Butovskii, G. Balázs, E. V. Peresypkina, A. V. Virovets, M. Seidl and M. Scheer, *Angew. Chem., Int. Ed.*, 2014, **53**, 7643.
- 3 (a) W. T. K. Chan, F. García, A. D. Hopkins, L. C. Martin, M. McPartlin and D. S. Wright, *Angew. Chem., Int. Ed.*, 2007, **46**, 3084; (b) J. E. Borger, A. W. Ehlers, M. Lutz, J. C. Slootweg and K. Lammertsma, *Angew. Chem., Int. Ed.*, 2014, **53**, 12836; (c) M. Arrowsmith, M. S. Hill, A. L. Johnson, G. Kociok-Köhn and M. F. Mahon, *Angew. Chem., Int. Ed.*, 2015, **54**, 7882; (d) J. E. Borger, A. W. Ehlers, M. Lutz, J. C. Slootweg and K. Lammertsma, *Angew. Chem., Int. Ed.*, 2015, **55**, 613.
- 4 (a) J. D. Masuda, W. W. Schoeller, B. Donnadieu and G. Bertrand, *Angew. Chem., Int. Ed.*, 2007, **46**, 7052; (b) O. Back, G. Kuchenbeiser, B. Donnadieu and G. Bertrand, *Angew. Chem., Int. Ed.*, 2009, **48**, 5530.
- 5 Selected publications on the insertion of p-block elements into the P–P bonds of  $P_4$ : (a) Y. Xiong, S. Yao, M. Brym and M. Driess, *Angew. Chem., Int. Ed.*, 2007, **46**, 4511; (b) J. J. Weigand, M. Holthausen and R. Fröhlich, *Angew. Chem., Int. Ed.*, 2009, **48**, 295; (c) M. H. Holthausen and J. J. Weigand, *J. Am. Chem. Soc.*, 2009, **131**, 14210; (d) G. Prabusankar, A. Doddi, C. Gemel, M. Winter and R. A. Fischer, *Inorg. Chem.*, 2010, **49**, 7976; (e) M. H. Holthausen and J. J. Weigand, *Chem. Soc. Rev.*, 2014, **43**, 6639.
- 6 (a) J.-P. Bezombes, P. B. Hitchcock, M. F. Lappert and J. E. Nycz, *Dalton Trans.*, 2004, 499; (b) B. M. Cossairt and C. C. Cummins, *New J. Chem.*, 2010, **34**, 1533; (c) S. Khan, R. Michel, J. M. Dieterich, R. A. Mata, H. W. Roesky, J.-P. Demers, A. Lange and D. Stalke, *J. Am. Chem. Soc.*, 2011, **133**, 17889; (d) N. A. Giffin, A. D. Hendsbee, T. L. Roemmele, M. D. Lumsden, C. C. Pye and J. D. Masuda, *Inorg. Chem.*, 2012, **51**, 11837; (e) S. Heinl, S. Reisinger, C. Schwarzmaier, M. Bodensteiner and M. Scheer, *Angew. Chem., Int. Ed.*, 2014, **53**, 7639.
- 7 (a) O. J. Scherer, T. Hilt and G. Wolmershäuser, *Organometallics*, 1998, **17**, 4110; (b) S. Pelties, D. Herrmann, B. de Bruin, F. Hartl and R. Wolf, *Chem. Commun.*, 2014, **50**, 7014; (c) S. Heinl and M. Scheer, *Chem. Sci.*, 2014, **5**, 3221; (d) C. Schwarzmaier, A. Y. Timoshkin, G. Balázs and M. Scheer, *Angew. Chem., Int. Ed.*, 2014, **53**, 9077; (e) D. W. Agnew, C. E. Moore, A. L. Rheingold and J. S. Figueiroa, *Angew. Chem., Int. Ed.*, 2015, **54**, 12673.
- 8 Selected main group element-substituted “butterfly  $P_4$ ” compounds with an  $E_2P_4$  core: (a) E. Niecke, R. Rüger and B. Krebs, *Angew. Chem., Int. Ed. Engl.*, 1982, **21**, 544; (b) R. Riedel, H.-D. Hausen and E. Fluck, *Angew. Chem., Int. Ed. Engl.*, 1985, **24**, 1056; (c) S. L. Hinchley, C. A. Morrison, D. W. H. Rankin, C. L. B. Macdonald, R. J. Wiacek, A. Voigt, A. H. Cowley, M. F. Lappert, G. Gunderson, J. A. C. Clyburne and P. P. Power, *J. Am. Chem. Soc.*, 2001, **123**, 9045.
- 9 (a) O. J. Scherer, T. Hilt and G. Wolmershäuser, *Angew. Chem., Int. Ed.*, 2000, **39**, 1425; (b) C. Eichhorn, PhD thesis, Kaiserslautern, 2003; (c) S. Deng, C. Schwarzmaier, C. Eichhorn, O. J. Scherer, G. Wolmershäuser, M. Zabel and M. Scheer, *Chem. Commun.*, 2008, 4064.
- 10 M. Scheer, S. Deng, O. J. Scherer and M. Sierka, *Angew. Chem., Int. Ed.*, 2005, **44**, 3755.
- 11 C. Schwarzmaier, S. Heinl, G. Balázs and M. Scheer, *Angew. Chem., Int. Ed.*, 2015, **54**, 13116.
- 12 S. Biswas, S. Haldar, P. K. Mandal, K. Goubitz, H. Schenk and R. Dabrowski, *Cryst. Res. Technol.*, 2007, **42**, 1029.
- 13 (a) M. Baudler, Y. Aktalay, K.-F. Tebbe and T. Heinlein, *Angew. Chem., Int. Ed. Engl.*, 1981, **20**, 967; in this context see also: (b) M. Baudler, *Angew. Chem., Int. Ed. Engl.*, 1982, **21**, 492.
- 14 DFT calculations were performed using Gaussian09; see ESI† for further details.
- 15 (a) A. R. Jupp and J. M. Goicoechea, *Angew. Chem., Int. Ed.*, 2013, **52**, 10064; (b) R. S. P. Turberville and J. M. Goicoechea, *Chem. Commun.*, 2012, **48**, 6100; (c) R. S. P. Turberville, A. R. Jupp, P. S. B. McCullough, D. Ergögmen and J. M. Goicoechea, *Organometallics*, 2013, **32**, 2234.
- 16 Selected examples for the insertion of isothiocyanates into transition metal-phosphorus bonds: (a) U. Segerer and E. Hey-Hawkins, *Polyhedron*, 1997, **16**, 2537; (b) U. Segerer, J. Sieler and E. Hey-Hawkins, *Organometallics*, 2000, **19**, 2445; (c) A. Antiñolo, S. García-Yuste, A. Otero and R. Reguillo-Carmona, *Eur. J. Inorg. Chem.*, 2009, 539; (d) W. Yi, J. Zhang, L. Hong, Z. Chen and X. Zhou, *Organometallics*, 2011, **30**, 5809.

