ChemComm

COMMUNICATION

Cite this: *Chem. Commun.,* 2016, 52, 8389

Received 31st March 2016, Accepted 3rd June 2016

DOI: 10.1039/c6cc02720b

www.rsc.org/chemcomm

Catalytic dehydrocoupling of amine-boranes and amines into diaminoboranes: isolation of a Pt(II), Shimoi-type, $\eta^{\texttt{1-BH}}$ complex \dagger

Marta Roselló-Merino,^a Raquel J. Rama,^a Josefina Díez^b and Salvador Conejero*^a

The platinum complex [Pt(I^tBu')(I^tBu)][BAr^F] is a very efficient catalyst in the synthesis of diaminoboranes through dehydrocoupling of amine-boranes and amines. Shimoi-type, η^{1} -BH complexes are key intermediates in the process.

In the last few years amine-boranes and related base-stabilised borane adducts have been shown to produce rich chemistry in which metal catalysed dehydrocoupling processes are involved.¹ During some of these reactions the B–H bonds of the amine borane establish an initial interaction with the metal centre to form complexes exhibiting η^1 or η^2 coordination modes, undergoing thereafter subsequent reactivities.^{1a,2}

Dehydrogenation of amine-boranes has been mainly focused on the production and release of dihydrogen and on the generation of amino-boranes that can undergo further dimerization, oligomerization or polymerization processes.^{1,2*j*,3} Less attention has been paid to the production of other boranes, through boron–boron coupling⁴ or the formation of diaminoboranes, $(NR_2)_2$ BH.⁵ With regard to the latter process, only a couple of catalytic processes have been reported by Alcaraz and Sabo-Etienne, using ruthenium^{5a,c} and by Hill,^{5b} with calcium and magnesium based catalysts to produce the corresponding diaminoboranes. Catalyst loadings of 2.5% (Ru, Ca, Mg), together with long reaction times (Ru), and even heating (Ca, Mg) were required.

In this communication we wish to report a very efficient platinum (n) complex that promotes diaminoborane formation at a catalyst loading of 0.5% in a few minutes for most of the substrates tested. In addition, we have been able to characterise

by means of X-ray diffraction studies the first Shimoi-type η^1 -BH complex of a Lewis base-borane adduct and platinum, which provides structural insights into a key intermediate in the dehydrogenation process of amine-boranes and amines.

Previously, we had reported that the $[Pt(I^tBu') (I^tBu)][BAr^F]$ complex (where I^tBu stands for 1,3-di-tert-butylimidazolylidene and I'Bu' its cyclometalated form), $1,6$ is able to promote the dehydrocoupling of dimethylamineborane $(NMe₂H·BH₃)$ into cyclic $\text{[NMe}_2\text{BH}_2\text{]}_2$ (Scheme 1).⁷ During NMR mechanistic studies, we observed that bis(dimethylamino)borane $(NMe₂)₂BH$ is also formed in small amounts and its yield increases as the concentration of free $NMe₂H$ in the reaction media increases. Therefore, we analysed the ability of complex 1 to act as a catalyst for the formation of diaminoboranes. COMMUNICATION

Constant Catalytic dehydrocoupling of amine-boranes and

annines into diaminoboranes: isolation of a Pt(u),

Shimoi-type, η ¹–BH complex²

Shimoi-type, η ¹–BH complex²

Shimoi-type, η ¹–BH c

We have first studied the benchmark reaction of tert-butylamineborane $(^tBuNH_2\cdot BH_3)$ and 1 equiv. of *tert*-butylamine (t BuNH₂) in CD₂Cl₂, using a catalyst loading of 0.5% (Scheme 2). The reaction was monitored both by NMR spectroscopy and by measuring the increase of gas pressure in a closed system due to the generated H_2 (see the ESI†). The ¹¹B NMR spectrum revealed the formation of a single species in nearly quantitative yield after 11.5 min showing a doublet signal at δ 25.8 ppm (1 J_{BH} = 127 Hz). In addition, the ¹H NMR spectrum exhibits a broad quartet signal centred at 4.12 ppm that sharpens upon $11B$ decoupling into a triplet ($J_{HH} = 8.2$ Hz). These values are in agreement with the formation of diaminoborane $(^t$ BuNH)₂BH,⁸ s1 (Scheme 2). The calculated TON and TOF values for this process are 400 and 2087 h^{-1} , the highest reported to the best of our knowledge (Table 1).

Scheme 1 Dehydrocoupling of $NMe₂H·BH₃$ catalysed by complex **1**.

 a Instituto de Investigaciones Químicas (IIQ), Departamento de Química Inorgánica, Centro de Innovación en Ouímica Avanzada (ORFEO-CINCA), CSIC and Universidad de Sevilla, Avda. Américo Vespucio 49, 41092 Sevilla, Spain. E-mail: sconejero@iiq.csic.es; Tel: +34 954489563

 b Laboratorio de Compuestos Organometálicos y Catálisis

⁽Unidad asociada al CSIC), Departamento de Química Orgánica e Inorgánica, Universidad de Oviedo, C/Julián Clavería 8, 33006, Oviedo, Spain

 \dagger Electronic supplementary information (ESI) available: Experimental section, H_2 evolution graphics and X-ray crystallographic data. CCDC 1468913. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c6cc02720b

Scheme 2 Catalytic synthesis of symmetric diaminoboranes

Entry^a	Amineborane	Amine	$%$ Yield ^b (isolated)	TON	TOF (h^{-1})
1	t BuNH ₂ ·BH ₃	t BuNH ₂	> 99(99)	400	2087
2	$NEt_2H·BH_3$	NEt ₂ H	>99(88)	400	2400
3	$(CH2)4NH·BH3$	(CH ₂) ₄ NH	95^{c} (81)	380	193
4	t BuNH ₂ ·BH ₃	NEt2H	> 99(99)	400	3692
5	t BuNH ₂ ·BH ₃	$\rm (CH_2)_4NH$	94(68)	376	376
6	$NEt2H3H3$	$\rm (CH_2)_4NH$	93	372	248
7	NMe ₂ H·BH ₃	BuNH ₂	78^{c} $(71)^{d}$	312	2023
8	NMe ₂ H·BH ₃	NEt,H	76 ^c	316	2257

^a Reaction conditions: CH₂Cl₂, rt. $\frac{b}{b}$ Yields determined by ¹¹B NMR spectroscopy with respect to all diaminoboranes formed. ^c Cyclic dimers $[NR_2BH_2]$ and other unidentified species constitute the remaining reaction products (see the ESI). d Isolated yield corresponds to all the possible products formed.

At the very end of the reaction, the catalyst remained the cyclometalated species 1, but is slowly hydrogenated into complex $[\mathrm{PtH(I^tBu})_2][\mathrm{BAT}^\mathrm{F}],$ 2. Other amine-boranes were tested under identical reaction conditions to form symmetrical diaminoboranes. Diethylamine-borane was also shown to be efficiently dehydrogenated in ca. 10 min in the presence of 1 equiv. of $NEt₂H$ and catalyst 1 (TON = 400, TOF = 2400 h^{-1}). On the other hand, pyrrolidine-borane (CH $_2)_{\rm 4}$ NH·BH $_3$ required longer reaction times (2 h) to be converted into $[(CH₂)₄N]₂BH$ (TON = 380, TOF = 193 h⁻¹). In the latter case, 5% of cyclic dimer $[(CH₂)₄NBH₂]₂$ is observed in the ¹¹B NMR spectra. Bulkier amine-boranes, such as ⁱPr₂NH·BH₃, do not undergo dehydrocoupling under similar reaction conditions, while heating at 60 $^{\circ}$ C for 12 h results in the formation of amino-borane $N^i Pr_2 BH_2$, with no evidence of the corresponding diaminoborane.

Asymmetrical diaminoboranes are challenging substrates to be produced catalytically in a selective manner. Hill et al. have recently reported a method for the preparation of this type of substance using group 2 metal catalysts.^{5b} Although in most of the cases the reaction proceeds with excellent selectivities, long reaction times (24–92 h) and mild heating are usually necessary. Complex 1 has shown very good catalytic behaviour in the formation of unsymmetrical diaminoboranes (Scheme 3). The reaction of t BuNH₂·BH₃ with NEt₂H takes place very fast (6.5 min, TON = 400, TOF = 3692 h^{-1}). According to NMR spectroscopy, borane $(^t$ BuNH)(NEt₂)BH is formed almost exclusively (ca. 4% of $({}^{t}$ BuNH)₂BH is observed by ¹H NMR). If the reaction is carried out in the inverse way, starting from $\text{NE}t_2\text{H-BH}_3$ and ${}^t\text{BuNH}_2$, the process is slower (ca. 20 min), but selectivities are comparable. The same behaviour is observed in the catalytic dehydrocoupling of $(CH_2)_4$ NH·BH₃ and ^tBuNH₂. In this case, the reaction

Scheme 3 Catalytic synthesis of asymmetric diaminoboranes. ^a Signal overlapping in the ${}^{1}H$ and ${}^{11}B({}^{1}H)$ NMR spectra precluded establishing the exact amount of each diaminoborane (see the ESI†).

exhibits good selectivities towards the asymmetric diaminoborane $(^{t}$ BuNH)[(CH₂)₄N]BH (92%), whereas the only symmetric borane detected is $[(CH₂)₄N]₂BH (8%)$, with complete conversion after 2 h at rt. When the reaction is carried out the other way around, that is, starting from t BuNH₂·BH₃ and (CH₂)₄NH, the selectivities are only slightly different (85% of $(^t$ BuNH)[$(CH_2)_4$ N]BH, 15% of $[(CH₂)₄N]₂BH$, although the reaction proceeds in 1 h. The combination of pyrrolidine and diethylamine substrates indicates that the reaction is not selective yielding all possible reaction products (NEt₂)₂BH, $[(CH₂)₄N]₂BH$ and $(NEt₂)[(CH₂)₄N]BH$ in a 1 : 5.4 : 3.1 ratio, respectively, independent of the two possible combinations utilised. A similar behaviour was observed in the reaction of $NMe₂H·BH₃$ with $NEt₂H$, although we could not establish the exact ratio due to signal overlapping in the ¹H NMR (see the ESI†). Nevertheless, good selectivities are produced in the reaction of $NMe₂H·BH₃$ and ${}^{t}BuNH₂$ (Scheme 3). Very interestingly, the reaction of ${}^t{\rm BunH}_2\cdot {\rm BH}_3$ with ${}^{\rm i}{\rm Pr}_2{\rm NH}$ generates diaminoborane (t BuNH)₂BH (s1) and amino-borane ${}^{\dot{i}}$ Pr₂N–BH₂ in ca. 5 min at rt. Since 1 does not catalyse the dehydrocoupling of ${}^{i}Pr_{2}NH \cdot BH_{3}$ at rt, ${}^{i}Pr_{2}N$ -BH₂ is probably formed through the reorganization of amino-borane t BuNH–BH₂ and ${}^{\mathrm{i}}$ Pr₂NH.⁹ Communication

MESCHI DH_y - SHRO (1973-2021)

HESCHI DH_Y - SHRO (1973-2022)

HESCHI DH_Y - SHRO (1974-2022)

HESCHI DHY - S

With regard to the mechanism by which these transformations occur, it has been previously shown that dehydrogenation of amine-boranes $NR_2H·BH_3$ leading to the corresponding aminoboranes $NR₂BH₂$ is a key step.⁵ Previously, we have reported that complex 1 is able to dehydrogenate $NMe₂H·BH₃$ into $NMe₂BH₂$ (that then dimerises) through a distinct mechanism that involves a first step in which the amine-borane interacts with the platinum atom through the BH protons (complex 3a in Scheme 4) followed by nucleophilic addition of free amine present in solution to the activated boron atom.⁷ According to DFT calculations the most stable coordination mode of the BH₃ moiety is η^1 -BH (Shimoi type complex) but, unfortunately, no experimental evidence for this

Scheme 4 Formation of complexes 3a,b.

ChemComm Communication

type of coordination could be provided. Here we further investigate these adducts by carrying out stoichiometric reactions between complex 1 and amine-boranes at 0 $^{\circ}$ C in CD₂Cl₂. ^tBuNH₂ BH₃ reacts with 1 leading to a new species that has been postulated as the η^1 -BH derivative 3b (Scheme 4). The BH₃ group resonates as a broad signal centred at 0.12 ppm that sharpens upon $11B$ broadband decoupling. This signal is shifted upfield from the free ^tBuNH₂·BH₃ (δ 1.36). In addition, $J_{\rm{PtH}}$ of the CH₂-Pt fragment has a value of 90 Hz, that is, 30 Hz smaller than in complex 1 (120 Hz), consistent with the coordination of a ligand trans to the CH_2 moiety.¹⁰ This value is smaller than that observed in dimethylamine-borane 3a (103 Hz) suggesting that the interaction with the platinum atom is stronger. The ¹¹B NMR spectrum shows a signal at -18.2 ppm that is shifted with respect to free amine-borane $(-21.1$ ppm). All these data agree well with the formulation of $3b$ as depicted in Scheme 4 .¹¹ Unfortunately, the instability of 3a and 3b due to their propensity to undergo dehydrocoupling precluded their isolation as pure compounds and further characterization. To avoid this problem, tertiary amine-borane $NMe₃$ ·BH₃ was used. However, no interaction with 1 was observed, probably due to steric constrains, thus allowing the establishment of a correlation between the bulkiness of the amine-borane and its interaction with 1 (stronger interaction: t BuNH₂·BH₃ > NMe₂H·BH₃ > NMe₃·BH₃). Consequently, the less hindered Lewis base stabilised borane $\mathrm{C_6H_5N\cdot BH_3}$ was a judicious choice. This borane reacts with 1 to yield 3c (Scheme 5) quantitatively by NMR spectroscopy. The main feature of 3c is the broad NMR signal centred at 1.25 ppm attributed to the $BH₃$ protons (2.55 ppm in free $C_6H_5N·BH_3$). The $^{11}B(^{1}H)$ spectrum exhibits a signal at -8.9 ppm, nearly 3 ppm down-field shifted with respect to $\rm C_6H_5N\cdot BH_3.$ The apparent coupling constant $^1\!J_{\rm B,H}$ is *ca.* 85 Hz, that is, 12 Hz smaller than in free $C_6H_5N·BH_3$, and compares well with previously described Shimoi-type complexes.^{11,12} The coupling constant of the CH₂–Pt protons with ¹⁹⁵Pt of 93 Hz is similar to that of complex 3**b**. A definite proof of the nature of this compound came from the solid-state structure obtained by X-ray diffraction studies. Colourless crystals suitable for this analysis were obtained by slow diffusion of a concentrated solution of 3c in CH_2Cl_2 into pentane at 0 °C. Fig. 1 depicts an ORTEP-type view of the cation of complex 3c. The complex is fourcoordinated, with two N-heterocyclic carbene units in the expected *trans* arrangement (C(1)–Pt(1)–C(12): 168.38(19)^o) one of which is cyclometalated. The fourth coordination site is occupied by the $C_6H_5N·BH_3$ ligand in which one of the hydride atoms of the $BH₃$ unit bridges the platinum nucleus. The Pt(1)–H(1B) and $H(1B)-B(1)$ bond distances are 1.96(5) and 1.03(2) Å, respectively. Open Communication could be provided. Here we further investigate

these absorbed on earlier care and the ratio and the ratio and the ratio and the ratio and the set of scheme and the ratio and the ratio and the ratio and

 $[BArF]$ tB \widehat{r} -*t*Bu $CD_2Cl_2/$ ri -*t*Bu $f_{\rm BL}$ 3_c

Fig. 1 ORTEP-type view of the cation of complex 3c. Hydrogen atoms, except those of the $BH₃$ unit, have been omitted for clarity (ellipsoids are drawn at 30% probability). Selected bonds (A) and angles $(°)$: Pt1-H1b 1.96(5), Pt1–C5 2.045(5), B1–H1b 1.03(2); Pt1–H1b–B1 147.48(485), C1–Pt1–C12 168.38(19).

The Pt(1)–H(1B)–B(1) angle of $147.48(485)^\circ$ together with the long $Pt(1) \cdot B(1)$ bond distance $(2.8436(5) A)$ indicates a negligible interaction between platinum and boron, and therefore this complex is best described as a η ¹-BH or Shimoi-type complex.^{12,13}

Variable temperature NMR spectroscopy was used to study the dynamic behaviour due to exchange between bridging and terminal BH protons. The resonance for the $BH₃$ protons collapses into the baseline at ca. 223 K. Upon cooling to 203 K, a new signal of relative integral 2 appears at 2.47 ppm, in the expected region for terminal BH protons. Nevertheless, at this temperature, the signal corresponding to the Pt–H–B proton is not discernible.¹⁴ Further cooling to 183 K provided evidence for a very broad resonance at $ca. -4$ ppm (approx. integral value of 1), with the terminal BH protons resonating at 2.44 ppm. With respect to the 11 B NMR spectra, a very broad signal centred at δ -12.2 is detected at this temperature.

For comparison purposes, we have analysed the interaction of $C_6H_5N·BH_3$ with the hydrogenated form of complex 1, the hydride derivative $[\mathrm{PtH(I^tBu)}_2][\mathrm{BAT}^F]$, 2.¹⁵ No evidence of interaction between 2 and $C_6H_5N·BH_3$ is observed at 298 K. At this temperature, the BH₃ protons resonate at δ 2.49, whereas the platinum-hydride signal appears at δ -25.47 exhibiting a $J_{\text{Pt,H}}$ of 2550 Hz, only marginally different from that of complex 2 in the absence of $C_6H_3N·BH_3$ ($J_{Pt,H}$ = 2564 Hz).¹⁵ However, the ¹H₃ MMP spectra at temperatures below 222 K indicate the 1 H NMR spectra at temperatures below 233 K indicate the presence of a new species, 4 (Scheme 6), coexisting with complex 2, in a 0.3 to 1 ratio (4 : 2). This compound is characterised by a signal in the hydride region at -21.85 ppm showing a reduced coupling constant with ¹⁹⁵Pt ($J_{Pt,H}$ = 1920 Hz) and another broad signal at 0.28 ppm (3H relative integral) that sharpens upon $11B$ decoupling. As the temperature decreases, the concentration of 4 increases at the expense of 2, reaching a maximum at 188 K (4:2 ratio, *ca.* 4:1). The ${}^{11}B{^1H}$ NMR spectra at all temperatures exhibit a single very broad signal at -12.3 ppm that does not show splitting upon coupling to 1 H below 208 K. Interestingly, even at 188 K the $BH₃$ group still shows fast exchange between terminal and bridging hydrogens. The dif-Scheme 5 Synthesis of complex 3c. **Ferent behaviour of compounds 1 and 2 toward the interaction**

Scheme 6 Equilibrium between complexes 2 and 4 in the presence of $C_6H_5N·BH_3.$

with $\rm{C_6H_5N\cdot BH_3}$ can be easily rationalised in terms of the different steric protection that the cyclometalated I^tBu ligand exerts on the platinum atom compared to its non-cyclometalated form. When the I^tBu ligand is cyclometalated the ${}^t\!{\rm Bu}$ group is tilted 16 away from the metal centre favouring the interaction with amine-boranes.

Once the amine-borane binds the metal in a Shimoi-type fashion, dehydrogenation leading to amino-boranes, NR_2-BH_2 , takes place through a mechanism that involves the intermediacy of boronium cations $(NHR_2)_2BH_2^+$ and the neutral platinum hydride [PtH(I^tBu')(I^tBu)], 5.⁷ It must be noted that we have not observed an interaction between complex 1 and amino-borane $NMe₂BH₂$ (either using cyclic dimer $[NMe₂BH₂]$ as a precursor¹⁷ or during monitoring the dehydrogenation reaction of 1 and $\text{NMe}_{2}\text{H}\cdot\text{BH}_{3}$ at 0 $^{\circ}$ C).⁷ At this point, we do not have further information on the mechanism by which the amino-borane is converted into diaminoboranes and the role of complex 1 in the process. Communication

Published on $\frac{1}{2}$ American Commons Article is licensed under a Creative Commons Article is an access Article is licensed under a Creative Commons Article is a set of the set of the Section Article is a

In summary, the coordinatively unsaturated $Pf(n)$ complex 1 is a very efficient catalyst for the synthesis of diaminoboranes achieving TON and TOF values of 400 and 3692 $\rm h^{-1},$ respectively, the highest reported to date. The process takes place through the initial coordination of the BH protons of amine-boranes to the platinum centre in an end-on mode (Shimoi type) that was demonstrated crystallographically in the pyridine \cdot BH $_3$ adduct 3c. Ongoing efforts are geared towards unveiling the mechanism by which amino-boranes are transformed into diaminoboranes.

Financial support (FEDER contribution) from the MINECO (Projects CTQ2013-45011-P and CTQ2014-51912-REDC) and the Junta de Andalucía (Project FQM-2126) is gratefully acknowledged.

Notes and references

- 1 (a) H. C. Johnson, T. N. Hooper and A. S. Weller, Top. Organomet. Chem., 2015, 49, 153; (b) E. M. Leitao, T. Jurka and I. Manners, Nat. Chem., 2013, 5, 817; (c) N. E. Stubbs, A. P. M. Robertson, E. M. Leitao and I. Manners, J. Organomet. Chem., 2013, 730, 84; (d) A. S. John, K. I. Goldberg and D. M. Heinekey, Top. Organomet. Chem., 2013, 40, 271.
- 2 (a) H. C. Johnson and A. S. Weller, Angew. Chem., Int. Ed., 2015, 54, 10173; (b) M. A. Esteruelas, A. M. López, M. Mora and E. Oñate, ACS Catal., 2015, 5, 187; (c) A. G. Algarra, L. J. Sewell, H. C. Johnson, S. A. Macgregor and A. S. Weller, Dalton Trans., 2014, 43, 11118;

(d) S. Muhammad, S. Moncho, E. N. Brothers and A. A. Bengali, Chem. Commun., 2014, 50, 5874; (e) L. J. Sewell, G. C. Lloyd-Jones and A. S. Weller, J. Am. Chem. Soc., 2012, 134, 3598; (f) C. Y. Tang, N. Phillips, M. J. Kelly and S. Aldridge, Chem. Commun., 2012, 48, 11999; (g) R. Kumar and B. R. Jagirdar, Inorg. Chem., 2013, 22, 28; (h) H. C. Johnson, A. P. M. Robertson, A. B. Chaplin, L. J. Sewell, A. L. Thompson, M. F. Haddow, I. Manners and A. S. Weller, J. Am. Chem. Soc., 2011, 133, 11076; (i) R. Dallanegra, A. P. M. Robertson, A. B. Chaplin, I. Manners and A. S. Weller, Chem. Commun., 2011, 47, 3763; (j) G. Alcaraz and S. Sabo-Etienne, Angew. Chem., Int. Ed., 2010, 49, 7170; (k) C. Y. Tang, A. L. Thompson and S. Aldridge, J. Am. Chem. Soc., 2010, 132, 10578; (l) T. M. Douglas, A. B. Chaplin, A. S. Weller, X. Yang and M. B. Hall, J. Am. Chem. Soc., 2009, 131, 15440.

- 3 (a) A. Staubitz, A. P. M. Robertson and I. Manners, Chem. Rev., 2010, 110, 4079; (b) N. C. Smythe and J. C. Gordon, Eur. J. Inorg. Chem., 2010, 509; (c) C. W. Hamilton, R. T. Baker, A. Staubitz and I. Manners, Chem. Soc. Rev., 2009, 38, 279.
- 4 H. C. Johnson, C. L. McMullin, S. D. Pike, S. A. Macgregor and A. S. Weller, Angew. Chem., Int. Ed., 2013, 52, 9776.
- 5 (a) C. J. Wallis, G. Alcaraz, A. S. Petit, A. I. Poblador-Bahamonde, E. Clot, C. Bijani, L. Vendier and S. Sabo-Etienne, Chem. – Eur. J., 2015, 21, 13080; (b) P. Bellham, M. S. Hill, G. Kociok-Köhn and D. J. Liptrot, Chem. Commun., 2013, 49, 1960; (c) C. J. Wallis, H. Dyer, L. Vendier, G. Alcaraz and S. Sabo-Etienne, Angew. Chem., Int. Ed., 2012, 51, 3646; (d) It has been reported that aromatic amine-boranes can produce diaminoboranes without a catalyst: H. Helten, A. P. M. Robertson, A. Staubitz, J. R. Vance, M. F. Haddow and I. Manners, Chem. – Eur. J., 2012, 18, 4665.
- 6 O. Rivada-Wheelaghan, B. Donnadieu, C. Maya and S. Conejero, Chem. – Eur. J., 2010, 16, 10323.
- 7 M. Roselló-Merino, J. López-Serrano and S. Conejero, J. Am. Chem. Soc., 2013, 135, 10910.
- 8 (a) P. Bellham, M. S. Hill and G. Kociok-Köhn, Organometallics, 2014, 33, 5716; (b) Y. Kawano, M. Uruichi, M. Shimoi, S. Taki, T. Kawaguchi, T. Kakizawa and H. Ogino, J. Am. Chem. Soc., 2009, 131, 14946.
- 9 A. P. M. Robertson, E. M. Leitao and I. Manners, J. Am. Chem. Soc., 2011, 133, 19322.
- 10 M. A. Ortuño, S. Conejero and A. Lledós, Beilstein J. Org. Chem., 2013, 9, 1352.
- 11 G. Alcaraz and S. Sabo-Etienne, Coord. Chem. Rev., 2008, 252, 2395.
- 12 (a) R. Dallanegra, A. B. Chaplin, J. Tsim and A. S. Weller, Chem. Commun., 2010, 46, 3092; (b) M. Shimoi, S. Nagai, M. Ichikawa, Y. Kawano, K. Katoh, M. Uruichi and H. Ogino, J. Am. Chem. Soc., 1999, 121, 11704.
- 13 (a) A. E. W. Ledger, C. E. Ellul, M. F. Mahon, J. M. J. Williams and M. K. Whittlesey, Chem. – Eur. J., 2011, 17, 8704; (b) R. Dallanegra, A. B. Chaplin and A. S. Weller, Angew. Chem., Int. Ed., 2009, 48, 6875; (c) Y. Kawano, K. Yamaguchi, S. Miyake, T. Kakizawa and M. Shimoi, Chem. – Eur. J., 2007, 13, 6920; (d) Y. Kawano, M. Hashiva and M. Shimoi, Organometallics, 2006, 25, 4420; (e) N. Merle, G. Koicok-Köhn, M. F. Mahon, C. G. Frost, G. D. Ruggerio, A. S. Weller and M. C. Willis, Dalton Trans., 2004, 3883; (f) T. Yasue, Y. Kawano and M. Shimoi, Angew. Chem., Int. Ed., 2003, 42, 1727.
- 14 A similar behavior has been observed in a ruthenium complex. See ref. 13a.
- 15 (a) M. A. Ortuño, P. Vidossich, S. Conejero and A. Lledós, Angew. Chem., Int. Ed., 2014, 53, 14158; (b) O. Rivada-Wheelaghan, M. Roselló-Merino, M. A. Ortuño, P. Vidossich, E. Gutiérrez-Puebla, A. Lledós and S. Conejero, Inorg. Chem., 2014, 53, 4257.
- 16 J.-N. Luy, S. A. Hauser, A. B. Chaplin and R. Tonner, Organometallics, 2015, 34, 5099.
- 17 G. Bénac-Lestrille, U. Helmstedt, L. Vendier, G. Alcaraz, E. Clot and S. Sabo-Etienne, Inorg. Chem., 2011, 50, 11039.