



Cite this: *Chem. Commun.*, 2016, 52, 10632

Received 16th May 2016,
Accepted 1st August 2016

DOI: 10.1039/c6cc04132a

www.rsc.org/chemcomm

Catalytic enantio- and diastereoselective Mannich reaction of α -substituted isocyanoacetates and ketimines†

Raquel de la Campa, Adam D. Gammack Yamagata, Irene Ortín, Allegra Franchino, Amber L. Thompson, Barbara Odell and Darren J. Dixon*


The highly diastereo- and enantioselective Mannich addition/cyclisation reaction of α -substituted isocyanoacetate ester pronucleophiles and (hetero)aryl and alkyl methyl ketone-derived ketimines using a silver acetate and a cinchona-derived amino phosphine binary catalyst system is reported.

The Mannich reaction is a powerful tool for the creation of carbon–carbon bonds¹ and since its discovery in the early 20th century great progress in the development of catalytic enantioselective variants has been made.² To date, the majority of studies have employed aldimines for the enantioselective preparation of amine derivatives possessing tertiary stereogenic centres.^{3,4} However recent efforts have focussed on the development of new catalytic Mannich reactions with ketimines for the stereoselective production of amine derivatives possessing tetrasubstituted stereogenic centres,⁵ the construction of which remains an important challenge in the field.⁶ To this end, our group recently developed a catalytic enantioselective Mannich reaction of isocyanoacetate esters and ketimines to generate trisubstituted chiral imidazolines possessing vicinal quaternary and tertiary stereogenic carbon atoms.⁷ Under the control of a binary catalyst system comprising silver oxide and a cinchona alkaloid-derived amino phosphine, impressive reactivity and high enantio- and diastereocontrol were achieved.⁸ As well as being important building blocks for the synthesis of α,β -diamino acids,⁹ substituted imidazolines, such as those produced by this chemistry, can be identified as the core of biologically active compounds.¹⁰

We recognised that the catalytic enantioselective Mannich reaction of α -substituted isocyanoacetate pronucleophiles with ketimines could provide a synthetically powerful approach for the enantioselective synthesis of imidazolines possessing two

vicinal tetrasubstituted stereocentres.¹¹ To this end, and in a further development of our ongoing research program, we wish to report our efforts leading to a catalytic enantioselective ketimine Mannich reaction of α -substituted isocyanoacetate esters affording chiral imidazolines with two contiguous fully substituted carbon stereocentres, promoted by a catalytic system made from a cinchona-derived amino phosphine ligand and a silver(i) salt (Scheme 1).

Initially we selected acetophenone-derived *N*-diphenylphosphinoyl ketimine **3a** as our model electrophile and investigated a range of 2-isocyanopropanoate ester pronucleophiles **2** using our previously optimised^{7a} catalyst system comprising quinine-derived amino phosphine **1a** (20 mol%) and Ag₂O (5 mol%) in ethyl acetate at $-20\text{ }^{\circ}\text{C}$ (Table 1). Methyl 2-isocyanopropanoate **2a** was first investigated as the pronucleophile. Very pleasingly after 60 hours, *N*-DPP protected *trans*-imidazoline **4a** was afforded as a single diastereomer. To allow full characterisation and determination of enantiomeric excess (ee) *via* HPLC analysis, DPP group removal using 1 M HCl was found to be necessary and imidazoline **5a** was afforded in 59% yield over the two steps, in 89% ee (entry 1). With reasonable reactivity and stereoselectivity identified, the performance of more bulky 2-isocyanopropanoate esters – specifically iso-propyl (**2b**), *tert*-butyl (**2c**), benzyl (**2d**) and benzhydryl (**2e**) 2-isocyanopropanoates – was studied (entries 2–5). With *tert*-butyl ester **2c** the high enantioselectivity (89% ee) was maintained and the reaction yield was slightly improved to 63% (entry 3),




Scheme 1 Enantioselective construction of chiral imidazolines with two contiguous tetrasubstituted stereocentres.

Chemistry Research Laboratory, Department of Chemistry, University of Oxford, Mansfield Road, Oxford, OX1 3TA, UK. E-mail: darren.dixon@chem.ox.ac.uk

† Electronic supplementary information (ESI) available: Procedures, characterisation data, copies of NMR spectra, HPLC traces, X-ray data and CIF file. CCDC 1493438. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c6cc04132a





Scheme 2 Application to the synthesis of fully substituted α,β -diamino acids.

substituent in the α -position in similarly high ee and yield (entries 12–15). The *tert*-butyl 2-isocyanoacetate **2g** also performed well in the reaction to give products **5t–5w** in excellent yield and ee (entries 16–19).¹⁴ NOe NMR experiments established the relative *trans* stereochemical configuration of products **5f**, **5o**, **5q** and **5v**. Absolute configuration was determined by comparison of the specific rotation of imidazoline **5y** with that reported in the literature.^{5c}

As a demonstration of the synthetic utility of this newly established method, product **5c** was hydrolysed to afford α,β -diamino acid **6c** (Scheme 2). Such compounds are difficult to access in enantioenriched form by other methods.⁹

To gain insight into the importance of the various functional groups possessed by amino phosphines **1**, a series of control experiments using modified ligand structures (**1d–1g**) were performed (Table 3). Diastereomeric ligand **1d** gave imidazoline **5c** with a large decrease in enantioselectivity (entry 2). Ligand **1e**, lacking the phosphine moiety, was significantly worse: product **5c** was obtained in low yield with poor enantioselectivity (entry 3). Ligand **1f** with an ester linker gave imidazoline **5c** in slightly decreased yield and enantioselectivity (entry 4). Finally ligand **1g** with a *N*-methyl group on the amide promoted the reaction with a similar yield to **1a** but much lower levels of enantioselectivity (entry 5).

In order to better understand the results from the catalyst modification studies and to determine how ligand **1a** binds to AgOAc, a crystal of complex **1a**-AgOAc was grown (Fig. 1).¹⁵ Single crystal X-ray diffraction determined that in the solid




Fig. 1 Structure of **1a**-AgOAc from single crystal X-ray diffraction studies showing the intramolecular H-bond (H...O = 1.896(3) Å; N...O = 2.752(4) Å). Displacement ellipsoids drawn at 30% probability; solvent and selected hydrogen atoms omitted for clarity.

state **1a** acts as a *P,N*-bidentate ligand, binding silver through both the phosphine and the quinuclidine N atom to create a 9-membered macrocycle. The other groups coordinating to silver are the acetate and a quinoline unit belonging to a symmetry related molecule of **1a** leading to the formation of polymeric chains in the solid state. The four-coordinate silver adopts a distorted tetrahedral geometry. An intramolecular hydrogen bond between the acetate group on silver and the transannular amide N–H at 1.896(3) Å is present with a N...O distance of 2.752(4) Å. The π system of the amide group is oriented towards the silver with a distance of 2.9571(18) Å between the amide nitrogen and the silver centre suggesting a possible interaction. Such an interaction would transfer charge resulting in acidification of the amide NH thus enhancing its H-bond donor ability.

³¹P and ¹H NMR studies of a 1 : 1 ratio of amino phosphine **1a** and AgOAc at –20 °C were also performed (see ESI† for full NMR studies). The ³¹P NMR experiments showed a quantitative binding of silver(I) in solution and ¹H NMR analysis of the same mixture revealed a large downfield shift (approximately 3.4 ppm relative to that of the free ligand) of the amide N–H consistent with presence of the same H-bonding interaction. Additionally, the protons adjacent to the quinuclidine nitrogen were also shifted downfield, consistent with coordination of the quinuclidine nitrogen to silver. In the solution phase there is no strong evidence that the quinoline N atom binds the Ag centre; the change of chemical shift for the proton adjacent to the quinoline nitrogen is only 0.07 ppm. When methyl isocyanoacetate is added to the solution of **1a** and AgOAc, the CH₂ group appears as a characteristic ABq system, providing evidence for complexation of the isonitrile to silver.

Building on the results of experiments using modified ligand structures (Table 3) and the structural data obtained from both the X-ray and NMR studies we propose a transition structure that rationalises the enantio- and diastereoselectivity observed in this Mannich reaction (Fig. 2). The N atom of the (*E*)-configured ketimine is postulated to make a key hydrogen bond to the amide N–H, which both orientates the electrophile and activates it towards nucleophilic addition. The aromatic ring of the H-bonded ketimine is oriented away from the quinuclidine and quinoline units on steric grounds and C–H/ π -interactions¹⁶

Table 3 Variation on ligand structure

Entry	1	Yield ^a	ee
1	1a	90	89
2	1d	89	20
3	1e	23	5
4	1f	75	72
5	1g	86	27

^a Yield over two steps for product **5c** (single diastereomer) after FCC.






Fig. 2 Proposed transition structure.

between it and the aromatic ring of the carboxamide could provide additional binding/organisation, possibly accounting for the high enantiofacial selectivity observed using this catalyst.^{7a,8} The ester group of the isocyanoacetate is oriented away from the aromatic group on the ketimine on steric grounds. Subsequent to the enantio- and diastereodetermining C–C bond forming step, ring closure and protonolysis would yield the *N*-DPP protected imidazoline with the observed stereochemical configuration. Satisfyingly, this transition state model, with its multipoint binding interactions, would also rationalise the stereochemical outcomes of the previously reported reactions with aldehydes^{8a} and ketones.^{8b}

In conclusion, we have developed a diastereo- and enantioselective Mannich reaction/cyclisation of α -substituted isocyanoacetates and ketimines catalysed by silver acetate and a cinchona-derived amino phosphine. The reaction scope is broad and includes substituted aryl, heteroaryl and alkyl methyl ketimines as well as isocyanoacetates with linear alkyl chains in the α -position. Hydrolysis of the product imidazolines provides access to fully substituted α,β -diamino acids. Information arising from X-ray and NMR studies, as well as control studies on the ligand, have allowed a detailed transition state model to be postulated.

The authors thank the EPSRC for a leadership fellowship to D. J. D., Merck, Sharp & Dohme (Hoddesdon, UK) for a CASE award to A. D. G. Y., and the EC for IEF to R. C. (PIEF-GA-2012-329689), IEF to I. O. (PIEF-GA-2010-275788) and PhD fellowship to A. F. (FP7/2007-2013, REA grant agreement 316955).

Notes and references

- 1 T. Akiyama, in *Comprehensive Organic Synthesis*, ed. P. Knochel and G. A. Molander, Elsevier, Oxford, 2nd edn, 2014, vol. 2, p. 629.
- 2 (a) S. Kobayashi and H. Ishitani, *Chem. Rev.*, 1999, **99**, 1069; (b) A. Cordova, *Acc. Chem. Res.*, 2004, **37**, 102; (c) N. Z. Burns and E. N. Jacobsen, in *Science of Synthesis, Stereoselective Synthesis*, ed. J. G. De Vries, G. A. Molander and P. A. Evans, Thieme, Stuttgart, 2011, vol. 2, p. 785.
- 3 Enantioselective Mannich reactions of aldimines: (a) K. Juhl, N. Gathergood and K. A. Jørgensen, *Angew. Chem., Int. Ed.*, 2001, **40**, 2995; (b) L. Bernardi, A. S. Gothelf, R. G. Hazell and K. A. Jørgensen, *J. Org. Chem.*, 2003, **68**, 2583; (c) T. Ooi, M. Kameda, J.-I. Fujii and K. Maruoka, *Org. Lett.*, 2004, **6**, 2397; (d) A. Okada,

- T. Shibuguchi, T. Ohshima, H. Masu, K. Yamaguchi and M. Shibasaki, *Angew. Chem., Int. Ed.*, 2005, **44**, 4564; (e) N. S. Chowdari, M. Ahmad, K. Albertshofer, F. Tanaka and C. F. Barbas III, *Org. Lett.*, 2006, **8**, 2839; (f) S. Kobayashi, R. Yazaki, K. Seki and Y. Yamashita, *Angew. Chem., Int. Ed.*, 2008, **47**, 5613; (g) J. Hernández-Toribio, R. G. Arrayás and J. C. Carretero, *J. Am. Chem. Soc.*, 2008, **130**, 16150; (h) D. Shang, Y. Liu, X. Zhou, X. Liu and X. Feng, *Chem. – Eur. J.*, 2009, **15**, 3678; (i) J. Hernández-Toribio, R. G. Arrayás and J. C. Carretero, *Chem. – Eur. J.*, 2010, **16**, 1153; (j) G. Liang, M.-C. Tong, H. Tao and C.-J. Wang, *Adv. Synth. Catal.*, 2010, **352**, 1851; (k) E. Hernando, R. G. Arrayás and J. C. Carretero, *Chem. Commun.*, 2012, **48**, 9622.
- 4 Enantioselective Mannich reactions of isocyanoacetate esters and aldimines: (a) X.-T. Zhou, Y.-R. Lin, L.-X. Dai, J. Sun, L.-J. Xia and M.-H. Tang, *J. Org. Chem.*, 1999, **64**, 1331; (b) X.-T. Zhou, Y.-R. Lin and L.-X. Dai, *Tetrahedron: Asymmetry*, 1999, **10**, 855; (c) J. Aydin, A. Rydén and K. J. Szabó, *Tetrahedron: Asymmetry*, 2008, **19**, 1867; (d) Z.-W. Zhang, G. Lu, M.-M. Chen, N. Lin, Y.-B. Li, T. Hayashi and A. S. C. Chan, *Tetrahedron: Asymmetry*, 2010, **21**, 1715; (e) S. Nakamura, Y. Maeno, M. Ohara, A. Yamamura, Y. Funanashi and N. Shibata, *Org. Lett.*, 2012, **14**, 2960; (f) P.-L. Shao, J.-Y. Liao, Y. A. Ho and Y. Zhao, *Angew. Chem., Int. Ed.*, 2014, **53**, 5435.
- 5 Enantioselective Mannich reactions of ketimines: (a) Y. Du, L.-W. Xu, Y. Shimizu, K. Oisaki, M. Kanai and M. Shibasaki, *J. Am. Chem. Soc.*, 2008, **130**, 16146; (b) L. C. Wieland, E. M. Vieira, M. L. Snapper and A. H. Hoveyda, *J. Am. Chem. Soc.*, 2009, **131**, 570; (c) G. Lu, T. Yoshino, H. Morimoto, S. Matsunaga and M. Shibasaki, *Angew. Chem., Int. Ed.*, 2011, **50**, 4382; (d) M. Hayashi, M. Sano, Y. Funahashi and S. Nakamura, *Angew. Chem., Int. Ed.*, 2013, **52**, 5557; (e) M. G. Núñez, A. J. M. Farley and D. J. Dixon, *J. Am. Chem. Soc.*, 2013, **135**, 16348; (f) Y.-Q. Wang, Y. Zhang, K. Pan, J. You and J. Zhao, *Adv. Synth. Catal.*, 2013, **355**, 3381; (g) M.-X. Zhao, H.-L. Bi, R.-H. Jiang, X.-W. Xu and M. Shi, *Org. Lett.*, 2014, **16**, 4566; (h) S. Zhang, L. Li, Y. Hu, Z. Zha, Z. Wang and T.-P. Loh, *Org. Lett.*, 2015, **17**, 1050; (i) S. Lin, Y. Kawato, N. Kumagai and M. Shibasaki, *Angew. Chem., Int. Ed.*, 2015, **54**, 5183; (j) S. Nakamura, R. Yamaji and M. Hayashi, *Chem. – Eur. J.*, 2015, **21**, 9615; (k) X. Liu, J. Zhang, L. Zhao, S. Ma, D. Yang, W. Yan and R. Wang, *J. Org. Chem.*, 2015, **80**, 12651; (l) S. Lin, N. Kumagai and M. Shibasaki, *Chem. – Eur. J.*, 2016, **22**, 3296; (m) T. Takeda, A. Kondoh and M. Terada, *Angew. Chem., Int. Ed.*, 2016, **55**, 4734.
- 6 J. Christoffers and A. Mann, *Angew. Chem., Int. Ed.*, 2001, **40**, 4591.
- 7 (a) I. Ortín and D. J. Dixon, *Angew. Chem., Int. Ed.*, 2014, **53**, 3462; (b) M. Hayashi, M. Iwanaga, N. Shiomi, D. Nakane, H. Masuda and S. Nakamura, *Angew. Chem., Int. Ed.*, 2014, **53**, 8411.
- 8 (a) F. Sladojevich, A. Trabocchi, A. Guarna and D. J. Dixon, *J. Am. Chem. Soc.*, 2011, **133**, 1710; (b) R. de la Campa, I. Ortín and D. J. Dixon, *Angew. Chem., Int. Ed.*, 2015, **54**, 4895.
- 9 A. Viso, R. Fernández de la Pradilla, M. Tortosa, A. García and A. Flores, *Chem. Rev.*, 2011, **111**, PR1.
- 10 H. Liu and D.-M. Du, *Adv. Synth. Catal.*, 2009, **351**, 489.
- 11 During the preparation of the present manuscript, the following article was published online: S. Nakamura, R. Yamaji and M. Iwanaga, *Chem. Commun.*, 2016, **52**, 7462.
- 12 Q.-A. Chen, D.-S. Wang and Y.-G. Zhou, *Chem. Commun.*, 2010, **46**, 4043.
- 13 Reaction between **2c** and 4'-bromopropiophenone-derived ketimine, followed by removal of the DPP group, gave the corresponding imidazoline in 62% overall yield and 70% ee.
- 14 Reaction between **3a** and valine-derived *tert*-butyl isocyanoacetate (isopropyl group at the α -position), followed by DPP deprotection, gave the corresponding imidazoline in 29% overall yield (8 days reaction time) and 98% ee.
- 15 CCDC 1493438.
- 16 S. Tsuzuki and A. Fujii, *Phys. Chem. Chem. Phys.*, 2008, **10**, 2584.

