



Stable bromoallene oxides†

Cite this: *Chem. Commun.*, 2016, 52, 11219

D. Christopher Braddock,* Areeb Mahtey, Henry S. Rzepa and Andrew J. P. White


Received 3rd August 2016,
Accepted 22nd August 2016

DOI: 10.1039/c6cc06395k

www.rsc.org/chemcomm

The first stable bromoallene oxides were obtained by the DMDO epoxidation of 1-bromo-1,3-di-*tert*-alkyllallenes, producing the first crystalline allene oxide of any kind. The epoxidations are regio-selective for the bromine-bearing $\Delta^{1,2}$ alkene, and also face selective producing single diastereomer *E*-olefin products.

The allene oxide functional group and its valence tautomeric cyclopropanone and oxyallyl cation forms are of significant interest.^{1,2} The first room-temperature stable allene oxide was reported in 1968 by peracid epoxidation of 1,3-di-*tert*-butyllallene **1** to give epoxide **2**.³ It was subsequently converted into its corresponding isomeric cyclopropanone **3** on heating thereby establishing their relative thermodynamic stabilities. However, despite this early success, only a handful of further allene oxides have been successfully characterised.^{4–10} Nevertheless, these studies have helped to establish that stable allene oxides can be obtained when (i) the epoxide is kinetically stabilised by steric protection against S_N2 attack, and (ii) when cyclopropanone formation *via* the presumed intermediacy of an oxyallyl cation is non-favoured.^{1c}



We have recently reported¹¹ that DMDO epoxidation of 3-*n*-alkyl-1-bromo-allenes **4** give rise directly to α,β -unsaturated carboxylic acids **5** by presumed spontaneous rearrangement of bromo-allene oxides – an unprecedented functional group in the literature¹² – to bromocyclopropanones and subsequent Favorskii rearrangement.


Department of Chemistry, Imperial College London, London, SW7 2AZ, UK.

E-mail: c.braddock@imperial.ac.uk; Fax: +44 (0)2075945805;

Tel: +44 (0)2075945772


† Electronic supplementary information (ESI) available: Experimental procedures and characterising data for 7–9, 11–24; copies of ¹H and ¹³C NMR spectra for all compounds including NOE, DEPT-135, correlation spectroscopy and bromide induced isotopic shifts where appropriate. CCDC 1485934 (9) and 1485935 (19). For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c6cc06395k

In that work, an analysis of the mechanism revealed that epoxidation of either olefin of the bromoallene would lead to the same product and so the site of initial epoxidation could not be identified. With the intention of identifying the preferred position of epoxidation in bromoallenes, we now report the first stable bromoallene oxides.




With the above literature precedents in mind, we selected known 1-bromo-1,3-di-*tert*-butyllallene **6** – reported as a colourless liquid – as a suitable substrate for epoxidation.¹³ However, in our hands it proved to be inseparable from its propargylic bromide isomer, was found to be somewhat volatile, and consequently epoxidation experiments proved inconclusive. Instead, we targeted novel adamantyl-containing bromoallenes **7–9**, which proved to overcome these difficulties (*vide infra*).

Bromoallene **7** was prepared by the initial lithiation of terminal alkyne **10** and addition to adamantancarboxaldehyde§ to give propargylic alcohol **11** (Scheme 1). Temperature control in this



Scheme 1 Synthesis of bromoallenes **7–9** (Ad = 1-adamantyl).





Scheme 2 Epoxidation of bromoallenes 7–9.




Fig. 1 X-ray crystal structure of bromoallene oxide 19.

transformation proved critical, otherwise significant quantities of ketone 12 and 1-adamantanemethanol were formed.¶¶ Using the conditions of Corey,¹³ alcohol 11 was transformed into bromoallene 7 along with the expected, but now separable, propargylic bromide 13.** Next, the action of the carbon tetrachloride–triphenylphosphine reagent combination¹⁴ on adamantancarboxaldehyde as a α,α,α -trisubstituted aldehyde provided ready access to vinyl dichloride 14¹⁵ free from any complicating dichloroalkane formation.¹⁶ Corey–Fuchs type acetylide anion formation,¹⁷ and *in situ* trapping¹⁸ with pivaldehyde and adamantancarboxaldehyde gave propargylic alcohols 15 and 17 respectively in good yields.†† Bromoallenes 8 and 9 were subsequently obtained by subsequent reaction with thionyl bromide along with the corresponding expected, and again separable, propargylic bromides 16 and 18.‡‡

The incorporation of adamantyl groupings rendered the three bromoallenes 7–9 to be solid, bromoallene 9 proved to be crystalline, and its structure was solved by X-ray crystallography.‡‡‡ With these compounds in hand, attention turned to epoxidation.¶¶¶

In the event, epoxidation of each of bromoallenes 7–9 with DMDO¹⁹ provided monoepoxidation products as *E*-1-bromoallene oxides 19–21|| as single diastereoisomers respectively, along with bromodiketones 22–24,*** and recovered bromoallene starting materials after column chromatography (Scheme 2).††† Importantly, under the conditions of the experiment and the isolation and purification regime, there was no evidence for the formation of any corresponding α,β -unsaturated carboxylic acids *via* intersecting bromoallene oxide–bromocyclopropanone–Favorskii rearrangement.¹¹ These experiments therefore establish that (i) bromoallene oxides of this type are kinetically stabilised with respect to rearrangement by the bulky alkyl substituents; (ii) for these bromoallenes, epoxidation occurs preferentially at the bromine-bearing $\Delta^{1,2}$ alkene and (iii) epoxidation occurs with complete face selectivity away from the alkyl grouping at C-3 of the allene.‡‡‡ Moreover, the uniformity of the product distributions obtained with the different bromoallenes 7–9 show that the *tert*-butyl and adamantyl groups behave as effectively identical and interchangeable substituents in this situation, but with the advantage that the incorporation of an adamantyl group increases the prospect of crystallinity. Indeed, bromoallene oxide 19 proved to be crystalline and its structure was determined by X-ray crystallography (Fig. 1). To the best of our knowledge this is the first X-ray crystallographic structural elucidation of any allene oxide.§§§²⁰

We probed the origins of the observed regioselectivity and diastereoselectivity using DFT theory (ω B97XD/TZVP, QZVP/PCM = acetone as continuum solvent).²¹ An NBO analysis on

representative reactant bromoallene 6 reveals the $\Delta^{2,3}$ localized occupied π -orbital is higher in energy (by 0.014 Hartree) than the $\Delta^{1,2}$ orbital, suggesting that the intrinsic electronic nucleophilic reactivity of the allene is at the 2,3 position. However, transition state models of the reaction with DMDO as electrophile reveals the 1,2 isomers are 1.2 (QZVP) or 1.4 (TZVP) kcal mol⁻¹ lower in ΔG_{298}^\ddagger for bromoallene 6 and 1.2 (TZVP) for bromoallene 9. For the simplest possible bromoallene (H₂C=C=CHBr, not shown) the 2,3 transition state at the π -face *anti*-periplanar to the Br is favoured by 0.4 kcal mol⁻¹ over the 1,2 isomer and by 1.3 kcal mol⁻¹ over 2,3-*syn*-periplanar attack (TZVP). Taken together, these calculations reveal that although the bromine atom is inherently deactivating for the $\Delta^{1,2}$ olefin, it also acts to sterically protect the *syn*-periplanar face of the $\Delta^{2,3}$ olefin. For bromoallenes 7–9, since the bulky alkyl group on C-1 protects the other face of $\Delta^{2,3}$ olefin and the C-3 alkyl group inhibits *Z*-1-bromoallene oxide formation, $\Delta^{1,2}$ attack leading to the *E*-1-bromoallene oxides 19–21 emerges as the most facile.

In conclusion, we have demonstrated that stable bromoallene oxides can be obtained by DMDO epoxidation of 1-bromo-1,3-di-*tert*-alkylallenes, and the first X-ray crystal structural elucidation of any allene oxide has thereby been accomplished. The epoxidation for these bromoallenes is selective for the bromine-bearing $\Delta^{1,2}$ alkene, and with exclusive face selectivity to produce *E*-1-bromoallene oxides.¶¶¶ The thermal stability of these bromoallene oxides is the subject of current investigations and will be reported in due course.

We thank Mr Jaren Soo for preliminary investigations with compound 6.

Notes and references

‡ Interestingly, a definitive structure for allene oxide 2 was not given in the original report and two geometrical isomeric structures were proffered. Our results (*vide infra*) support this to be the *E*-allene oxide as drawn.

§ Readily prepared from 1-adamantanemethanol by PCC oxidation.‡

¶ See ESI† for experimental details. See also ref. 21.

|| We invoke an unprecedented irreversible hydride transfer from the initially formed secondary alkoxide AdCH(C \equiv C^{*t*}Bu)OLi to still unreacted AdCHO in a concerted six-membered TS to explain this observation. In this process steric compression of the adamantyl-bearing sp³ centre is relieved, and the formation of conjugated ynone 12 acts as a thermodynamic driving force. Formation of the lithium alkoxide from alcohol 11 by the action of *n*-BuLi, and subsequent addition of AdCHO gave also ynone 12.



** Unwanted bromide **13** was found to slowly isomerise to bromoallene **7** on standing as a neat liquid, or this could be induced by the action of LiCuBr₂ in refluxing THF. As a representative experiment, after 20 h at reflux, allene **7** and bromide **13** were obtained as a 1.35:1 mixture respectively.

†† Temperature control was found to be essential to prevent hydride transfer to give products of the type RCOC≡CAD and RCH₂OH.

‡‡ Bromides **16** and **18** also isomerised on standing (tlc analysis) into their respective bromoallenes.

§§ The regioisomeric structures of bromoallenes **7** and **8** follow by the nature of the S_N2' reaction in which they are formed (ref. 13). These structures are further supported by characteristic NOE enhancements and HMBC correlations.‡

¶¶ At the outset, in a simplistic model, we predicted that epoxidation of a bromoallene would be electronically disfavoured at the Δ^{2,3} alkene position due to the deactivating donation of its π-cloud into the periplanar C–Br σ* orbital. However, it was unclear whether the mesomeric activating release of a bromine lone pair into the Δ^{1,2} alkene would overcome its inductive deactivating effect.

||| The bromoallene oxides **19–21** showed characteristic allene oxide vinyl resonances in their ¹H NMR spectra (CDCl₃) at 4.84, 4.97 and 4.83 ppm respectively allowing immediate identification of regiochemistry of epoxidation. Their respective ¹³C NMR resonances (CDCl₃) at C-3 (100.2, 100.3, 100.4 ppm) match perfectly with an allene oxide of the type ^tBuCH=C(–O)CHR (100.3 ppm, ref. 6). Characteristic IR stretches for all bromides **19–21** at 1806 cm^{–1} show them to be allene oxides (see ref. 1c).

*** We invoke the formation of bromodiketones **22–24** via further epoxidation of bromoallene oxides **19–21** to give their respective bromospirodi-epoxides, followed by ring-opening rearrangement to the diketones with bromide anion loss from C-1 and gain at C-3 (for the ring-opening rearrangement of spirodi-epoxides of trisubstituted allenes with halide anions see ref. 10b). They were assigned these structures on the basis of their characteristic diketone stretches in their IR spectra (22: 1713, 1694 cm^{–1}; 23: 1719, 1690 cm^{–1}; 24: 1717, 1692 cm^{–1}), the presence of two carbonyl resonances in their ¹³C NMR spectra (22: 206.3, 190.9 ppm; 23: 206.3, 191.0 ppm; 24: 205.9, 191.2 ppm), and a ¹³C NMR resonance at ca. 59 ppm (22: 58.8 ppm; 23: 58.2 ppm; 24: 59.1 ppm) that displays a bromine induced isotopic shift (for 22, Δδ = 1.6 ppb).‡ NOEs from H → R (22, NOESY; 23, 4.58%) confirm the bromide has shifted rather than a possible hydride shift (interchanging R and R'). However, their expected molecular ions with the characteristic bromine isotope pattern could not be observed by mass spectrometry, and only m/z 135 (Ad⁺) was observed by EI⁺ or CI⁺ methods (for 23, 24).

††† The yields represent the optimized yields for the bromoallene oxides using 3.0 equivalents of DMDO where the total mass recoveries from these experiments are >90%. The use of 1.5 equivalents gave the bromoallene oxides **19–21** in 21%, 19% and 21% isolated yield respectively, with (for **7** and **8**) 8% and 6% of **22** and **23**, and recovered starting materials **7** (68%) and **8** (74%) for >95% total mass recovery. The use of 5.0 equivalents (for **7** and **8**) led instead to the predominant formation of the corresponding bromodiketones **22** and **23** where no bromoallene oxides were observed.

‡‡‡ These results are also congruent with the reported epoxidations of tri(alkyl)substituted allenes (ref. 4 and 10b) where (i) the most substituted alkene is epoxidised, and (ii) the epoxidation occurs with face selectivity for the resulting E-olefin.

§§§ A search of the Cambridge Structural Database (version 5.37, Feb-2016 update) for an allene oxide moiety returned only 3 hits, all of which are epoxyC60fullerenes. The O–C bond lengths in the epoxide moiety of bromoallene oxide **19** show a marked asymmetry. Whilst that to the sp³ carbon [O1–C1 1.453(6) Å] is close to the average C–O bond length in epoxides with two sp³ carbon centres [1.446 Å], that to the sp² carbon of the adjacent C=C double bond is significantly shorter [O1–C2 1.388(7) Å]. The C–C bond of the epoxide moiety [C1–C2 1.435(8) Å] is also shorter than the average for epoxides with two sp³ carbon centres [1.467 Å]. The C2–C3 double bond [1.304(8) Å] is unchanged from the average seen in allene systems [1.317 Å] (data for average bond lengths of C=C double bonds in allene units, and in epoxides with two sp³ carbon centres came from statistical analyses of searches of the Cambridge Structural Database).

¶¶¶ The question of the preferred site of epoxidation in 1-bromo-3-alkyl allenes (ref. 11) remains experimentally unresolved however.

- For reviews see: (a) G. L'abbé, *Angew. Chem., Int. Ed. Engl.*, 1980, **19**, 276–289; (b) W. Smadja, *Chem. Rev.*, 1983, **83**, 263–320; (c) T. H. Chan and B. S. Ong, *Tetrahedron*, 1980, **36**, 2269–2289.
- For computational studies see: B. A. Hess, Jr., U. Eckart and J. Fabian, *J. Am. Chem. Soc.*, 1998, **120**, 12310–12315 and references cited therein.
- (a) R. L. Camp and F. D. Greene, *J. Am. Chem. Soc.*, 1968, **90**, 7349. A contemporaneous report on the peracid epoxidation of isomeric 1,1-di-*tert*-butylallene showed that 2,2-di-*tert*-butylcyclopropanone was formed instead: (b) J. K. Crandall and W. H. Machleder, *J. Am. Chem. Soc.*, 1968, **90**, 7347–7349. 1,1-Di-*tert*-butylallene oxide was later characterised in solution after low temperature epoxidation (ref. 4).
- 1,1,3-Tri-*tert*-butylallene oxide reported as a 10:1 mixture with an oxetane by glpc, and 1,3-di-*tert*-butyl-1-methylallene oxide as the major component of a four component mixture: J. K. Crandall, W. W. Conover, J. B. Komin and W. H. Machleder, *J. Org. Chem.*, 1974, **39**, 1723–1729. These trisubstituted allene oxides were reported to be resistant to conversion to their isomeric cyclopropanones.
- 1-*tert*-Butylallene oxide as a 'reasonably pure compound' by reduced pressure bulb-to-bulb distillation: (a) T. H. Chan and B. S. Ong, *J. Org. Chem.*, 1978, **53**, 2994–3001 and references cited therein. This allene oxide was reported to polymerise on standing in solution. This allene oxide has been prepared as a single enantiomer: (b) T. Konoike, T. Hayashi and Y. Araki, *Tetrahedron: Asymmetry*, 1994, **5**, 1559–1566.
- For a stable 3-*tert*-butylallene oxide from a fulvene *endo*-peroxide isolated by preparative tlc see: I. Erden, J. Drummond, R. Alstad and F. Xu, *Tetrahedron Lett.*, 1993, **34**, 1255–1258.
- Some non *tert*-butylallene oxides could be generated photochemically and detected by ¹H NMR methods in solution: (a) L. E. Breen, N. P. Schepp and C.-H. E. Tan, *Can. J. Chem.*, 2005, **83**, 1347–1351. See also: (b) M. D. Clay, J. Durber and N. P. Schepp, *Org. Lett.*, 2001, **3**, 3883–3886; (c) M. W. Konecny and N. P. Schepp, *Org. Biomol. Chem.*, 2009, **7**, 4437–4443 and references cited therein.
- For the isolation and characterization of a natural allene oxide as an unstable intermediate in the metabolism of lipid hydroperoxides see: (a) A. R. Brash, S. W. Baertschi, C. D. Ingram and T. M. Harris, *Proc. Natl. Acad. Sci. U. S. A.*, 1988, **85**, 3382–3386 and references cited therein. For *E* vs. *Z* isomers see: (b) A. R. Brash, W. E. Boeglin, D. F. Stec, M. Voehler, C. Schneider and J. K. Cha, *J. Biol. Chem.*, 2013, **288**, 20797–20806. See also: (c) N. V. Medvedeva, S. K. Latypov, A. A. Balandina, L. S. Mukhtarova and A. N. Grechkin, *Russ. J. Bioorg. Chem.*, 2005, **31**, 595–596.
- For representative use of allene oxides as intended transient intermediates see: (a) M. Shipman, H. R. Thorpe and I. R. Clemens, *Tetrahedron*, 1998, **54**, 14265–14282; (b) I. R. Clemens, M. Shipman and H. R. Thorpe, *Synlett*, 1995, 1065–1066; (c) S. J. Kim and J. K. Cha, *Tetrahedron Lett.*, 1988, **29**, 5613–5616 and references cited therein.
- For the first report of a stable spirodi-epoxide (from bisepoxidation of allenes) see: (a) J. K. Crandall, W. H. Machleder and M. J. Thomas, *J. Am. Chem. Soc.*, 1968, **90**, 7346–7347. For a recent representative use of spirodi-epoxides in cascade reactions see: (b) R. Sharma, M. Manpadi, Y. Zhang, H. Kim, N. G. Ahkmedov and L. J. Williams, *Org. Lett.*, 2011, **13**, 3352–3355.
- D. C. Braddock, J. Clarke and H. S. Rzepa, *Chem. Commun.*, 2013, **49**, 11176–11178.
- Ethyl 2-bromo-3-(diphenylmethylene) oxirane-2-carboxylate (*i.e.*, a bromoallene oxide ester) was reported in a study of ketenes and aliphatic diazo compounds: H. Staudinger and T. Reber, *Helv. Chim. Acta*, 1921, **4**, 3–23.
- E. J. Corey and N. W. Boaz, *Tetrahedron Lett.*, 1984, **25**, 3055–3058.
- R. Appel, *Angew. Chem., Int. Ed.*, 1975, **14**, 801–811.
- This useful building block has only been reported once previously (by a different method): J. Buendia, B. Darses and P. Dauban, *Angew. Chem., Int. Ed.*, 2015, **54**, 5697–5701.
- The CCl₄-PPh₃ combination on aldehydes typically gives rise to dichloromethane and dichloroalkene mixtures: R. Rabinowitz and R. Marcus, *J. Am. Chem. Soc.*, 1962, **84**, 1312–1313.
- There are surprisingly only limited literature precedents for the alkyl lithium mediated conversion of 1,1-dichloroalkenes into



- terminal alkynes. For a representative example see: F. Bertrand, B. Quiclet-Sire and S. Z. Zard, *Angew. Chem., Int. Ed.*, 1999, **38**, 1943–1946.
- 18 There is only one example in the literature that traps the presumed lithium acetylide from the action of alkyl lithiums on 1,1-dichloroalkenes with aldehydes as electrophiles: Z. Li and S. Z. Zard, *Org. Lett.*, 2009, **11**, 2868–2871.
- 19 R. W. Murray and M. Singh, *Org. Synth.*, 1997, **74**, 91.
- 20 A lithiated oxirane with considerable asymmetry has recently been characterised by crystallography: A. Salomone, F. M. Perna, A. Falcicchio, S. O. Nilsson Lill, A. Moliterni, R. Michel, S. Florio, D. Stalke and V. Capriati, *Chem. Sci.*, 2014, **5**, 528–538.
- 21 For all managed computational research data, including input and output files, see: D. C. Braddock, A. Mahtey, H. S. Rzepa and A. J. P. White, *Imperial College HPC data repository*, 2016. Collection doi: 10.14469/HPC/442.

