## ChemComm



**View Article Online** 

## COMMUNICATION



Cite this: Chem. Commun., 2016, 52, 12040

Received 17th August 2016, Accepted 12th September 2016

DOI: 10.1039/c6cc06761a

www.rsc.org/chemcomm

## $Br_2F_7^-$ and $Br_3F_{10}^-$ : peculiar anions showing $\mu_2$ - and $\mu_3$ -bridging F-atoms<sup>+</sup>

S. I. Ivlev,<sup>a</sup> A. J. Karttunen,\*<sup>b</sup> R. V. Ostvald\*<sup>c</sup> and F. Kraus\*<sup>a</sup>

RbCl and CsCl react with BrF<sub>3</sub> yielding the corresponding decafluoridotribromates(III), MBr<sub>3</sub>F<sub>10</sub> (M = Rb, Cs), which were structurally characterized for the first time. The Br<sub>3</sub>F<sub>10</sub><sup>-</sup> anion is surprisingly not linear but contains a  $\mu_3$ -bridging fluorine atom and seems to be the first example of  $\mu^3$ -F bridging of Br atoms. The compounds are highly reactive and cannot be handled in glassware. As for the tetrafluoridobromates themselves, they are powerful oxidizers and thus suitable for the dry-chemical recycling of precious metals and additionally feature a significantly higher BrF<sub>3</sub> content.

Under appropriate conditions fluorine is able to react with almost any element and a rich diversity of compounds result. In fluorine-containing molecules and molecular ions, the fluorine atom clearly prefers terminal positions, although many coordination compounds incorporate fluorine atoms as bridging ligands.<sup>1</sup> The most common type of F-bridging is the  $\mu_2$ -connection (either straight mono( $\mu$ -F), or bent di( $\mu$ -F)).<sup>2-13</sup> The highly strained tri( $\mu$ -F)-bridges between various elements<sup>1</sup> are scarce with only some examples in the literature.<sup>2-13</sup> Bridges among three atoms  $(\mu_3$ -F) are even more rare;<sup>14-21</sup> however,  $\mu_4$ -F<sup>22,23</sup> and even cage-like  $\mu_6$ -F<sup>22,24,25</sup> coordination types are known. In the vast majority of these compounds the fluorine atom bridges either metal atoms (thus, forming homoor hetero (oligo-)nuclear complexes), or, more rarely, metal and nonmetal atoms. In comparison, compounds featuring  $\mu_2$ -F bridges between two nonmetal atoms are even more uncommon; examples are known for H,<sup>26</sup> Kr,<sup>27,28</sup> Xe,<sup>29</sup> Br,<sup>4,30</sup> and I,<sup>31</sup> as well as for several metalloid atoms:  $B^{32}_{,33} As^{33}_{,33} Sb^{34}_{,34}$  and  $Si^{35}_{,35}$  To the best of our knowledge, the  $\mu_3$ -F type among nonmetals was fully established only for Xe<sup>36</sup> and I.<sup>20,21</sup>

Here, we report the results of our investigations on the non-common fluorine bridging in rubidium and cesium decafluoridotribromates(III): RbBr<sub>3</sub>F<sub>10</sub> (compound 1) and CsBr<sub>3</sub>F<sub>10</sub> (compound 2), respectively, as well as in rubidium heptafluoridodibromate( $\mathbf{m}$ ), RbBr<sub>2</sub>F<sub>7</sub> (compound 3). The compounds were synthesized during our research on the metal tetrafluoridobromate(III) series - powerful oxidizers which are promising for the dry-chemical recycling of noble metals.<sup>4,37,38</sup> MBr<sub>3</sub>F<sub>10</sub> and  $MBr_2F_7$  (M = Rb, Cs) were first reported by Stein;<sup>30</sup> however, their structures could only be deduced by Raman spectroscopy. Also, military personnel tried the synthesis of  $Br_3F_{10}^{-}$  but did not succeed.<sup>41</sup> The  $Br_3F_{10}^{-}$  (and  $Br_2F_7^{-}$ ) anion was reported to be chainlike with Br-µ-F-Br connections. We continued our previous works on BrF<sub>4</sub><sup>-</sup> compounds<sup>4</sup> and started to investigate compounds of higher BrF3 content to reveal how the chain elongation influences the structure of the anion and its reactivity. However, single crystal X-ray diffraction analyses, vibrational spectroscopy, as well as quantum chemical calculations showed that the Br<sub>3</sub>F<sub>10</sub><sup>-</sup> anions are not chain-like but contain  $\mu_3$ -bridging F atoms.

Compounds 1, 2 and 3 were synthesized using stoichiometric amounts of MCl (M = Rb, Cs) and BrF<sub>3</sub>, according to eqn (1), which was previously applied in the synthesis of RbBrF<sub>4</sub>:<sup>38</sup>

$$6MCl + (6n + 2)BrF_3 \rightarrow 6MBr_nF_{(3n+1)} + Br_2 + 3Cl_2$$
 (1)

 $RbBr_3F_{10}$  and  $CsBr_3F_{10}$  were obtained as yellowish-colored crystalline solids. Since both compounds are structurally isotypic, we present a detailed structural description only for compound 1 (Rb). Further details of the compounds presented here, such as powder X-ray patterns, Rietveld refinements, thermal analyses, experimentally observed as well as calculated Raman and IR spectra and band assignments, are available from the ESI.<sup>†</sup>

Compound 1 crystallizes in the monoclinic space group type  $P2_1$  (no. 4) with a = 7.6219(3) Å, b = 8.2593(4) Å, c = 8.4645(4) Å,

<sup>&</sup>lt;sup>a</sup> Anorganische Chemie, Philipps-Universität Marburg, Hans-Meerwein-Straße 4, 35032 Marburg, Germany. E-mail: florian.kraus@chemie.uni-marburg.de

<sup>&</sup>lt;sup>b</sup> Department of Chemistry, Aalto University, 00076 Aalto, Finland.

E-mail: antti.i.karttunen@aalto.fi

<sup>&</sup>lt;sup>c</sup> Department of Chemical Technology of Rare and Radioactive Metals, Tomsk Polytechnic University, 30 Lenina avenue, Tomsk 634050, Russia. E-mail: ostvald@tpu.ru

<sup>†</sup> Electronic supplementary information (ESI) available: Experimental and computational details and results from quantum chemical calculations. CCDC 1499499–1499501. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c6cc06761a



Fig. 1 The asymmetric unit of  $RbBr_3F_{10}.$  Displacement ellipsoids are shown at 70% probability level at 110 K.

 $\beta$  = 114.414(1)°, *V* = 485.21(4) Å<sup>3</sup>, and *Z* = 4 at 110 K. Further crystallographic details are given in Table 2. The asymmetric unit and the crystal packing of compound 1 are shown in Fig. 1 and Fig. S10 (ESI†), respectively.

An unexpected and unusual feature of compounds 1 and 2 is the shape of the  $Br_3F_{10}^-$  anion. It contains a  $\mu_3$ -bridging fluorine atom and not two  $\mu_2$ -bridging fluorine atoms as was previously deduced from Raman spectroscopic investigations on the compound.<sup>30</sup> To the best of our knowledge, RbBr<sub>3</sub>F<sub>10</sub> and CsBr<sub>3</sub>F<sub>10</sub> are the first documented compounds which incorporate fluorine as a  $\mu_3$ -bridging ligand between Br atoms.

The bromine atoms in RbBr<sub>3</sub>F<sub>10</sub> are coordinated by four fluorine atoms, each in an almost square-planar shape. The F–Br–F angles are observed in the range of 86.8(1) to 95.8(1)°, which is typical for known tetrafluoridobromates(III): KBrF<sub>4</sub>,<sup>12,39</sup> RbBrF<sub>4</sub>,<sup>38</sup> CsBrF<sub>4</sub>,<sup>4</sup> CsBr<sub>2</sub>F<sub>7</sub>,<sup>4</sup> and Ba(BrF<sub>4</sub>)<sub>2</sub>.<sup>37</sup> The Br atoms are located almost exactly in the virtual planes formed by the corresponding fluorine atoms (the distances from the least-squares planes are only 0.0026(4), 0.0033(5), and 0.0095(4) Å). The  $\mu_3$ -F atom is slightly (0.458(2) Å for compound 1, 0.370(8) Å for compound 2) above the virtual plane built by the three Br atoms. The Br– $\mu_3$ -F–Br angles are observed in the interval from 113.2(1) to 120.1(1)°. All three planar BrF<sub>3</sub>-units are tilted

towards each other, and the angles between the virtual planes are equal to 65.46(6), 66.22(7), and 69.10(7)°. The point group of the  $Br_3F_{10}^{-}$  anion is  $C_1$ ; however, its symmetry is very close to  $D_3$ . In DFT calculations on the isolated anion point group  $D_3$  is obtained. We also carried out two gas-phase DFT calculations on the  $Br_3F_{10}^{-}$  anion to compare the relative energetics of the  $\mu_3$ -bridging, experimentally observed structure and the previously postulated chain-like structure with Br-µ-F-Br connections. At the DFT-PBE0/def2-TZVP level of theory, the D<sub>3</sub>-symmetric,  $\mu_3$ -bridging structure is energetically 15 kJ mol<sup>-1</sup> more favorable than the  $C_{2h}$ -symmetric, chainlike structure (structural data are available in the ESI<sup>+</sup>). As expected, the Br-F bonds in the *trans*-position with respect to the  $\mu_3$ -F atom are shortened by circa 0.1 Å (for both compounds) in comparison to the other terminal F atoms. The coordination sphere of the Br<sub>3</sub>F<sub>10</sub><sup>-</sup> anion is also interesting and is discussed in the ESI<sup>+</sup> for reasons of brevity. Selected atomic distances and angles of the Br<sub>3</sub>F<sub>10</sub> anions, experimentally observed as well as theoretically predicted, are given in Table 1.

It is interesting to note that the cation type seems to have a rather low impact on the molecular structure of the  $Br_3F_{10}^{-1}$  anion. All corresponding Br–F bond lengths (including those to  $\mu_3$ -F as well as to *trans*-F atoms) do not change significantly ( $3\sigma$ ) if the Rb atoms are exchanged by Cs atoms. However, the bond angles seem to be more susceptible to the cation size and undergo a noticeable change resulting also in the change in the  $\mu_3$ -F distance from the virtual Br(1)–Br(2)–Br(3) plane.

Compound 3, rubidium heptafluoridodibromate(III), appears as a yellowish crystalline solid. However, the product of reaction (1) contains a mixture of RbBrF<sub>4</sub>, RbBr<sub>2</sub>F<sub>7</sub>, and BrF<sub>3</sub> rather than pure compound 3 (see the ESI†), which confirms the results obtained by Stein.<sup>30</sup> RbBr<sub>2</sub>F<sub>7</sub> crystallizes in the monoclinic space group type  $P2_1/c$  (no. 14) with a = 7.5109(3) Å, b = 7.8759(3) Å, c = 13.6898(5) Å,  $\beta = 123.118(2)^\circ$ , V = 678.26(5) Å<sup>3</sup>, and Z = 4 at 110 K. It is structurally isotypic to  $CsBr_2F_7$ .<sup>4</sup> The crystallographic details of RbBr<sub>2</sub>F<sub>7</sub> are given in Table 2. The asymmetric unit and the crystal packing of compound 3 are shown in Fig. 2 and Fig. S11 (ESI†) respectively.

 $RbBr_2F_7$  contains a  $Br_2F_7^-$  anion with the fluorine atom F(1) acting as a  $\mu_2$ -bridging ligand between the two bromine atoms. The  $Br-\mu$ -F distances are 2.115(2) and 2.145(2) Å, and are

Table 1	Selected bo	nd lengths	and angles f	or the Br <sub>3</sub> F <sub>1</sub>	$_{\rm D}^-$ anion.	The atom	labels	correspond	to those	in Fig. 1.	The [	DFT-PBE0	data	for th	e ideal
D <sub>3</sub> -symm	netric Br <sub>3</sub> F <sub>10</sub> <sup>-</sup>	anion have	e been calcul	ated for the	gas-phas	e (see the	ESI for	computatio	nal details	s)					

	Value (Å/°)						
Parameter	RbBr <sub>3</sub> F <sub>10</sub>	$CsBr_3F_{10}$	$Br_3F_{10}^{-}$ (DFT-PBE0)				
Br(1)-µ <sub>3</sub> -F	2.243(3)	2.238(10)	2.30				
$Br(2)-\mu_3$ -F	2.248(2)	2.246(9)	2.30				
$Br(3)-\mu_3$ -F	2.320(3)	2.329(10)	2.30				
Br-trans-F (F7, F8, F10)	$1.745(2) \cdots 1.752(2)$	$1.746(6) \cdot \cdot \cdot 1.767(7)$	1.76				
Br–F (other)	$1.837(2) \cdots 1.874(2)$	$1.824(10) \cdots 1.878(7)$	1.84				
$Br(1)-\mu_3$ -F- $Br(2)$	120.1(1)	122.0(3)	120				
$Br(2) - \mu_3 - F - Br(3)$	114.7(1)	115.0(4)	120				
$Br(3)-\mu_3$ -F-Br(1)	113.2(1)	115.1(4)	120				
Distance from $\mu_3$ -F to Br(1)-Br(2)-Br(3) plane	0.458(2)	0.370(8)	0				
Tilting angles between $F_3M-\mu$ -F-MF <sub>3</sub> least-squares planes	65.46(6)	62.1(3)	81.8				
	66.22(7)	66.2(2)					
	69.10(7)	72.0(2)					

	Compound 1	Compound 2	Compound 3
Empirical formula	Br <sub>3</sub> F <sub>10</sub> Rb	Br <sub>3</sub> F <sub>10</sub> Cs	Br <sub>2</sub> F <sub>7</sub> Rb
Color and appearance	Colorless plates, yellow	vish powders	
$M/g \text{ mol}^{-1}$	515.20	562.60	378.29
Crystal system	Monoclinic		
Space group	$P2_1$ (no. 4)		$P2_{1}/c$ (no. 14)
a/Å	7.6219(3)	7.4399(16)	7.5109(3)
<i>b</i> /Å	8.2593(4)	8.4037(17)	7.8759(3)
c/Å	8.4645(4)	8.8365(18)	13.6898(5)
$\beta$ / $^{\circ}$	114.414(1)	115.31(3)	123.118(2)
$V/\text{\AA}^3$	485.21(4)	519.4(2)	678.26(5)
Ζ	2		4
$ ho_{ m calc}/ m g\  m cm^{-3}$	3.526	3.597	3.705
$ ho_{\rm exp}/{\rm g~cm}^{-3}$	$3.687\pm0.006$	$3.460\pm0.026$	$3.786\pm0.007$
$\lambda/\text{\AA}$	0.71073 (Mo-K <sub>α</sub> )		
T/K	110	100	110
$R_{\rm int}, R_{\sigma}$	0.0444, 0.0465	0.0472, 0.1449	0.0533, 0.0309
$R(F^2)$ (all data), $wR(F^2)$ (all data)	0.0294, 0.0509	0.0730, 0.0732	0.0336, 0.0507
S (all data)	1.024	0.790	1.055
Flack parameter x	0.050(6)	0.00(2)	_
No. of reflections, parameters, constraints, restraints	4132, 128, 0, 1	3195, 128, 0, 1	2035, 92, 0, 0
$2\theta$ range refined (min, max)	2.643, 35.843	2.912, 31.900	3.138, 30.362
$\Delta \rho_{\rm max}$ , $\Delta \rho_{\rm min}/{\rm e}$ Å <sup>-3</sup>	1.22, -1.45	1.43, -1.21	0.94, -0.94
$(\Delta \sigma)_{\rm max}$	0.001	0.000	0.001
ICSD number	431741	431740	431739



essentially equal to 2.113(1) and 2.143(1) Å as observed in  $CsBr_2F_7$ ;<sup>4</sup> however, the Br–µ-F–Br angle is 134.75(9)° being *circa* 6° smaller than the analogous angle of 140.27(6)° reported for the Cs compound. The *trans*-F–Br distances are equal to 1.767(2) and 1.780(2) Å, while the other terminal Br–F bond lengths lie in the range of 1.845(2) to 1.890(2) Å, therefore, being equal to the corresponding bonds in  $CsBr_2F_7$  within the  $3\sigma$  criterion. The F–Br–F angles are observed within the interval from 87.46(7) to 95.65(7)° (87.74(6) to 95.12(5)° in  $CsBr_2F_7$ ). These facts imply again that these counter-ions do not significantly influence the molecular structure of the fluoridobromate anions. The cell parameters together with the selected bond lengths and angles for  $RbBr_2F_7$  and  $CsBr_2F_7$  as well as for

Fig. 2 The asymmetric unit of  $RbBr_2F_7.$  Displacement ellipsoids are shown at 70% probability level at 110 K.

Table 3Selected bond lengths and angles in  $RbBr_2F_7$ ,  $CsBr_2F_7$ , and  $CsAu_2F_7$ . The atom labels correspond to those in Fig. 2, and were changed in thecases of  $CsBr_2F_7$  and  $CsAu_2F_7$  to make the comparison possible

	Value (Å∕°)	Value (Å/°)					
Parameter	$\text{RbBr}_2\text{F}_7(P2_1/c)$	$\mathrm{CsBr}_{2}\mathrm{F}_{7}\left(P2_{1}/c\right)^{4}$	$\mathrm{CsAu}_{2}\mathrm{F}_{7}\left(C2/c\right)^{40}$				
a/Å	7.5109(3)	7.7078(1)	11.365(6)				
b/Å	7.8759(3)	8.0218(2)	10.820(15)				
c/Å	13.6898(5)	14.1584(3)	7.374(3)				
$\beta/^{\circ}$	123.118(2)	122.742(2)	123.40(3)				
M(1)-F(1)	2.145(2)	2.143(1)	1.988(8)				
M(1) - F(3)	1.871(2)	1.868(2)	1.89(2)				
M(1) - F(4)	1.855(2)	1.858(2)	1.92(2)				
M(1) - F(5)	1.767(2)	1.769(1)	1.86(2)				
M(2) - F(1)	2.115(2)	2.112(2)	1.988(8)				
M(2) - F(2)	1.890(2)	1.884(2)	1.89(2)				
M(2) - F(6)	1.845(2)	1.849(1)	1.92(2)				
M(2) - F(7)	1.780(2)	1.779(2)	1.86(2)				
M(1) - F(1) - M(2)	134.75(9)	140.27(6)	130.08(4)				
F(5) - M(1) - F(1)	175.17(8)	176.18(6)	176.0(6)				
F(7) - M(2) - F(1)	176.04(8)	176.44(7)	176.0(6)				
Tilting angle between $F_3M-\mu$ -F-MF <sub>3</sub> planes	65.31(6)	61.90(4)	48.1(3)				

 $M = Br (RbBr_2F_7, CsBr_2F_7); Au (CsAu_2F_7).$ 

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.

Article. Published on 19 September 2016. Downloaded on 8/17/2025 8:03:29 AM

**Den Access** 

 $CsAu_2F_7$  (which is similar in terms of the anion geometry but is not structurally isotypic)<sup>40</sup> are shown in Table 3.

In comparison to the Br<sub>3</sub>F<sub>10</sub><sup>-</sup> anions, a shortening of the Br- $\mu$ -F-Br bridges of *circa* 0.1 to 0.2 Å is observed which may be due to smaller Coulomb repulsion between the two Br atoms. The Br-*trans*-F distances in the Br<sub>2</sub>F<sub>7</sub><sup>-</sup> anion are on average 0.025 Å longer than those in Br<sub>3</sub>F<sub>10</sub><sup>-</sup>. The other terminal fluorine atoms are not so susceptible to the change of the inner environment and the corresponding Br-F bonds show only a slight elongation of approximately 0.01 Å.

All compounds reported in this work may be convenient carriers for BrF<sub>3</sub> due to its high mass content (72.4, 79.7, 73.0% by mass in RbBr<sub>2</sub>F<sub>7</sub>, RbBr<sub>3</sub>F<sub>10</sub> and CsBr<sub>3</sub>F<sub>10</sub> respectively). So, the "problematic" BrF<sub>3</sub> can be replaced by these comparatively easy to handle solid compounds. To date there has been no indication for the existence of such interhalide anions in compounds with cations other than Rb or Cs, which is possibly due to the cation size. Also, it is unknown if suitable reaction conditions can be found that would lead to the formation of the highly symmetrical Br<sub>4</sub>F<sub>13</sub><sup>-</sup> anion, with a  $\mu_4$ -bridging F atom. Investigations in those directions are ongoing.

## Notes and references

- 1 T. Birk, ChemistryOpen, 2013, 2, 13-16.
- 2 C. Bour, J. Monot, S. Tang, R. Guillot, J. Farjon and V. Gandon, Organometallics, 2014, 33, 594–599.
- 3 M. Inomata and Y. Suenaga, Acta Crystallogr., Sect. E: Struct. Rep. Online, 2014, 70, m359–360.
- 4 S. Ivlev, P. Woidy, V. Sobolev, I. Gerin, R. Ostvald and F. Kraus, Z. Anorg. Allg. Chem., 2013, 639, 2846–2850.
- 5 S. P. Sarish, H. W. Roesky, M. John, A. Ringe and J. Magull, *Chem. Commun.*, 2009, 2390–2392.
- 6 E. Tomat, L. Cuesta, V. M. Lynch and J. L. Sessler, *Inorg. Chem.*, 2007, **46**, 6224–6226.
- 7 G. B. Nikiforov, C. Knapp, J. Passmore and A. Decken, J. Fluorine Chem., 2006, 127, 1398–1404.
- 8 H. Hao, C. Cui, H. W. Roesky, G. Bai, H.-G. Schmidt and M. Noltemeyer, *Chem. Commun.*, 2001, 1118–1119.
- 9 J. Gindl, M. A. Said, P. Yu, H. W. Roesky, M. Noltemeyer and H.-G. Schmidt, *Isr. J. Chem.*, 1999, **39**, 125–128.
- 10 D. M. Dawson, R. A. Henderson, A. Hills and D. L. Hughes, J. Chem. Soc., Dalton Trans., 1992, 973–976.
- 11 R. H. Crabtree, G. G. Hlatky and E. M. Holt, *J. Am. Chem. Soc.*, 1983, **105**, 7302–7306.
- 12 A. J. Edwards and G. R. Jones, J. Chem. Soc. A, 1970, 1491-1497.

- 13 H. Schmidbaur and H. F. Klein, Angew. Chem., 1966, 78, 750.
- 14 M. V. Ivanova, T. Köchner, H. P. A. Mercier and G. J. Schrobilgen, *Inorg. Chem.*, 2013, 52, 6806–6819.
- 15 D. S. Tereshchenko, I. V. Morozov, A. I. Boltalin, E. V. Karpova, T. Y. Glazunova and S. I. Troyanov, *Crystallogr. Rep.*, 2013, 58, 68–77.
- 16 E.-Q. Gao, N. Liu, A.-L. Cheng and S. Gao, Chem. Commun., 2007, 2470-2472.
- 17 W. Uhl, F. Breher, B. Neumüller, A. Lützen, W. Saak and J. Grunenberg, *Organometallics*, 2001, **20**, 5478–5484.
- 18 B. F. Straub, F. Rominger and P. Hofmann, *Inorg. Chem.*, 2000, **39**, 2113–2119.
- 19 S. Herold and S. J. Lippard, Inorg. Chem., 1997, 36, 50-58.
- 20 A. R. Mahjoub, A. Hoser, J. Fuchs and K. Seppelt, Angew. Chem., Int. Ed., 1989, 28, 1526–1527.
- 21 A. R. Mahjoub, A. Hoser, J. Fuchs and K. Seppelt, *Angew. Chem.*, 1989, **101**, 1528–1529.
- 22 H. W. Roesky and I. Haiduc, J. Chem. Soc., Dalton Trans., 1999, 2249–2264.
- 23 N. Buchholz and R. Mattes, Angew. Chem., Int. Ed., 1986, 25, 1104–1105.
- 24 Q.-M. Wang and T. C. W. Mak, Chem. Commun., 2000, 1435-1436.
- 25 D. Stalke, F.-Q. Liu and H. W. Roesky, *Polyhedron*, 1996, 15, 2841–2843.
- 26 D. Mootz and K. Bartmann, Angew. Chem., Int. Ed., 1988, 27, 391-392.
- 27 R. J. Gillespie and G. J. Schrobilgen, Inorg. Chem., 1976, 15, 22-31.
- 28 J. F. Lehmann, D. A. Dixon and G. J. Schrobilgen, *Inorg. Chem.*, 2001, 40, 3002–3017.
- 29 B. A. Fir, M. Gerken, B. E. Pointner, H. P. A. Mercier, D. A. Dixon and G. J. Schrobilgen, *J. Fluorine Chem.*, 2000, **105**, 159–167.
- 30 L. Stein, J. Fluorine Chem., 1985, 27, 249-256.
- 31 K. O. Christe, in Book of Abstracts, 249th ACS National Meeting, American Chemical Society, Denver, Colorado, 2015, p. INOR 362.
- 32 T. W. Hudnall, C.-W. Chiu and F. P. Gabbaï, Acc. Chem. Res., 2009, 42, 388–397.
- 33 K. O. Christe, X. Zhang, R. Bau, J. Hegge, G. A. Olah, G. K. S. Prakash and J. A. Sheehy, *J. Am. Chem. Soc.*, 2000, **122**, 481–487.
- 34 J. F. Lehmann, G. J. Schrobilgen, K. O. Christe, A. Kornath and R. J. Suontamo, *Inorg. Chem.*, 2004, 43, 6905–6921.
- 35 M. Lehmann, A. Schulz and A. Villinger, Angew. Chem., Int. Ed., 2009, 48, 7444–7447.
- 36 J. H. Holloway, V. Kaucic, D. Martin-Rovet, D. R. Russell, G. J. Schrobilgen and H. Selig, *Inorg. Chem.*, 1985, 24, 678–683.
- 37 S. Ivlev, V. Sobolev, M. Hoelzel, A. J. Karttunen, T. Müller, I. Gerin, R. Ostvald and F. Kraus, *Eur. J. Inorg. Chem.*, 2014, 6261–6267.
- 38 S. Ivlev, A. Karttunen, R. Ostvald and F. Kraus, Z. Anorg. Allg. Chem., 2015, 641, 2593–2598.
- 39 W. G. Sly and R. E. Marsh, Acta Crystallogr., 1957, 10, 378-379.
- 40 R. Schmidt and B. G. Müller, Z. Anorg. Allg. Chem., 2004, 630, 2393–2397.
- 41 T. Groshens, R. Hollins, Organic Perfluorohalogenate Salts; New Energetic Materials, Technical Report, DTRA-TR-14-39, June 2014, www.dtic.mil/cgi-bin/GetTRDoc?AD=ADA604532.