 Cite this: *Chem. Commun.*, 2016, 52, 9972

Correction: Beyond "turn-on" readout: from zero background to signal amplification by combination of magnetic separation and plasmon enhanced fluorescence

Suqin Gong and Yunsheng Xia*

DOI: 10.1039/c6cc90336c

www.rsc.org/chemcomm

Correction for 'Beyond "turn-on" readout: from zero background to signal amplification by combination of magnetic separation and plasmon enhanced fluorescence' by Suqin Gong and Yunsheng Xia, *Chem. Commun.*, 2016, **52**, 9660–9663.

The authors regret that axis labels and units were omitted from the graphs in Fig. 2C and F in the original article, and the labels 'C' and 'F' are also missing from the respective parts of the image. A corrected version of Fig. 2 is presented herein.

Fig. 2 TEM (A) and HRTEM (B) images of the hybrid (AgNP-QD) superstructures. (C) DLS sizes of the Ag NPs (top) and the hybrid AgNP-QD superstructures (bottom). Fluorescence emission spectra for assaying moderate concentrations of HAase (0–0.15 ng mL⁻¹) in the presence (D) and absence (E) of the Ag NPs. (F) Plots of the fluorescence intensities versus HAase concentrations in the presence and absence of the Ag NPs. (G) Plots of the scattering intensities (at 550 nm) versus HAase concentrations in the presence and absence of Ag NPs. Fluorescence emission spectra for assaying ultra-trace concentrations of HAase (0–7.0 pg mL⁻¹) in the presence (H) and absence (I) of the Ag NPs.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

Key Laboratory of Functional Molecular Solids, Ministry of Education, College of Chemistry and Materials Science, Anhui Normal University, Wuhu, 241000, China.
E-mail: xiayuns@mail.ahnu.edu.cn